

## **Archive ouverte UNIGE**

https://archive-ouverte.unige.ch

Article scientifique

Article 2022

Published version

**Open Access** 

This is the published version of the publication, made available in accordance with the publisher's policy.

\_ \_ \_

# The influence of small-sided football games with numerical variability in external training load

Sousa, Honorato; Gouveia, Élvio Rúbio; Marques, Adilson; Sarmento, Hugo; Pestana, Marcelo; Quintal, Tomás; Lopes, Helder; Ihle, Andreas

### How to cite

SOUSA, Honorato et al. The influence of small-sided football games with numerical variability in external training load. In: Sustainability, 2022, vol. 14, n° 2, p. 1000. doi: 10.3390/su14021000

This publication URL:	https://archive-ouverte.unige.ch/unige:165400
Publication DOI:	<u>10.3390/su14021000</u>

© The author(s). This work is licensed under a Creative Commons Attribution (CC BY) <u>https://creativecommons.org/licenses/by/4.0</u>



Article



# The Influence of Small-Sided Football Games with Numerical Variability in External Training Load

Honorato Sousa <sup>1</sup>, Élvio Rúbio Gouveia <sup>2,3,4,\*</sup>, Adilson Marques <sup>5,6</sup>, Hugo Sarmento <sup>1</sup>, Marcelo Pestana <sup>2</sup>, Tomás Quintal <sup>2</sup>, Helder Lopes <sup>2,7</sup> and Andreas Ihle <sup>4,8,9</sup>

- <sup>1</sup> University of Coimbra, Research Unit for Sport and Physical Activity (CIDAF), Faculty of Sport Sciences and Physical Education, 3004-504 Coimbra, Portugal; honoratosousa@hotmail.com (H.S.); hugo.sarmento@uc.pt (H.S.)
- <sup>2</sup> Department of Physical Education and Sport, University of Madeira, 9000-072 Funchal, Portugal; jm.desporto@gmail.com (M.P.); tomasquintal9@gmail.com (T.Q.); hlopes@staff.uma.pt (H.L.)
- Interactive Technologies Institute, LARSYS, 9020-105 Funchal, Portugal
- <sup>4</sup> Center for the Interdisciplinary Study of Gerontology and Vulnerability, University of Geneva, 1211 Geneva, Switzerland; andreas.ihle@unige.ch
- <sup>5</sup> The Interdisciplinary Centre for the Study of Human Performance (CIPER), Faculty of Human Kinetics, University of Lisbon, 1649-004 Lisbon, Portugal; amarques@fmh.ulisboa.pt
- <sup>6</sup> Instituto de Saúde Ambiental (ISAMB), University of Lisbon, 1649-026 Lisbon, Portugal
- <sup>7</sup> CiTUR-Centro de Investigação, Desenvolvimento e Inovação em Turismo, 9000-082 Funchal, Portugal
- <sup>8</sup> Department of Psychology, University of Geneva, 1211 Geneva, Switzerland <sup>9</sup> Swige National Contra of Commutance in Paccarsh LW//S Overcoming Values
- Swiss National Centre of Competence in Research LIVES–Overcoming Vulnerability: Life Course Perspectives, 1015 Lausanne, Switzerland
- \* Correspondence: erubiog@staff.uma.pt; Tel.:+351-291705249

**Abstract:** Small-sided games (SSGs) are a motivational strategy and effective training method to develop skills and physical fitness at optimal intensity. This study investigates the influence of variability in the number of players on the physiological response using SSGs. The sample was composed of 10 field players with an average age of  $15.9 \pm 0.50$  years old. Three sessions were held on 3 different days to collect information. The 10 min SSG exercise with temporal numerical variability was repeated twice on each day. The numerical ratio of players involved changed every 2 min without disrupting the practice. In a game space with  $35 \times 25$  m, the exercise always started in a situation of 3 vs. 3. External training load metrics (i.e., physical activity counts, activity intensity, and energy expenditure) were assessed using Actigraph WGT3X accelerometers. The 3 vs. 3 format game had higher physical activity intensity. Consequently, the 3 vs. 3 format in young football players had higher energy expenditure (i.e., a higher METs) than that in the other studied game formats. This study confirms the influence of SSGs with numerical variability in external training load.

Keywords: physiological response; external load; small-sided games; numerical variability

#### 1. Introduction

Small-sided games (SSGs), also referred to as skill-based conditioned games [1], gamebased training [2], or reduced and conditioned games [3], are game situations that generate practice conditions to induce specific game behaviours. SSGs are a sustainable training method to develop skills and physical fitness at optimal intensity.

These training methods are based on the variation of practice conditions (depending on training objective), namely, variability in the number of players and game space, and the adaptation or conditioning of the game regulation (among others), but maintaining the logic and reality of the basic principles of the game [4-6].

One of the foci of most significant interest in the use of SSGs in the training process is based on their use to develop the physical dimension of the practitioner's abilities,



Citation: Sousa, H.; Gouveia, É.R.; Marques, A.; Sarmento, H.; Pestana, M.; Quintal, T.; Lopes, H.; Ihle, A. The Influence of Small-Sided Football Games with Numerical Variability in External Training Load. *Sustainability* **2022**, *14*, 1000. https://doi.org/ 10.3390/su14021000

Academic Editor: José Carmelo Adsuar Sala

Received: 13 November 2021 Accepted: 14 January 2022 Published: 17 January 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). particularly in situations of pressure and fatigue [7,8]. SSGs are capable of generating highintensity activities (short sprint, constant tackles, greater proximity to the opponent, and consequently an increase in the number of duels), involving various types of movements, similar to those that exist in the real competitive context [9].

Although the investigation of SSGs arose in the last decade, this training method has been used since the 1970s. However, in the last decade, interest in this type of situation has grown again, intending to increase the participation of players in training, and consequently the physical and physiological load [10-15].

The growing popularity of the SSGs comes from a set of benefits that can be used in football at all ages and levels, which replicate the required movements, physiological intensity, and technical resources of a competitive game [9,10,16,17] while triggering decision making about pressure and fatigue in players [7].

It is possible to measure exercise intensity in SSGs through player movement, and the physiological responses and perception of effort [18]. The coach controls a set of variables that influence exercise intensity in SSGs [19], such as field area, the number of players, coach feedback, training planning (continuous or interval and rest-time manipulation), modification of the rules, and the use of goals and goalkeepers (GK) [9,20–22].

Although there are studies that address the issue of numerical variability in SSGs, they were mostly conducted through the use of jokers [10,23,24], which is momentary variability. The joker is often used by teachers or coaches to manipulate the SGG in terms of numerical relationship, which can be defined as supporting both teams in the offensive or defensive process of the game [24,25]. The dynamics that a joker provides to training situations never happen in a real game context, as this element always plays in the team that has the ball, and there is no reversibility of functions, which is precisely the opposite of what the game is and what this study intends to present.

Physiological analysis of SSGs in football training has been around for over 30 years [26]. More recently, with technological advances and greater access to a wide range of instruments (accelerometers, GPS, heart-rate monitoring), it has led to load assessment and monitoring being more accessible, reliable, and accurate [7,27–30].

Some studies indicate that SSG formats with different numbers of players induce different physiological and perceptual characteristics [31–33]. However, these studies only verified the influence of the numerical change in team players while always maintaining numerical equality between teams. Given that football is currently played in the search (attack) and impediment (defence) of constant numerical imbalances, further research is needed to clarify the impact of team responses on numerical imbalances using SSGs [34,35]. To the best of our knowledge, few studies considered unbalanced formats or a changeable numerical ratio of players in the SSGs on the external load of practitioners [23,34,36,37]. Therefore, this study investigates the influence of variability in the number of players using reduced game formats on the physiological response. We identify if there is a change in the physiological response of players in five different SSG formats with numerical variability (3 vs. 3, 4 vs. 3, 5 vs. 3, 3 vs. 4, and 3 vs. 5).

#### 2. Materials and Methods

#### 2.1. Participants

The sample was composed of 10 male field players with an average age of  $15.9 \pm 0.50$  years old. Ten field players were twice examined on 3 different assessment days. In total, 24 observations were considered in the analyses. All participants were affiliated with the local football association in the last 5 seasons and belonged to the same team. They had a regular weekly program of 3 training sessions of 75 min each and the competitive moment at the end of the week (regional or local championship that qualifies a team for the final national championship phase). The study sample size was determined following a priori power analysis, which indicated that, to detect a large effect size of r = 0.40 with an alpha probability of 0.05, a power of 0.85, and a total of 5 measurements, the sample size

would need to comprise 10 participants. GPower, (Heinrich Heine University, Düsseldorf, Germany; 3.1.9.7 software) was used in the calculations [38]

All players were informed about the research objectives, its requirements, and the potential benefits and risks. Participation was voluntary. Parents or legal guardians signed informed consent, and players gave their verbal assent. All procedures followed the guidelines stated in the Declaration of Helsinki [39] and had been approved by the ethics commission: Ethical Committee of the Faculty of Human Kinetics, University of Lisbon, approval code: CEIFMH No: 35/2021; approval date: 6 July 2021.

#### 2.2. Study Design and Procedures

Three sessions were held for the data collection on three different days, starting at the same time and following a 24 h interval between each of the collection sessions. The presented SSG format was variable, with a total duration of 10 min and changing the numerical ratio of players in the game every 2 min without interrupting the practice. The exercise always started as a 3 vs. 3 + GK game format (0 to 2 min), then 4 vs. 3 + GK (2 to 4 min), 5 vs. 3 + GK (4 to 6 min), 3 vs. 4 + GK (6 to 8 min), and 3 vs. 5 + GK (8 to 10 min).

The order in which these changes took place was randomised differently for the three days of the study (except for the initial 3 vs. 3 + GK format). The purpose was to keep uncertainty about the sequence of the situations to ensure that there were no strategic behaviours on the part of the participants who played the format. Every 2 min, the team structure was reconfigured, entering players in a delimited zone (in the middle of the game space) or leaving the nearest line where they were. However, the original three elements of each team that started the 3 vs. 3 + GK remained throughout the exercise. The new elements that entered always played at the same time each day. The entire 10 min exercise were repeated twice. Teams remained the same throughout the study, and the prior knowledge of the coach was used for the composition and structuring of the teams on the basis of the capacities and usually performed positions or functions for an equitable distribution of individual value, adopting similar strategies and following the recommendations [40].

The sessions started with a 20 min warm-up, with a general mobilisation of all body segments for a musculoskeletal activation, followed by an exercise to maintain ball possession without goals but using GKs as external supports to the game space. These two exercises were later complemented with dynamic stretching and speed exercises. After the warm-up, the used SSG formats lasted 10 min, followed by 5 min of passive rest to perform a second series of 10 min. The sessions ended with a 10 min cooldown using a series of static stretches.

The dimensions of the game space were  $35 \times 25$  m using smaller football goals that were 6 m wide and 2 m high. The only changes to regulation were removing the offside rule and replacing the ball when it left the end line, always starting by being carried out by the GK of the team to which the ball belonged, regardless of whether it was a corner. In these replacements, the GKs had the freedom to use hands or feet. To keep play intensity high and increase the effective playing time, several balls were placed around the field, and several collaborators in the study were strategically positioned to collect and give the ball whenever it left the game space. The teachers or coaches and those responsible for the study were only authorised to provide verbal encouragement feedback. They were expressly prohibited from giving any information or indications on strategic issues or tactical-technical behaviour.

#### 2.3. External Training Load Metrics

External training loads were assessed using the ActiGraph wGT3X-BT accelerometer (ActiGraph, LLC, Pensacola, FL, USA). Participants wore the accelerometer over the right anterior superior iliac spine. The accelerometer was initialised to store activity counts at the beginning of each 10 min exercise. The captured raw acceleration data were converted into various objective activities using ActLife Monitoring System software. ActiLife5's cut point data scoring option allows for users to break out the proportion of time that each

dataset encompasses into different categories of activity. Data were processed using a 10 s epoch. Activity counts were analysed by the vector magnitude counts and determined physical activity intensity in the following categories: (1) light physical activity intensity (760–1952 counts), (2) moderate physical activity intensity (1953–5724 counts), (3) vigorous physical activity intensity (5725–9498 counts), and (4) very vigorous physical activity intensity (9499 and higher counts). Energy expenditure was also analysed using MET rates.

#### 2.4. Statistical Analyses

First, the Friedman test was used to investigate the change in young football players' internal and external training load across five different small-sided game formats with numerical variability (i.e., 3 vs. 3, 4 vs. 3, 5 vs. 3, 3 vs. 4, and 3 vs. 5). Second, to investigate the change in internal and external training load between each game format (i.e., 3 vs. 3 vs. 4 vs. 3, 3 vs. 3, 3 vs. 5, 4 vs. 3 vs. 5 vs. 3, 4 vs. 3 vs. 3 vs. 4, 5 vs. 3 vs. 5, 4 vs. 3 vs. 5) we used the Wilcoxon signed-rank test. Data analysis assumptions were verified. The level of confidence was set at 95%. Data were analysed using IBM SPSS statistics 26.

#### 3. Results

Participants' characteristics are summarised in Table 1.

Table 1. Participants' characteristics.

Physical Tests	Mean	Std. Deviation	95% Confidence Interval for Mean			
Age (years)	15.9	0.50	15.6–16.3			
Weight (kg)	60.5	7.60	55.7-65.3			
Height (cm)	171.5	6.57	167.3–175.7			
BMI (kg/m <sup>2</sup> )	20.5	2.12	19.2–21.9			
Bar suspension (s)	32.8	11.64	25.4-40.2			
Sit-ups (n)	26.7	1.72	25.6–27.8			
Sit and Reach Bilateral (cm)	33.1	7.59	28.3–37.9			
Dynamometry Right hand (kg)	35.6	4.12	33.0–3.2			
Flamingo Balance (n)	11.9	4.06	9.3–14.5			
Sprint 5 m (s)	1.0	0.06	0.97–1.0			
Sprint 10 m (s)	1.7	0.06	1.7–1.8			
Sprint 35 m (s)	5.0	0.16	4.9–5.1			
YYIR1 (s)	1151.7	462.21	858.0-1445.3			
T Test (s)	9.9	0.52	9.6–10.2			
Counter Movement Jump (cm)	28.6	2.8	26.8-30.4			
Squat Jump (cm)	28.9	2.5	27.3–30.5			

Table 2 depicts the results of the Friedman test, indicating that, in the external training load metrics, there was a statistically significant difference in the vector magnitude counts,  $(\chi^2 (4, n = 72) = 79.12, p < 0.001)$ , steps counts  $(\chi^2 (4, n = 72) = 10.88, p < 0.001)$ , light physical activity intensity counts  $(\chi^2 (4, n = 72) = 20.26, p < 0.001)$ , moderate physical activity intensity counts  $(\chi^2 (4, n = 72) = 69.79, p < 0.001)$ , vigorous physical activity intensity counts  $(\chi^2 (4, n = 72) = 9.88, p = 0.043)$ , very vigorous physical activity intensity counts  $(\chi^2 (4, n = 72) = 9.88, p = 0.043)$ , very vigorous physical activity intensity counts  $(\chi^2 (4, n = 72) = 72.44, p < 0.001)$ , and METs  $(\chi^2 (4, n = 72 = 109.86, p < 0.001)$ , across the different studied game formats.

Table 2. External training load metrics.												
External Load Metrics	3 vs. 3 (1) ( <i>n</i> = 24)		4 vs. 3 (2) ( <i>n</i> = 24)		5 vs. 3 (3) ( <i>n</i> = 24)		3 vs. 4 (4) ( <i>n</i> = 24)		3 vs. 5 (5) ( <i>n</i> = 24)		+ (p)	‡
	$\mathbf{M} \pm \mathbf{Std.}$	Med.	$\mathbf{M} \pm \mathbf{Std.}$	Med.	$\mathbf{M} \pm \mathbf{Std.}$	Med.	$\mathbf{M} \pm \mathbf{Std.}$	Med.	$\mathbf{M} \pm \mathbf{Std.}$	Med.		
Vector magnitude	18,144.18 ± 3018.63	18,534.60	16,109.92 ± 2259.99	16095.10	16,069.22 ± 2189.49	16,141.55	15,486.84 ± 2887.96	15,109.35	16,658.36 ± 2102.17	16,616.40	<0.001	1 > 2, 3, 4 and 5; 2 < 5; 4 < 5
Steps (counts)	$244.00\pm17.51$	244.00	$227.31\pm22.23$	230.00	$227.04\pm22.43$	229.00	$215.04\pm26.96$	220.00	$229.14\pm24.18$	233.00	< 0.001	1 > 2, 3, 4 and 5; $2 > 4;$ 3 > 4; 4 < 5
SPA counts	$0,00\pm0,00$	0.00	$0.005\pm0.04$	0.00	$0.00\pm0.00$	0.00	$0.01\pm0.05$	0.00	$0.002\pm0.02$	0.00	0.046	n.s.
LPA counts	$0.00\pm0.00$	0.00	$0.02\pm0.06$	0.00	$0.00\pm0.00$	0.00	$0.02\pm0.07$	0.00	$0.002\pm0.02$	0.00	< 0.001	1 < 2 and 4; 2 > 3 and 5; 3 < 4; 4 > 5
MPA counts	$0.08\pm0.14$	0.00	$0.33\pm0.30$	0.33	$0.31\pm0.31$	0.17	$0.42\pm0.37$	0.34	$0.27\pm0.35$	0.17	< 0.001	1 < 2, 3, 4 and 5; 3 < 4; 4 > 5
VPA counts	$1.14\pm0.39$	1.17	$1.18\pm0.41$	1.17	$1.29\pm0.36$	1.33	$1.14\pm0.34$	1.17	$1.27\pm0.40$	1.33	0.043	1 < 3 and 5; 2 < 3; 3 > 4; 4 < 5
VVPA counts	$0.79\pm0.41$	0.83	$0.45\pm0.34$	0.33	$0.39\pm0.36$	0.33	$0.43\pm0.42$	0.33	$0.44\pm0.37$	0.33	<0.001	1 > 2, 3, 4 and 5 1 > 2, 3, 4
MET rates	$7.85\pm0.72$	7.89	$6.81\pm0.94$	6.85	$6.73\pm0.87$	6.89	$6.48 \pm 1.09$	5.78	$7.02\pm0.92$	7.08	<0.001	and 5; 2 > 4; 2 < 5; 3 > 4; 3 < 5; 4 < 5

*† p*-value from the Friedman test to investigate the change in young football players' internal and external training load across five different small-sided game formats; *‡* Wilcoxon signed-rank test indicating the individual differences in internal and external training load between each game format.

#### 3.1. Vector Magnitude Counts

The Wilcoxon signed-rank test revealed statistically significant higher vector magnitude counts in 3 vs. 3 in comparison to those in 4 vs. 3 (z = -4.99, p < 0.001), 5 vs. 3 (z = -5.28, p < 0.001), 3 vs. 4 (z = -5.35, p < 0.001), 3 vs. 5 (z = -5.14, p < 0.001).

#### 3.2. Light Physical Activity Intensity Counts

There were also statistically significant higher vector magnitude counts in 3 vs. 5 in comparison to those in 4 vs. 3 (z = -2.30, p = 0.021) and 3 vs. 4 (z = -3.72, p < 0.001). There were statistically significant lower light physical activity intensity counts in 3 vs. 3 in comparison to those in 4 vs. 3 (z = -2.53, p = 0.011), and 3 vs. 4 (z = -2.64, p = 0.008). We also identified statistically significant lower light physical activity intensity counts in 4 vs. 3 in comparison to those in 5 vs. 3 (z = -2.53, p = 0.011) and 3 vs. 5 (z = -2.11, p = 0.035). In addition, the 3 vs. 4 format had statistically significant higher light physical activity intensity counts in comparison to those in 5 vs. 3 (z = -2.12, p = 0.034) and 3 vs. 5 (z = -3.30, p = 0.001).

#### 3.3. Moderate Physical Activity Intensity Counts

There were statistically significant lower moderate physical activity intensity counts in 3 vs. 3 in comparison to those in 4 vs. 3 (z = -5.64, p < 0.001), 5 vs. 3 (z = -5.49, p < 0.001), 3 vs. 4 (z = -6.26, p < 0.001), 3 vs. 5 (z = -4.77, p < 0.001). In the 3 vs. 4 format, there were statistically significant higher moderate physical activity intensity counts in comparison to those in 5 vs. 3 (z = -2.12, p = 0.034), and 3 vs. 5 (z = -3.30, p < 0.001).

#### 3.4. Vigorous Physical Activity Intensity Counts

There were statistically significant lower vigorous physical activity intensity counts in 3 vs. 3 in comparison to those in 5 vs. 3 (z = -2.68, p = 0.007) and 3 vs. 5 (z = -2.40, p = 0.016). We also identified statistically significant higher vigorous physical activity intensity counts in 5 vs. 3 in comparison to those in 4 vs. 3 (z = -2.17, p = 0.030) and 3 vs. 4 format (z = -2.46, p = 0.014). Lastly, the 3 vs. 5 format had statistically significant higher vigorous physical activity intensity counts in comparison to those in the 3 vs. 4 format (z = -2.55, p = 0.011).

#### 3.5. Very Vigorous Physical Activity Intensity Counts

There were statistically significant higher very vigorous physical activity intensity counts in 3 vs. 3 in comparison to those in 4 vs. 3 (z = -5.51, p < 0.001), 5 vs. 3 (z = -6.02, p < 0.001), 3 vs. 4 (z = -5.99, p < 0.001), 3 vs. 5 (z = -6.12, p < 0.001).

#### 3.6. METs

There was statistically significant higher energy expenditure (METs) in 3 vs. 3 in comparison to that in 4 vs. 3 (z = -7.17, p < 0.001), 5 vs. 3 (z = -7.20, p < 0.001), 3 vs. 4 (z = -7.23, p < 0.001), 3 vs. 5 (z = -7.00, p < 0.001). We also identified statistically significant higher METs in 4 vs. 3 in comparison to that in 3 vs. 4 format (z = -2.58, p = 0.010). In addition, there were statistically significant higher MET counts in 3 vs. 5 in comparison to those in 4 vs. 3 (z = -2.07, p = 0.039), 5 vs. 3 (z = -2.43, p = 0.015), and 3 vs. 4 (z = -4.27, p < 0.001). Lastly, the 5 vs. 3 format had statistically significant higher MET counts in comparison to that in the 3 vs. 4 format (z = -2.12, p = 0.034).

#### 4. Discussion

This study analysed the effect of different SSG formats with numerical variability on young football players' external training load metrics. In general, the 3 vs. 3 format game had higher physical activity counts (i.e., higher vector magnitude) and requested more time spent in very vigorous physical activity intensity than the 4 vs. 3, 5 vs. 3, 3 vs. 4, and

3 vs. 5 formats did. Consequently, the 3 vs. 3 format in young football players provided higher energy expenditure (i.e., higher METs) than that of other studied game formats.

In the last decade, SSGs have been simultaneously studied as effective training methods with promising results in skills and fitness dimensions. SSGs provide optimal work intensities and appropriate training structures. However, it remains unclear which practice conditions related to the number of players, space, or rules better impact the physiological response.

For example, Clemente et al. [4] proposed a set of methodological suggestions and recommendations to insert SSGs into the training process during a sporting season. When changing the number of players, introducing different conditions to the game rules and modifying the dimensions of the game space, different physiological responses occur. In addition, Little [9], in a review of research concerning conditioning with football drills, claimed that SSGs that involve more players result in appropriate intensities for lactate threshold development. On the other hand, the 3 vs. 3 and 4 vs. 4 game formats can be used to develop maximal oxygen consumption. These results suggest that game formats with fewer players may train and develop aerobic endurance that is specific to high-intensity football. Our study partly supports these findings, since we found statistically significant higher very vigorous physical activity intensity counts and energy expenditure in 3 vs. 3 compared to those in other game formats.

To the best of our knowledge, no other study has considered a similar design to investigate the influence of variability in the number of players on the external training load. The work with the most similar results to our study was reported by Praça, Custódio, and Greco [41]. They concluded that the physical demands of players were reduced when adding an element to a team compared to situations in which they played in numerical equality. The same study concluded that if the teacher or coach's goal is to reduce physical demand or energy expenditure (for reasons such as proximity to a competition or even recovery training after a game), numerical superiority situations are helpful in organising consequent reduction in training intensity.

The present study found that the 3 vs. 3 format game provided higher physical activity counts, increased the time spent in very vigorous physical activity intensity, and requested higher energy expenditure than that of other unbalanced game formats (i.e., 4 vs. 3, 5 vs. 3, 3 vs. 4, or 3 vs. 5). Concerning a teaching-learning approach, some important messages to the practice should be considered from our study to sustain the players' performance in the training sessions. First, when there are modifications in the number of players in action, namely, creating an unbalanced game situation, changes are seen in the external training load and the intensity of the activity. Indeed, some studies agreed with the use of SSGs to induce changes in the external load in relatively short periods, suggesting that the ability to develop the players' physical profile while training the tactical-technical elements might make SSGs a very sustainable training method [42]. Second, if teachers or coaches design SSG for conditioning, the 3 vs. 3 format may more significantly impact physiological and physical levels, and request higher energy expenditure. Introducing one or two players into a team, i.e., creating an unbalanced situation, may reduce the intensity levels in both teams. Third, considering that 3 vs. 3 may increase the time spent in very vigorous physical activity intensity, caution about overtraining is needed in implementing this game format to develop skills and physical fitness.

Despite the recognised potential of SSGs, caution is needed in implementing this method in a training periodisation process. Improper programming can hide shortcomings in performance and the development of physical abilities. It is crucial to strictly control the dynamics of SSGs, i.e., the game's duration when designing them, and consequent framing them in the periodisation and management of training.

Some limitations should be acknowledged when interpreting our results. First, even though three sessions were held for data collection on 3 different days to strengthen the reliability of the data collection, the main focus was on the influence of the formats and not on the players' performance, and only one team under 17 was included. Second, the

scarce literature on the topic using a similar experimental design limits the discussion of the present results. More longitudinal and experimental studies are needed to better understand how small but significant modifications in the number of players can affect external training load and activity intensity.

This study also has some important strengths that must be underlined. First, there are concrete, practical implications that can help teachers or coaches when designing SSGs for improving players' physical profile while training the tactical-technical elements. Second, the uniqueness of the study design, using a changeable SSG format with a duration of 10 min and changing the numerical ratio of players in the game every 2 min without interrupting the practice, opens new avenues on exploring the potential of the SSGs keeping at the same time the main features of the real game. Lastly, the instrument used to collect the external training load could capture and record continuous accurate information about the quantity and intensity of the physical activity.

#### 5. Conclusions

This study emphasised the influence of SSGs with numerical variability in the external training load in young football players. The 3 vs. 3 format provides higher physical activity, increases the time spent in very vigorous physical activity intensity, and requests higher energy expenditure than that of the other studied unbalanced game formats, namely, the 4 vs. 3, 5 vs. 3, 3 vs. 4, or 3 vs. 5 formats. This study reinforces that unbalanced game formats may require a reduction in training intensity, favouring the development of other components related to individual tactics.

**Author Contributions:** Conceptualization, H.S. (Honorato Sousa) and É.R.G.; methodology, H.S. (Honorato Sousa), M.P. and T.Q.; formal analysis, H.S. (Honorato Sousa) and É.R.G.; writing—original draft preparation, H.S. (Honorato Sousa), É.R.G., M.P. and T.Q.; writing—review and editing: É.R.G., A.M., H.S. (Hugo Sarmento), H.L. and A.I.; supervision: É.R.G. and A.I. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was supported by the Swiss National Centre of Competence in Research LIVES, "Overcoming Vulnerability: Life Course Perspectives," granted by the Swiss National Science Foundation (grant number: 51NF40-185901). A.I. acknowledges support from the Swiss National Science Foundation (grant number: 10001C\_189407). É.R.G. acknowledges support from LARSyS—the Portuguese national funding agency for science, research, and technology (FCT) pluriannual funding 2020–2023 (reference: UIDB/50009/2020).

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki and proved by the ethics commission: Ethical Committee of the Faculty of Human Kinetics, University of Lisbon, approval code: CEIFMH No: 35/2021; approval date: 6 July 2021.

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study. Written informed consent has been obtained from the patient(s) to publish this paper.

Data Availability Statement: Not applicable.

Acknowledgments: The authors would like to thank the field team (Romualdo Caldeira, Pedro Luz, and Francisco Fernandes) for the technical assistance in data collection and management. We are especially grateful to the younger soccer players who for their participation in this study, and to the Clube Futebol Andorinha from Santo António, Madeira for providing us with the sports facilities to carry out the study. All individuals included in this section have consented to the acknowledgement.

Conflicts of Interest: The authors declare no conflict of interest.

#### References

- 1. Gabbett, T.J. Skill-based conditioning games as an alternative to traditional conditioning for rugby league players. *J. Strength Cond. Res.* **2006**, *20*, 309. [CrossRef]
- 2. Gabbett, T.; Jenkins, D.; Abernethy, B. Game-based training for improving skill and physical fitness in team sport athletes. *Int. J. Sports Sci. Coach.* 2009, *4*, 273–283. [CrossRef]

- 3. Davids, K.; Araújo, D.; Correia, V.; Vilar, L. How small-sided and conditioned games enhance acquisition of movement and decision-making skills. *Exerc. Sport Sci. Rev.* **2013**, *41*, 154–161. [CrossRef] [PubMed]
- 4. Clemente, F.M.; Martins, F.M.L.; Mendes, R.S. Periodization based on small-sided soccer games: Theoretical considerations. *Strength Cond. J.* 2014, *36*, 34–43. [CrossRef]
- Hill-Haas, S.V.; Dawson, B.; Impellizzeri, F.M.; Coutts, A.J. Physiology of small-sided games training in football: A systematic review. Sports Med. 2011, 41, 199–220. [CrossRef]
- 6. Sgrò, F.; Bracco, S.; Pignato, S.; Lipoma, M. Small-Sided Games and Technical Skills in Soccer Training: Systematic Review and Implications for Sport and Physical Education Practitioners. *J. Sports Sci.* **2018**, *6*, 9–19. [CrossRef]
- Gabbett, T.J.; Mulvey, M.J. Time-motion analysis of small-sided training games and competition in elite women soccer players. J. Strength Cond. Res. 2008, 22, 543–552. [CrossRef]
- 8. Bangsbo, J. Physiology of training. In Science and Soccer; Reilly, T., Williams, A.M., Eds.; Routledge: London, UK, 2003; pp. 47–58.
- 9. Little, T. Optimizing the use of soccer drills for physiological development. Strength Cond. J. 2009, 31, 67–74. [CrossRef]
- 10. Clemente, F.M. Small-Sided and Conditioned Games in Soccer Training: The Science and Practical Applications, 1st ed.; Springer Imprint: Singapore, 2016.
- Martín Martínez, I.; Reigal Garrido, R.E.; Chirosa Ríos, L.J.; Hernández Mendo, A.; Chirosa Ríos, I.; Martín Tamayo, I.; Guisado Barrilao, R. Efectos de un programa de juegos reducidos en la percepción subjetiva del esfuerzo en una muestra de chicas adolescentes. *Cuad. Psicol. Deporte* 2015, 15, 89–98. [CrossRef]
- 12. Casamichana, D.; Castellano, J.; Hernández-Mendo, A. Generalizability Theory applied to the study of the physical profile during reduced games with different spatial orientation in soccer. *Int. J. Sport Sci.* 2014, *10*, 195–205.
- Arslan, E.; Kilit, B.; Clemente, F.M.; Soylu, Y.; Sögüt, M.; Badicu, G.; Akca, F.; Gokkaya, M.; Murawska-Ciałowicz, E. The Effects of Exercise Order on the Psychophysiological Responses, Physical and Technical Performances of Young Soccer Players: Combined Small-Sided Games and High-Intensity Interval Training. *Biology* 2021, 10, 1180. [CrossRef]
- 14. Teixeira, J.E.; Forte, P.; Ferraz, R.; Leal, M.; Ribeiro, J.; Silva, A.J.; Barbosa, T.M.; Monteiro, A.M. Monitoring accumulated training and match load in football: A systematic review. *Int. J. Environ. Res. Public Health* **2021**, *18*, 3906. [CrossRef]
- 15. Younesi, S.; Rabbani, A.; Clemente, F.; Sarmento, H.; JFigueiredo, A. Session-to-session variations in external load measures during small-sided games in professional soccer players. *Biol. Sport* **2021**, *38*, 185–193. [CrossRef] [PubMed]
- 16. Owen, A. Physiological and Technical Analysis of Small-Sided Conditioned Training Games within Professional Football; SAGE Publications: Wrexham, Welsh, 2003.
- 17. Gamble, P. A skill-based conditioning games approach to metabolic conditioning for elite rugby football players. *J. Strength Cond. Res.* **2004**, *18*, 491–497. [CrossRef]
- Arslan, E.; Orer, G.; Clemente, F. Running-based high-intensity interval training vs. small-sided game training programs: Effects on the physical performance, psychophysiological responses and technical skills in young soccer players. *Biol. Sport* 2020, 37, 165–173. [CrossRef] [PubMed]
- 19. Balsom, P.; Lindholm, T.; Nilsson, J.; Ekblom, B. Precision Football; Polar Electro Oy: Kempele, Finland, 1999.
- Jeffreys, I. The use of small-sided games in the metabolic training of high school soccer players. *Strength Cond. J.* 2004, 26, 77–78. [CrossRef]
- Sarmento, H.; Clemente, F.M.; Harper, L.D.; Costa IT da Owen, A.; Figueiredo, A.J. Small sided games in soccer–a systematic review. Int. J. Perform. Anal. Sport 2018, 18, 693–749. [CrossRef]
- Clemente, F.M.; Ramirez-Campillo, R.; Sarmento, H.; Praça, G.M.; Afonso, J.; Silva, A.F.; Rosemann, T.; Knechtle, B. Effects of small-sided game interventions on the technical execution and tactical behaviors of young and youth team sports players: A systematic review and meta-analysis. *Front. Psychol.* 2021, *12*, 667041. [CrossRef]
- 23. Hill-Haas, S.V.; Coutts, A.J.; Dawson, B.T.; Rowsell, G.J. Time-motion characteristics and physiological responses of small-sided games in elite youth players: The influence of player number and rule changes. *J. Strength Cond. Res.* **2010**, *24*, 2149–2156. [CrossRef]
- 24. Castellano, J.; Silva, P.; Usabiaga, O.; Barreira, D. The influence of scoring targets and outer-floaters on attacking and defending team dispersion, shape and creation of space during small-sided soccer games. *J. Hum. Kinet.* **2016**, *51*, 153–163. [CrossRef]
- 25. Serra-Olivares, J.; González-Víllora, S.; García-López, L.M.; Araújo, D. Game-based approaches' pedagogical principles: Exploring task constraints in youth soccer. J. Hum. Kinet. 2015, 46, 251–261. [CrossRef]
- MacLaren, D.; Davids, J.; Isokawa, M.; Mellor, S.; Reilly, T. Physiological strain in 4-a-side soccer. In Science and Football I; Reilly, T., Lees, A., Davids, K., Murphy, W.J., Eds.; E & FN Spon: London, UK, 1988; pp. 76–80.
- 27. Reilly, T. An ergonomics model of the soccer training process. J. Sports Sci. 2005, 23, 561–572. [CrossRef]
- 28. Castagna, C.; Belardinelli, R.; Impellizzeri, F.M.; Abt, G.A.; Coutts, A.J.; D'Ottavio, S. Cardiovascular responses during recreational 5-a-side indoor-soccer. *J. Sci. Med. Sport* **2007**, *10*, 89–95. [CrossRef] [PubMed]
- 29. Jensen, J.M.; Randers, M.B.; Krustrup, P. Intermittent high-intensity drills improve in-seasonal performance of elite soccer players. In *Science and Football VI*; Routledge: London, UK, 2008; pp. 322–327.
- 30. Boyd, L.J.; Ball, K.; Aughey, R.J. The reliability of minimaxx accelerometers for measuring physical activity in australian football. *Int. J. Sports Physiol. Perform.* 2011, *6*, 311–321. [CrossRef] [PubMed]
- 31. Rampinini, E.; Impellizzeri, F.M.; Castagna, C.; Abt, G.; Chamari, K.; Sassi, A.; Marcora, S.M. Factors influencing physiological responses to small-sided soccer games. *J. Sports Sci.* 2007, 25, 659–666. [CrossRef] [PubMed]

- 32. Arslan, E.; Alemdaroglu, U.; Koklu, Y.; Hazir, T.; Muniroglu, S.; Karakoc, B. Effects of passive and active rest on physiological responses and time motion characteristics in different small sided soccer games. *J. Hum. Kinet.* **2017**, *60*, 123–132. [CrossRef]
- Guard, A.N.; McMillan, K.; MacFarlane, N.G. The influence of relative playing area and player numerical imbalance on physical and perceptual demands in soccer small-sided game formats. *Sci. Med. Footb.* 2021, 1–7. [CrossRef]
- Sampaio, J.E.; Lago, C.; Gonçalves, B.; Maçãs, V.M.; Leite, N. Effects of pacing, status and unbalance in time motion variables, heart rate and tactical behaviour when playing 5-a-side football small-sided games. J. Sci. Med. Sport 2014, 17, 229–233. [CrossRef]
- Torres-Ronda, L.; Gonçalves, B.; Marcelino, R.; Torrents, C.; Vicente, E.; Sampaio, J. Heart rate, time-motion, and body impacts when changing the number of teammates and opponents in soccer small-sided games. *J. Strength Cond. Res.* 2015, 29, 2723–2730. [CrossRef] [PubMed]
- Bekris, E.; Gkisis, I.; Milonys, E.; Anagnostakos, K.; Sarakinos, A.; Sambanis, M. The Physiological and Technical-Tactical Effects of an Additional Soccer Player's Participation in Small Sided Games Training (No. IKEEART-2015-3845); Aristotle University of Thessaloniki: Thessaloniki, Greece, 2012.
- Evangelos, B.; Eleftherios, M.; Aris, S.; Ioannis, G.; Konstantinos, A.; Natalia, K. Supernumerary in small sided games 3vs3 & 4vs4. J. Phys. Educ. Sport 2012, 12, 398.
- Faul, F.; Erdfelder, E.; Lang, A.-G.; Buchner, A. G\*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav. Res. Methods* 2007, *39*, 175–191. [CrossRef] [PubMed]
- 39. World Medical Association. Declaration of Helsinki. Int. J. Pharm. Med. 2000, 14, 279–281. [CrossRef]
- 40. Casamichana, D.; Castellano, J. Time–motion, heart rate, perceptual and motor behaviour demands in small-sides soccer games: Effects of pitch size. *J. Sports Sci.* **2010**, *28*, 1615–1623. [CrossRef]
- 41. Praça, G.M.; de Oliveira Custódio, I.J.; Greco, P.J. Superioridade numérica altera a demanda física de jogadores de Futebol durante Pequenos Jogos. *Rev. Bras. Cineantropometria Desempenho Hum.* **2015**, *17*, 269–279. [CrossRef]
- Owen, A.L.; Wong, D.P.; Paul, D.; Dellal, A. Effects of a periodized small-sided game training intervention on physical performance in elite professional soccer. J. Strength Cond. Res. 2012, 26, 2748–2754. [CrossRef]