

This publication URL: Publication DOI:

https://archive-ouverte.unige.ch/unige:172646 10.1016/j.pec.2023.107894

© This document is protected by copyright. Please refer to copyright holders for terms of use.

Patients preferences for communication during video consultations

Sanae Mazouri-Karker, Olivia Braillard, Robin Lüchinger, Nadia Bajwa, Sophia Achab, Patricia Hudelson, Mélissa Dominicé Dao, Noëlle Junod-Perron



PII: S0738-3991(23)00274-4

DOI: https://doi.org/10.1016/j.pec.2023.107894

Reference: PEC107894

To appear in: Patient Education and Counseling

Please cite this article as: Sanae Mazouri-Karker, Olivia Braillard, Robin Lüchinger, Nadia Bajwa, Sophia Achab, Patricia Hudelson, Mélissa Dominicé Dao and Noëlle Junod-Perron, Patients preferences for communication during video consultations, *Patient Education and Counseling*, (2023) doi:https://doi.org/10.1016/j.pec.2023.107894

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier.

### Patients preferences for communication during video consultations

Sanae Mazouri-Karker<sup>1</sup>, Olivia Braillard<sup>2</sup>, Robin Lüchinger<sup>3</sup>, Nadia Bajwa<sup>2,3</sup> Sophia Achab<sup>2,4</sup>, Patricia Hudelson<sup>2</sup>, Mélissa Dominicé Dao<sup>2,3</sup>, Noëlle Junod-Perron<sup>2,3</sup>,

1. E-health and Telemedicine Division, Geneva University Hospitals, Geneva, Switzerland

2. Primary Care Division, Geneva University Hospitals, Geneva, Switzerland

3. Faculty of Medicine, University of Geneva, Geneva, Switzerland

4. ReConnecte Unit, Addiction Division, University Hospitals, Geneva, Switzerland

SANAE MAZOURI-KARKER, Diagnostic Department, eHealth and Telemedicine Service, Geneva University Hospitals. Geneva, Switzerland. ORCID: https://orcid.org/0000-0003-3754-8742

Email: dr.mazouri@cmtb.ch

OLIVIA BRAILLARD, Department of Primary Care Medicine, Geneva University Hospitals, Geneva, Switzerland. ORCID: https://orcid.org/ 0000-0003-2720-3900

Email: olivia.braillard@hcuge.ch

ROBIN LUECHINGER, Faculty Member at the Unit of Development and Research in Medical Education, Faculty of Medicine, University of Geneva, Geneva, Switzerland.

ORCID : https://orcid.org/0000-0002-6169-2704

Email: robin.luechinger@unige.ch

NADIA M. BAJWA, Residency Program Director in the Department of General Pediatrics at the Children's Hospital, Geneva University Hospitals, and Faculty Member at the Unit of Development and Research in Medical Education (UDREM), Faculty of Medicine, University of Geneva, Geneva, Switzerland. ORCID : http://orcid.org/0000-0002-1445-4594

Email: nadia.bajwa@hcuge.ch

SOPHIA ACHAB, Clinical Director, Treatment Centre ReConnecte, Department of Psychiatry, University Hospitals of Geneva, Switzerland.Head, Clinical and Sociological Research Unit (WHO

Collaborating Centre for Training and Research in Mental Health), Department of Psychiatry, Faculty of Medicine, University of Geneva, Switzerland https://orcid.org/0000-0002-3861-3297

Email: sophia.achab@hcuge.ch

PATRICIA HUDELSON-PERNEGER, Department of Primary Care Medicine, Geneva University Hospitals and Faculty of Medicine, University of Geneva. ORCID: https://orcid.org/0000-0003-2601-7479

Email: Patricia.Hudelson@hcuge.ch

MELISSA DOMINICE DAO, Department of Primary Care Medicine, Geneva University Hospitals and Faculty of Medicine, University of Geneva. ORCID : https://orcid.org/0000-0002-0839-277X

Email: melissa.dominice@hcuge.ch

NOELLE JUNOD PERRON, Medical Directory, Geneva University Hospitals and Faculty Member at the Unit of Development and Research in Medical Education, Faculty of Medicine, University of Geneva, Geneva, Switzerland. ORCID : https://orcid.org/0000-0002-9124-8663

Email : Noelle.Junod@hcuge.ch

# Contact information for the corresponding author

### SANAE MAZOURI-KARKER<sup>1</sup>

Address : CMTB Medical Centre Terre-Bonne, Crassier road 7, 1262 Eysins, Switzerland

Email dr.mazouri@cmtb.ch

Fax: no fax

Phone number: +41 763980178

### Abstract

ObjectiveThe aim of our simulation-based study was to explore patient preferences for physician behaviours in video consultations

### Methods

We conducted an exploratory study in outpatient setting in Geneva, Switzerland.Patients were invited to watch two variations videos of six simulated physician communication behaviours (camera framing, gaze orientation, initial talk at the opening phase, privacy reminder, pauses, empathy)and to indicate which one they preferred

# Results

417 patients watched three different video-recorded encounters. Most patients preferred framing with both face and bust (50.7%) versus face alone (21.8%). They valued eye gazing towards the camera (42.9%) versus eye gazing shifting between screen and camera (13%). The social talk related to the connection quality was appreciated (43.1% vs 17.1%) as well as the privacy reminder (80.8% vs 6.5%).Patients preferred short rather than long pauses after physician's statements (63.9 vs 14.9%) as well as expressive rather than neutral nonverbal behaviour (46.7% vs 17.6%).

# Conclusion

Our results confirm that patients prefer the use of video specific communication behaviours recommended by experts except for shifting eye gaze and long pauses after physician's statements.

# **Practice Implications**

Given the increasing use of video consultations, video communication "best practices" should be systematically addressed in physician training

# Key words

Video consultation; telemedicine; ehealth; physician's behaviour; patient centered communication

### Introduction

Telemedicine (TLM) includes remote diagnosis, treatment, care, and patient education with the use of several telecommunications tools [1, 2]. Remote consultations can be synchronous (using tools such as video, telephone) or asynchronous (use of text-message, email). Telemedicine can facilitate communication among health professionals as well as between professionals and their patients by improving access to care and specialized advice [3, 4]. Technological evolution has enabled a rapid expansion of telemedicine, providing healthcare professionals with new ways to communicate with their patients beyond the traditional model of in-person visits [5], by transforming personal face-to face communication [6] and creating a new health management approach and communication context [7, 8], particularly for chronic diseases where videoconferencing has been found to be feasible and effective [9]. Videoconferencing has also offered opportunities for remote physical examination of patients with chronic conditions such as heart failure, [10] and with similar outcomes when used to replace or augment traditional care [11]. Experiences during the covid pandemic, during which remote

consultations were more frequent, suggest that teleconsultations by phone or video may also be an effective alternative to face-to-face consultations for many patients attending primary care and mental health services [12-20].

The quality of verbal and nonverbal physician-patient communication is a critical predictor of treatment outcomes including physician-patient agreement, understanding of health problems, satisfaction with care, and patient adherence [21, 22]. Many studies have analysed the quality of physician-patient communication in face-to-face consultation [23-25] but less is known on the quality of verbal and nonverbal communication in synchronous telemedicine (via videoconferencing or phone calls). It is known that compared to face-to face consultations, TLM can influence health outcomes through changes in the form and content of physician-patient communication. Indeed, the characteristics of the patient, the physician and the context of the medical consultation can influence the nature and content of the physician-patient communication which, in return, will influence health behaviours [26]. This process depends on the mode used for the consultation: in person, by telephone, by video, etc.

Studies in out-patient psychiatry suggest that there is only a slight decrease in patient satisfaction with video consultations compared to a face-to-face consultations [27], no differences in verbal content between the four modes of consultation (face-to-face, telephone, hands-free telephone and video conferencing system) and positive reactions from both patients and psychiatrists towards the use of video conferencing for the consultations [28]. Studies in internal medicine also found that content and patient satisfaction were similar for in-person and remote consultations [29-31].

Nonetheless, there is evidence that video consultations differ on a number of important dimensions than can affect the quality and outcomes of the interaction, including patient participation, decision sharing, expression of empathy, and attention to verbal cues [32]. Other studies showed that physicians and patients use more verbal cues to establish or improve interpersonal relationship in order to compensate the reduction in nonverbal cues in video consultations [33-35]. In an effort to optimize communication during video consultations, best practice recommendations have been issued by a number of experts and national medical associations (Table 1) [33, 36-42]. However, there is a lack of information on patient perspectives regarding these communication strategies, and it is possible that patients may be sensitive to additional elements of the context and interactions during a video consultation.

In an effort to fill this gap, we aimed to describe patient preferences regarding physicians' communication behaviours in simulated video consultations, identify patient socio-demographic factors associated with their preferences, and explore the reasons for their preferences.

#### Methods

#### Design and setting

We conducted an online survey between October and December 2021 in Geneva, Switzerland.

We recruited patients from three outpatient settings that provide consultations for patients presenting with medical and trauma-related problems that do not require hospitalization: a walk-in clinic at Geneva University Hospitals (providing approximately 35,000 consultation per year), and two walk-in clinics situated in the suburbs of Geneva (each providing 26,000 to 35,000 consultations per year).

### Participants and procedure

Patients waiting for a medical consultation who were French-speaking and over 18 years of age were invited to take part in an online survey on a tablet provided by one of 8 research assistants. Research assistants explained the study objectives and procedures, including the anonymous analysis and reporting of study findings, and obtained written consent to participate. Patients who could not read and understand French were excluded.

The study was granted a waiver of ethical exemption by the Ethical Committee of the Canton of Geneva since it did not involve collecting any personal health information (article 2 of the Swiss Federal Act on Research involving Human Beings) [43].

### Development of videos displaying telemedicine-related behaviours

We reviewed the current guidelines on how to communicate during video-based consultations [33, 36, 37, 39, 40] and identified a list of six behaviours that were specific to video-based consultations or for which there was a lack of consensus among experts. Nonverbal communication behaviours included: framing of the doctor, level and orientation of eye gazing, duration of pauses, nonverbal expression of empathy. Verbal communication behaviours included confidentiality issues (privacy reminder) and initial social talk (questions about the patient's comfort and environment). (Table 2).

Five physicians (3 women, 5 men; aged 35-50 years) involved in primary care teaching were selected to act in the videos. They were of different ages (35 to 50 years) and gender (three women and three men). They were expected to demonstrate good communication skills independently of the use of EHR. Physicians were paired with simulated patients of varying ages and gender who work at the Geneva Faculty of Medicine, with the aim of limiting gender or age biases. Video scripts focused on 6 common complaints in primary care. The distribution of complaints, physicians and patients are shown in Table 2.

We wrote a script for each video-specific behaviour. For each behaviour, two sequences were recorded, each showing variations on the behaviour. Physician-actors were asked to closely follow the script and replicate exactly the same verbal and nonverbal communication for each of the two sequences, with the exception of the video-related behaviour variations (i.e duration of the pauses...). The simulated encounters were video-recorded from a patient perspective by a professional videographer in the presence of two investigators in order to ensure that the acting physicians respected the instructions. The sequences were repeated until the verbal and nonverbal communication unrelated to video-related behaviours and variations in video-related behaviours). We performed a manipulation check asking three experienced primary care physicians, blinded to the study objectives, to identify and validate the video-recorded variations of the different video-related behaviours. We also asked them to check whether the physician-actors displayed similar verbal and nonverbal communication unrelated to the video in the different sequences for each scenario.

# Procedure

Participants were invited to watch three different pairs of videos presented randomly and integrated in an online survey on a tablet in the waiting room. They were asked to watch the videos as if they were the patient in the consultation. It took them approximatively 15 minutes to complete the survey: 10 minutes to watch videos, indicate their preferences, and write down the reasons for their preferences, and 5 minutes to answer socio-demographic information. Each set of videos included two different sequences displaying variations of the tested video-related behaviour. Both the videos and sequences were presented randomly. After watching the video set, patients selected their preference directly on the tablet (option 1 or 2 or no preference). In case of a preference, they were asked to write down the reasons.

#### Outcomes, and other measures

The main outcome was patients' preference among two variations regarding the six video-related behaviours. However, if they could not reach a decision, they could indicate "no preference." At the end of the survey, patients were also asked to provide socio-demographic information such as age, gender, civil status, nationality, and employment. We also recorded study location (hospital versus community emergency service) in order to account for potential location-related biases.

We also noted any reasons spontaneously offered by patients for their preferences.

#### Sample size estimate

We aimed for a sample size that would have a 90% power of detecting a 20% difference in the proportion of patients choosing a specific sequence compared to the null hypothesis (50% of participants choose the sequence). This led to a sample size of 200 patients. Given that each patient had to assess only three of the six video sets, we doubled this number to include 400 patients.

### Statistical analysis

We used Stata software version xx.0 for the analysis. We used proportions to summarize patient preferences for a sequence in each video set [44]. Difference of preferences were calculated using Chi square test with p<0.05. We conducted a multivariate analysis to investigate the association between patients' preferences and their sociodemographic data.

Reasons given for preferences were analysed as "congruent" or "non-congruent" with respect to the focal behaviour in the video-recorded sequence: e.g. if a patient reported that he/she preferred the face framing because the doctor looked closer, we coded this reason as congruent since the patient's preference was related to the focal behaviour in the video; it was coded as non-congruent if the patient reported that they preferred the fame framing video because they liked the doctor's tone of voice [44] . Four investigators (SM, MDD, PH, NJP) double coded patients' reasons for their preference (2 investigators per theme) and differences were resolved by discussion among the four investigators. Results were expressed in percentages.

### Results

417 patients watched videos of 3 different standardized encounters illustrating specific behaviours (two variations of each behaviour). The majority of patients were below 45 years old, employed and of Swiss or European nationality (Table 3).

Patient preferences are indicated in Figure 1. Patients preferred the following communication behaviours recommended by experts: framing that includes the bust and face, addressing confidentiality issues at the beginning of the encounter, making the patient feel comfortable by asking questions about the quality of the connection (sound and image) and increased expression of nonverbal empathy in response to the patient's distress. Patients did not show a preference for eye gazing towards the patient and longer pauses between sentences, which are also recommended by experts.

In multivariate analyses, non-European nationality was associated with a preference for longer pauses (OR 3.1 (1.14-8.40) p=0.03) and having no preference regarding asking about the quality of the connection (OR 0.23 (0.06-0.20, p = 0.02).

Reasons given for preferences (Table 4) were largely concordant for the video sequences regarding nonverbal empathy behaviours, moderately concordant for confidentiality issues, and pauses. There were more discordant reasons given for eye gazing, framing and to a lesser extent for confidentiality issues.

Discussion and conclusion

### Discussion

The aim of our simulation-based study was to explore patient preferences for physician behaviours in video consultations with respect to verbal and nonverbal communication. We found that a majority of patients showed a preference for expert-recommended communication behaviours, including framing of the bust and face, addressing confidentiality issues at the beginning of the encounter, making the patient comfortable by asking questions about the quality of the sound and image, and increased expression of nonverbal empathy in response to the patient's distress. Themes that were not chosen by patients although recommended by experts were: eye gazing towards the camera and longer pauses between physicians' sentences. The only sociodemographic factor associated with patient preferences was having a non-European nationality.

#### Nonverbal communication

Nonverbal behaviour, a key component of communication, includes several elements such as interpersonal distance, body position, gestures, facial expression and eye contact.[45]. Such elements have been shown to be important to understanding clinical interactions [31, 32, 33], In our study, patients' preferences for a framing of the face and the chest may be due to the fact that it closely resembles the way patients see their physician in the consultation room and allows patients to situate the physician's environment. It may also facilitate perception of nonverbal communication such as interpersonal distance, gestures, and body position that indicate to which extent physicians engage and direct their attention to patients [41].

Visual contact is another important nonverbal behaviour [46]. Several authors recommended simulating eye contact during video consultations by gazing at the camera rather than at the screen as a potential way to reinforce patient centred communication [38, 41, 46]. Indeed, the position of the webcam can sometimes make a mutual gaze particularly difficult. We found that a similar number of patients preferred eye gazing towards the screen or were indifferent to the type of eye gazing. Our findings differ from another study reporting that patients from both Lebanon and Japan largely favoured eye gazing at the camera and linked it to higher communication and interpersonal skills ratings [47]. Our findings may be due to the fact that the difference between screen and camera gazing in our videos was very subtle. Indeed, only a minority of patients identified that eye gazing was different between the two video-recorded consultation sequences.

Use of pauses or silence has been encouraged to facilitate patient contribution both in the gathering information and explanation phases of the consultation [48]. In video-mediated interaction, there are short periods of latency (milliseconds-long delays in image or sound) that can interfere with conversation turn-taking, [49] and therefore some experts have recommended longer pauses during physician speech to avoid misunderstanding and speech overlaps [39]. However, we found that patients did not favour longer pauses during a video-consultation about a positive occult faecal blood (colorectal cancer screening). It may be that when discussing a worrisome result, longer pauses have the potential to increase the patient's anxiety. Delays or longer pauses may also be interpreted as a problem with the video system as in face-to-face conversations, participants interact with no delay, each person speaking when the other has finished. Interestingly, Shaw et al. found that short periods

of latency in video consultations were tolerated or ignored by participants whereas there was actually more overlapping talk or interruption with longer periods of latency [49]. Finally, the association between non-European origin and a preference for longer pauses between sentences may be related to the fact that non-European patients may have less command of the French language and prefer a slower information rate. Longer pauses may allow them time for translation and integration of information and facilitate trust-building between non- French speaking patients and their physicians [50].

Finally, patients in our study preferred increased expression of nonverbal empathy in response to the patient's distress displayed in the video-consultation. This is not surprising since poor body language and nonverbal communication are often mentioned as barriers to telehealth adoption [51-54]. Indeed, new technologies tend to negatively influence patient-physician communication through limitation of both patient participation and of sensory and nonverbal expressions [55]. Our findings support expert recommendations to use more paraverbal cues, gestures and active listening during video consultations to compensate the sensory and nonverbal limitations inherent to video consultations and to reinforce verbal communication.[42]

### Verbal communication

The opening phase of a consultation in which the physicians put the patient at ease and identifies the topics the patient wishes to discuss, is of crucial importance. It can impact the accuracy and efficiency of the consultation as well as on the nature of the physician-patient relationships [48]. The opening phase of video consultations requires additional verbal talk in the absence of mutual gaze, handshaking and walking with the patient to the consultation room [56]. In our study, most patients preferred the video sequence where confidentiality issues were addressed and more than half gave a congruent reason for their choice. Other studies have also found that patients are concerned about security and privacy [57]. In order to respond to these concerns, experts recommend addressing confidentiality at the very beginning of the video consultation [38] by taking verbal consent for the video, proceeding to mutual identification (including naming the people present in the room) and reassuring the patient that the video call is confidential and secure [38, 42].

Finally, while experts recommend checking the quality of the connection at the beginning of the consultation [41, 49, 58], only some patients in our study appreciated physicians' verbal attempts at

small talk and questions about the quality of the video connection; a large percent of patients indicated no preference. Our findings somewhat support expert recommendations regarding checking the quality of the connection at the beginning of the consultation [41, 49, 58]. Indeed, some patients and physicians can experience stress at the beginning of the video encounter because of the different many steps needed to establish a connection [49, 58]. Paying attention to this issue specific to video consultations is another way of demonstrating interest and respect regarding patient's comfort. The fact that non-European patients were less sensitive to this issue may be explained by the fact that they may be used to experience video calls of poor quality since internet speed may be lower in other parts of the world [59]. However, the design of the study did not allow us to test this hypothesis.

### Limitations

Our study has several limitations. As this was a simulation-based study, participants were the raters of observed differences in communication styles. As un-trained raters, their impressions may have been influenced by multiple factors limiting the generalizability of our results to the real-world setting [60].

Participants were patients in out-patient waiting rooms in a French-speaking city of Switzerland who were able to understand and read French. Their preferences may not be representative of patients elsewhere. Patients were mostly younger than 45 years old and educated. We collected little information about patients' social and economic status apart from nationality, level of education, and no information about health status. Therefore, we are not able to assess the impact of social, linguistic and economic status, health conditions or health and digital literacy on patients' preferences regarding physicians' behaviour in video consultations. Patients were asked to watch the videos as if they were the patient in the video consultation, but their own health problems could have influenced their preferences. We also did not examine the influence of gender and age match between participating patients and simulated patients in the videos. However, the use of real patients as raters increased the fidelity of our simulation study.

Patients' preferences were explored through video-recorded simulated consultations addressing specific health issues (e.g., patients' preferences regarding nonverbal empathy were explored in an encounter where the patient expressed high distress, the confidentiality issue was addressed for a sensitive topic (condom break). It is possible that the topic of the consultation may have influenced patients' preferences since communication is both context dependent and goal oriented [61, 62].

Nonetheless, we chose these topics because we considered that some video-related behaviours recommended by experts were specifically relevant to these clinical situations.

We only considered the socio-demographic factors of patients, but probably those of physicians also have an influence on patients' preferences. Also, **our analysis did not take into account the multilevel structure of our data set. Future analyses could cluster participants based on sociodemographic characteristics and the survey location to give more insight into patient preferences**.

Finally, our study design only allowed for patients' preferences as outside observers, our findings cannot be generalized to the real world where elements influencing communication are multiple and complex, including the duration and the quality of the patient-physician relationship. Future studies should be conducted during live video consultations with real patients using the different communication behaviours that were simulated in this study.

### Conclusion

Our study suggests that patients are attentive to and have preferences for a number of physician communication behaviours that are recommended by experts for use during video consultations. Although our findings should be repeated in different contexts of care and among more groups of patients with diverse ages, cultural, socio-economic backgrounds and levels of digital literacy and cross-checked with results of studies exploring patients' preferences as actors in real consultations.

# Practice Implications

Given the increasing use and importance of remote consultations, video communication "best practices" should be systematically addressed in physician training.

### **Declaration of Competing Interest**

This manuscript has not been previously published and is not under consideration in the same or substantially similar form in any other peer-reviewed media. All authors listed have contributed sufficiently to the project, and all those who are qualified to be authors are listed in the author by-line. No conflicts of interest, financial or other, exist.

# Acknowledgements

Contributors: The authors wish to thank the "Private Foundation of Geneva University Hospitals" for

supporting our work as well as all the participants who took part into the study (patients, physicians,

and research assistants).

# References

- 1. Chen, M.-J., et al., A telehealth service model for the treatment of hypertension. Journal of Telemedicine and Telecare, 2013. **19**(5): p. 238-241.
- 2. Stanberry, B., *Telemedicine: barriers and opportunities in the 21st century.* Journal of Internal Medicine, 2000. **247**(6): p. 615-628.
- 3. Johansson, A.M., I. Lindberg, and S. Söderberg, *Patients' experiences with specialist care via video consultation in primary healthcare in rural areas.* International journal of telemedicine and applications, 2014. **2014**.
- 4. Sims, J.M., *Communities of practice: Telemedicine and online medical communities.* Technological Forecasting and Social Change, 2018. **126**: p. 53-63.
- 5. Matusitz, J. and G.-M. Breen, *Telemedicine: its effects on health communication*. Health communication, 2007. **21**(1): p. 73-83.
- Turner, J.W., et al., Media Attitudes vs. Use: The Contribution of Context to the Communication Environment in Telemedicine. Health Care Management Review, 2003. 28(2): p. 95-106.
- Gustke, S.S., et al., *Patient Satisfaction with Telemedicine*. Telemedicine Journal, 2000. 6(1): p. 5-13.
- 8. Munos, B., et al., *Mobile health: the power of wearables, sensors, and apps to transform clinical trials.* Annals of the New York Academy of Sciences, 2016. **1375**(1): p. 3-18.
- 9. Mallow, J.A., et al., *The Use of Video Conferencing for Persons with Chronic Conditions: A Systematic Review*. Ehealth Telecommun Syst Netw, 2016. **5**(2): p. 39-56.
- 10. Seuren, L.M., et al., *Physical Examinations via Video for Patients With Heart Failure: Qualitative Study Using Conversation Analysis.* J Med Internet Res, 2020. **22**(2): p. e16694.
- 11. Albritton, J., et al., *Video Teleconferencing for Disease Prevention, Diagnosis, and Treatment.* Annals of Internal Medicine, 2021. **175**(2): p. 256-266.
- 12. Fisk, M., A. Livingstone, and S.W. Pit, *Telehealth in the Context of COVID-19: Changing Perspectives in Australia, the United Kingdom, and the United States.* J Med Internet Res, 2020. **22**(6): p. e19264.
- 13. Lam, A., et al., *Videoconsultation to overcome barriers during COVID-19.* Ann Acad Med Singap, 2021. **50**(1): p. 77-83.
- 14. Mann, D.M., et al., *COVID-19 transforms health care through telemedicine: evidence from the field.* Journal of the American Medical Informatics Association, 2020. **27**(7): p. 1132-1135.
- Mazouri-Karker, S., *Revue Médicale Suisse : Télémédecine à l'ère du COVID-19 : une révolution ? Expérience des hôpitaux universitaires de Genève.* Revue Médicale Suisse, 2020.
   16(706): p. 1695-1698.
- Ohannessian, R., T.A. Duong, and A. Odone, *Global Telemedicine Implementation and Integration Within Health Systems to Fight the COVID-19 Pandemic: A Call to Action.* JMIR Public Health Surveill, 2020. 6(2): p. e18810.
- 17. Petrazzuoli, F., et al., *Patient consultations during SARS-CoV-2 pandemic a mixed-method cross-sectional study in 16 European countries.* Rural & Remote Health, 2022. **22**(4).

- 18. Portnoy, J., M. Waller, and T. Elliott, *Telemedicine in the era of COVID-19.* The Journal of Allergy and Clinical Immunology: In Practice, 2020. **8**(5): p. 1489-1491.
- 19. Rockwell, K.L. and A.S. Gilroy, *Incorporating telemedicine as part of COVID-19 outbreak response systems.* Am J Manag Care, 2020. **26**(4): p. 147-148.
- 20. Carrillo de Albornoz, S., K.-L. Sia, and A. Harris, *The effectiveness of teleconsultations in primary care: systematic review.* Family Practice, 2022. **39**(1): p. 168-182.
- 21. DiMatteo, M.R., *The physician—patient relationship: effects on the quality of health care.* Clinical obstetrics and gynecology, 1994. **37**(1): p. 149-161.
- Hall, J.A. and M.C. Dornan, *Meta-analysis of satisfaction with medical care: description of research domain and analysis of overall satisfaction levels.* Social science & medicine, 1988.
   27(6): p. 637-644.
- 23. Bensing, J., *Doctor-patient communication and the quality of care.* Social science & medicine, 1991. **32**(11): p. 1301-1310.
- 24. Buller, M.K. and D.B. Buller, *Physicians' Communication Style and Patient Satisfaction*. Journal of Health and Social Behavior, 1987. **28**(4): p. 375-388.
- 25. Ong, L.M.L., et al., *Doctor-patient communication: A review of the literature.* Social Science & Medicine, 1995. **40**(7): p. 903-918.
- 26. Miller, E.A., *Telemedicine and doctor–patient communication: a theoretical framework for evaluation.* Journal of Telemedicine and Telecare, 2002. **8**(6): p. 311-318.
- 27. Dongier, M., et al., *Telepsychiatry: psychiatric consultation through two-way television. A controlled study.* The Canadian Journal of Psychiatry, 1986. **31**(1): p. 32-34.
- 28. Ball, C., et al., *A comparison of communication modes in adult psychiatry*. Journal of Telemedicine and Telecare, 1995. **1**(1): p. 22-26.
- 29. Agha, Z., D.L. Roter, and R.M. Schapira, *An evaluation of patient-physician communication style during telemedicine consultations*. J Med Internet Res, 2009. **11**(3): p. e36.
- 30. Liu, X., et al., *Doctor-Patient communication: a comparison betweenTelemedicine consultation and face-to-face consultation*. Internal Medicine, 2007. **46**(5): p. 227-232.
- 31. Tates, K., et al., *The effect of screen-to-screen versus face-to-face consultation on doctorpatient communication: an experimental study with simulated patients.* Journal of medical Internet research, 2017. **19**(12): p. e8033.
- 32. Street Jr, R.L., E.J. Wheeler, and W.T. McCaughan, *Specialist–primary care provider–patient communication in telemedical consultations.* Telemedicine Journal, 2000. **6**(1): p. 45-54.
- Chua, I.S., V. Jackson, and M. Kamdar, Webside manner during the COVID-19 pandemic: maintaining human connection during virtual visits. Journal of Palliative Medicine, 2020.
   23(11): p. 1507-1509.
- 34. Walther, J.B., *Interpersonal effects in computer-mediated interaction: A relational perspective.* Communication research, 1992. **19**(1): p. 52-90.
- 35. Walther, J.B., T. Loh, and L. Granka, *Let me count the ways: The interchange of verbal and nonverbal cues in computer-mediated and face-to-face affinity.* Journal of language and social psychology, 2005. **24**(1): p. 36-65.
- 36. ACH. Academy of Communication in Healthcare. 2020 [cited 2022 19 Dec]; Available from: https://achonline.org/COVID-19/Telemedicine.
- Begasse de Dhaem, O. and C. Bernstein, *Headache virtual visit toolbox: the transition from bedside manners to webside manners.* Headache: The Journal of Head and Face Pain, 2020.
   60(8): p. 1743-1746.
- 38. BJGP. Video consultations. guide for practice. 2020 [cited 2022 19 Dec]; by Trisha Greenhalgh, on behalf of the IRIHS research group at the University of Oxford, with input from Clare Morrison of Scottish Government Technology Enabled Care Programme and Professor Gerald Koh Choon Huat from National University of Singapore]. Available from: <u>https://bjgplife.com/video-consultations-guide-for-practice/</u>.
- 39. Holstead, R.G. and A.G. Robinson, *Discussing serious news remotely: navigating difficult conversations during a pandemic.* JCO Oncology Practice, 2020. **16**(7): p. 363-368.

- 40. Houchens, N. and R. Tipirneni, *Compassionate communication amid the COVID-19 pandemic.* Journal of Hospital Medicine, 2020. **15**(7): p. 437-439.
- 41. Gunner, C., I. Thomas, and C. Morrison. *Video consultation skills: an overview for clinicians*.
   2020 [cited 2022 21 Dec]; Available from: <u>https://www.youtube.com/watch?v=QQ5V1-OoY1g</u>.
- 42. Jiménez-Rodríguez, D., et al. *Consensus on Criteria for Good Practices in Video Consultation: A Delphi Study*. International Journal of Environmental Research and Public Health, 2020. **17**, DOI: 10.3390/ijerph17155396.
- 43. Confederation, S., *Federal Act on Research involving Human Beings*, T.F.A.o.t.S. Confederation, Editor. 2014: CH.
- 44. StataCorp. *Stata Statistical Software*. [15] 2017.
- 45. Blanch-Hartigan, D., et al., *Measuring nonverbal behavior in clinical interactions: A pragmatic guide.* Patient education and counseling, 2018. **101**(12): p. 2209-2218.
- 46. Gorawara-Bhat, R. and M.A. Cook, *Eye contact in patient-centered communication*. Patient education and counseling, 2011. **82**(3): p. 442-447.
- 47. Helou, S., et al., *Physician eye contact in telemedicine video consultations: A cross-cultural experiment.* International Journal of Medical Informatics, 2022. **165**: p. 104825.
- 48. Silverman, J., S. Kurtz, and J. Draper, *Skills for communicating with patients*. 2016: crc press.
- 49. Shaw, S.E., et al., *Video consultations between patients and clinicians in diabetes, cancer, and heart failure services: linguistic ethnographic study of video-mediated interaction.* Journal of medical Internet research, 2020. **22**(5): p. e18378.
- 50. Egorskaya, A. and E. Zarnitsyna, *Improving communication skills in medical practice by mastering foreign languages*. Международный журнал гуманитарных и естественных наук, 2020(7-3): p. 72-74.
- 51. Armfield, N.R., et al., *Humour sans frontieres: the feasibility of providing clown care at a distance.* Telemedicine and e-Health, 2011. **17**(4): p. 316-318.
- 52. Demiris, G., S.M. Speedie, and S. Finkelstein, *Change of patients' perceptions of TeleHomeCare*. Telemedicine Journal and e-Health, 2001. **7**(3): p. 241-248.
- 53. Ehlers, D.K., J.L. Huberty, and G.-J. de Vreede, *Can an evidence-based book club intervention delivered via a tablet computer improve physical activity in middle-aged women?* Telemedicine and e-Health, 2015. **21**(2): p. 125-131.
- 54. Westra, I. and F. Niessen, *Implementing real-time video consultation in plastic surgery*. Aesthetic plastic surgery, 2015. **39**(5): p. 783-790.
- 55. Miller, E.A., *The technical and interpersonal aspects of telemedicine: effects on doctor– patient communication.* Journal of telemedicine and telecare, 2003. **9**(1): p. 1-7.
- 56. Stommel, W., C. Licoppe, and M. Stommel, *Difficult to assess in this manner. An'ineffective'showing sequence in post-surgery video consultation.* 2020.
- 57. Almathami, H.K.Y., K.T. Win, and E. Vlahu-Gjorgievska, *Barriers and Facilitators That Influence Telemedicine-Based, Real-Time, Online Consultation at Patients' Homes: Systematic Literature Review.* J Med Internet Res, 2020. **22**(2): p. e16407.
- 58. Greenhalgh, T., et al., *Infrastructure Revisited: An Ethnographic Case Study of how Health Information Infrastructure Shapes and Constrains Technological Innovation.* J Med Internet Res, 2019. **21**(12): p. e16093.
- 59. Speedtest. *Median Country Speeds November 2022*. 2022 [cited 2022 21 Dec]; Speedtest Global Index]. Available from: <u>https://worldpopulationreview.com/country-rankings/internet-speeds-by-country</u>.
- 60 Schmuckler, M. A. (2001). What is ecological validity? A dimensional analysis. Infancy, 2(4), 419-436.
- 61. Essers, G., et al., *Identifying context factors explaining physician's low performance in communication assessment: an explorative study in general practice.* BMC family practice, 2011. **12**(1): p. 1-8.

62. Salmon, P. and B. Young, *Creativity in clinical communication: from communication skills to skilled communication.* Medical education, 2011. **45**(3): p. 217-226.

# Appendices

# Table 1 Expert-based, best practice recommendations regarding video consultations

Expert-based recommendations	References
Recommendations with consensus	0
Ensure professional backdrop, appropriate environment	[33, 37, 39, 41, 42]
Describe the setting and members present, ensure who	[36, 37, 39, 41, 42]
is present/hearing the discussion	
Presentations and confirmation of patient identity (if not	
known)	
Inquire about the patient's environment (interference,	[33, 36-38, 42]
etc)	
Small talk (initial small talk to put the patient at ease)	[33, 36, 42]
Announce transitions, use active listening skills	[38, 39, 41]
Address confidentiality issues and privacy concerns	[38, 39, 41]
Inform about the tool platform (recording or not, etc)	
Express nonverbal empathy (facial expression, etc.)	[33, 36, 37, 39, 41, 42]
Discuss modalities for next consultation, when to consult	[33, 36, 38, 39, 41, 42]
in presence	
Send summaries and documents by email or other	[37, 39]
electronic mean	
Recommendations lacking consensus	

Framing	Head and bust [33, 39]
	Face [36]
Eye gazing orientation	Camera [33, 37, 39, 41]
	Screen [38, 42]
Take pauses	brief pauses [33]
	Longer pauses [37, 39]

 Table 2 Physician, patient and consultation characteristics for all six-communication related

# behaviour

	Dimension	Physician	Simulated patient	Торіс
1	Framing	Female 50 yrs old	Male 63 yrs old	Transmission of high cholesterol blood values
2	Eye gazing	Male 47 yrs old	Male 35 yrs old	History taking of abdominal bloating and stress and work
3	Initial social talk (related or not to quality of connection sound and image)	Male 38 yrs old	Female 60 yrs old	Beginning of an encounter after 6 months without consultation
4	Confidentiality	Female 35 yrs old	Female 22 yrs old	Beginning of an encounter (afraid to be pregnant)
5	Pauses	Male 35 yrs old	Female 50 yrs old	Transmission of a positive faecal blood test
6	Nonverbal empathy	Female 42 yrs old	Female 77 yrs old	Patient distressed by her husband's hospitalization

# Table 3 Patients' sociodemographic data

	N (%)	
Age (yrs)		
< 25	98 (23.5)	
25-44	176 (42.2)	
45-64	122 (29.3)	
<u>&gt;</u> 65	21 (5.0)	
Gender		
Female	227 (54.4)	
Civil status		
Single	210 (50.4)	
Married	145 (34.8)	
Divorced-separated	57 (13.7)	
Widowed	5 (1.2)	
Nationality		
Swiss	233 (55.9)	
Europe	116 (27.8)	
South America	27 (6.5)	
Africa	23 (5.5)	
Asia	12 (2.9)	
North America	3 (0.7)	
Occupation	$\mathbf{\alpha}$	
Without profession	16 (3.8)	
Student	99 (23.7)	
Employed	217 (52.0)	
Independent	30 (7.2)	
Retired	22 (5.3)	
Unemployed	15 (3.6)	
Other	18 (4.3)	

Figure 1 Patients' preferences regarding physicians' video-specific communication behaviours (%)



### Table 4 Percentage of patients' reasons congruent with the intention regarding video

### consultation behaviour

	Behaviour	Total comments N	Congruent reasons N (%)	Examples
Nonverbal	Framing	113	26 (23)	More proximity, framing, framing more human, posture
	Eye gazing	82	4 (5)	Eye gazing
	Pauses	138	70 (51)	Less pauses/interruptions, less hesitant, quicker, more direct
	Empathy	122	98 (80)	Empathetic, implicated, more understanding, listening more
	Confidentiality	156	87 (56)	Physician check whether patient feels free to speak, make the patient feel comfortable, check whether patient is alone
Verbal	Environment	113	38 (34)	Checks whether the patient is comfortable, check the video sound, the environment

# Highlights

- Most patients preferred when physicians addressed confidentiality issues (reminder of nonrecording of data, checking that patient is alone) and inquired about the quality of the connection at the beginning of the consultation.
- Although patients were sensitive to increased expression of nonverbal empathy in response to the patient' distress, they did not favour longer pauses during a video consultation.
- Patients did not favour or were indifferent to the eye gazing towards the camera.
- These findings are in line with the video-related communication behaviours recommended by experts except for eye gazing and pause duration.