



Article scientifique

Article

2010

Published version

Open Access

This is the published version of the publication, made available in accordance with the publisher's policy.

Exercise and the skeleton: How it works and what it really does

Bonnet, Nicolas; Ferrari, Serge Livio

How to cite

BONNET, Nicolas, FERRARI, Serge Livio. Exercise and the skeleton: How it works and what it really does. In: IBMS boneKEy, 2010, vol. 7, n° 7, p. 235–248. doi: 10.1138/20100453

This publication URL: <https://archive-ouverte.unige.ch/unige:33137>

Publication DOI: [10.1138/20100453](https://doi.org/10.1138/20100453)

PERSPECTIVES

Exercise and the Skeleton: How It Works and What It Really Does

Nicolas Bonnet and Serge L. Ferrari

Service of Bone Diseases, Department of Rehabilitation and Geriatrics, Geneva University Hospital, Geneva, Switzerland

Abstract

Exercise and nutrition are among the most commonly advocated lifestyle measures to improve skeletal health. A variety of exercise regimens has emphasized the beneficial role of speed, strength, power, endurance and coordination in improving bone mass and structure, and the effects of these regimens vary depending on age and are maximized during childhood and adolescence. Exercise influences the skeleton primarily by its direct impact on bone and by improving muscle mass and strength, which exerts further strains on the skeleton. These strains are sensed by mechanoreceptors, primarily on osteocytes, whose nature remains incompletely understood but which ultimately transduce the mechanical signals into biological signals, a process that involves the Wnt- β -catenin canonical signaling pathway. This biological signal is able to trigger bone remodeling by directing osteoblast activity and osteoclastic resorption. The role of microcracks in the mechanotransduction pathway and in the initiation of bone repair remains to be elucidated. Moreover, exercise induces changes in circulating levels of hormones such as growth hormone (GH) and insulin-like growth factor (IGF)-1, which exert anabolic effects on muscle and bone. A better understanding of the potential and limitations of the effects of physical activity on the skeleton, and of the molecular mechanisms mediating these effects, will lead to the development of new bone anabolic agents. *IBMS BoneKEy*. 2010 July;7(7):235-248.

©2010 International Bone & Mineral Society

Introduction

Bone has two main functions as a metabolic and structural tissue. The metabolic demands on the skeleton are regulated largely through calciotropic hormones whereas the structural functions of bone are determined primarily by genetic factors and adaptation mechanisms to its loading environment. From an historical point of view, it is important to note that the work of Wolff concentrated on the principle that static stressing of bone provoked alterations in shape, whereas Wilhelm Roux was the first to propose in 1881 a dynamic concept of functional adaptation by which the skeleton optimizes its structure to meet mechanical demands (1). Frost then developed the concept of the "mechanostat" describing the piezoelectric properties of the bone and other signaling mechanisms through which bone strength is increased where it is most needed. The importance of

bone adaptation to exercise is such that osteoporosis can be defined as a failure of the adaptive response to maintain the structure needed to withstand daily loading (2).

The beneficial effects of exercise on the mechanical properties of bone tissue are determined by the structure (macroarchitecture and microarchitecture) and the intrinsic quality of bone (the latter involving mainly hydroxyapatite and collagen content, which will not be the focus of this review). The loads applied on bone tissue during exercise can be divided into four categories: axial compression, bending, shearing and twisting. Each of these forces will affect bone tissue differently. For example, compression applies a homogenous strain on the surface of a transverse section of bone whereas twisting applies a high strain on the external surface but no significant forces on the internal

surface of bone. During exercise there is a combination of these different forces, making it difficult to predict the resulting effects on bone structure. However, several animal and human studies have shown that exercise influences both modeling and remodeling of the cortical and trabecular bone compartments. Loading increases periosteal deposition and slows endosteal resorption or causes endosteal deposition depending upon the skeletal location. Strains produce modeling mostly in the proximal region of limb bones (3;4), changing bones' geometry as evaluated by the total tissue volume, cortical or endocortical cross-sectional area, and moment of inertia. In rodents, exercise also increases bone volume fraction (BV/TV) as a result of increasing trabecular number and thickness, and decreasing trabecular separation (5-7). Similar effects have been reported in humans by measuring apparent trabecular bone parameters with QCT, pQCT or MRI (8;9). Finally, there is abundant evidence that mechanical properties are improved in response to loading, mostly through increases in periosteal apposition and particularly in the young (10;11) (see below).

Interestingly, bone turnover changes may tend towards bone formation in response to exercise of moderate intensity, but towards bone resorption with intensive training (12). In the latter case, Haversian (intracortical) bone remodeling could be increased, particularly in the distal region of the weight-bearing bones where stresses and strains are higher, creating microdamage. This in turn would induce osteocyte apoptosis that may initiate Haversian remodeling (13). The true nature of the message transmitted by osteocyte apoptosis is still unknown (this is one of the new interests in the field of mechanical bone signaling) (14). This remodeling process produces a transient increase in porosity, leaving the bone insufficiently strong to resist further strains produced by repeated loading, which would be a mechanism for stress fractures.

The Influence of Age on the Skeletal Response to Exercise in Humans

A quick overview of published data on exercise and bone in rodents and humans shows mainly beneficial effects (15;16). However a closer look reveals discordant results particularly depending on the age of practice (Fig. 1) (17). A number of published studies indicate that the most effective period during which to observe a larger increase of bone mineral mass with exercise is during adolescence, especially during the pre-pubertal years (18;19). Hence a prospective study in 120 children aged 8 years showed that 3 jumps of 61 cm per week over the course of 7 months doubled the percent change of bone mineral content (BMC) at the femoral neck and spine compared to a control group (20). Jumping was also effective in children aged 13 years in improving bone accrual in a sex-specific manner (21). Girls practicing gymnastics exhibit higher bone mineral density (BMD) and bone diameter at the midshaft tibia and femur compared to girls playing football (22). As summarized in a recent review, physical activity should include activities that generate high ground-reaction forces, such as jumping, skipping and running and last at least 7 to 20 months, to increase bone mass by about 2% at weight-bearing sites (femoral neck, spine) (23). Greater spinal BMC and apparent BMD (aBMD), as well as trabecular volumetric density and bone strength in the peripheral skeleton, are also observed in adult elite gymnasts (24), suggesting that the effects of exercise on bone mass growth are retained in adults, providing a true benefit as they reach perimenopause. However, it has been reported that more than half of the BMC gain (+3.5% after 7 months of exercise versus controls) induced by jumping in pre-pubertal boys and girls (25) was lost over 8 years, that is, even before these subjects reached peak bone mass. To understand this phenomenon, the 'Achilles tendon of the exercise effect,' it should be kept in mind that 80% of peak bone mass is genetically determined (26;27). Hence an initially greater bone mass (just after the intervention) may subsequently be remodeled to ultimately reach its genetically determined target.




Fig. 1. Schematic description of the effect of physical activity on bone mass throughout life. The red curve represents a continuous exercise effect; otherwise a punctuated effect of exercise could be attenuated over time. The yellow curve represents a sedentary person engaging in normal activities (such as walking). Adapted from Blanchet C, Chaire L, Chagnon A, Thibault G. *Activité physique et santé osseuse. Avis du comité scientifique de Kino-Québec*. Québec: Gouvernement du Québec, Ministère de l'Éducation, du Loisir et du Sport; 2008:1-42.

Nutrition may influence the effect of exercise in children and adults. Calcium supplements in prepubertal girls have been shown to interact with their level of physical activity on bone mass gain (28). Moreover, exercise had no effect in children with a low calcium intake (452.8 mg/day), a mild effect in those with moderate intake (762.9 mg/day) and a greater effect in those with a high calcium intake (1073 mg/day) (29). Similar observations have been reported for protein intake and exercise on BMD and microarchitecture in young boys (30), which could be explained by a dual stimulation of insulin-like growth factor (IGF)-1 production by proteins and exercise.

The effects of exercise in adults are less prominent. In a meta-analysis pooling 14 studies, exercise resulted in a slightly higher BMD at the spine (weighted mean difference 0.016 g/cm², CI, 0.005 to 0.027, p=0.005) (31). In older populations there are no published, randomized, prospective studies demonstrating positive effects of exercise on BMD. The lack of clear exercise effects in this population may have several

explanations. First is the difficulty in the elderly of practicing exercise with an impact of sufficient intensity to have a bone-forming effect (see below) (32). Moreover, a lack of motivation and fatigue limit compliance and the effectiveness of conventional exercise training. Second, lower muscle mass and strength (sarcopenia) in the elderly limits the strain exerted on the skeleton (33). Finally, the number of osteocytes and/or their sensitivity to mechanical strain may decline with age. The osteogenic potential of mesenchymal stem cells also declines with age as a result of changing expression of Wnts and PPARs, with a related increase of bone marrow adipogenicity (34;35). Nevertheless, despite no positive effect on BMD, several studies have demonstrated a reduction of fracture rate in elderly people practicing an activity, probably by preventing falls (36-38).

Mechanisms of Exercise-Related Bone Modeling/Remodeling

Exercise influences the skeleton by three main mechanisms: a direct impact on bone

that is translated into biological signals by mechanoreceptors; an indirect impact by improving muscle mass and strength that secondarily stimulates these

mechanoreceptors; and an indirect impact by inducing changes in hormone levels (such as calciotropic hormones, leptin, etc.) and local factors (Fig 2).




Fig. 2. Schematic description of the effect of mechanical stimuli on the skeleton through structural adaptation of bone and the maintenance of physiological homeostasis by employing the mineral reservoir embedded in the bone structure. During exercise load is transmitted to the skeleton through direct stimulation of the bone mechanosensor and by indirect stimulation through dynamic muscle activity. Hormones from fat and the liver modulate loading by affecting bone and muscle growth as well as muscle performance, and act indirectly through potential changes in the mineral reservoir. Recent data also underscore a paracrine role of muscle factors on bone adaptation in response to loading as well as the importance of neurons' capacity to change their function, chemical profile, or structure in response to mechanical stimuli, improving skeletal modeling/remodeling responses. Adapted from Sievänen H. Hormonal influences on the muscle-bone feedback system: a perspective. *J Musculoskelet Neuronal Interact.* 2005 Jul-Sep;5(3):255-61.

Direct effects of exercise on bone

As recently reviewed by Rizzoli *et al.* (23), most of the exercises demonstrating a significant effect on bone mass acquisition are those with impact, whereas exercises such as swimming or bicycling exert no significant effects on bone. For this reason, studies on bone adaptation to exercise performed in the past 30 years have focused on the effects of direct mechanical stimuli on bone, which has led to three main observations. First, bone adaptive responses require dynamic rather than static

mechanical stimuli. Thus bone formation increases with the degree of bone deformation, however, up to a certain level of deformation the bone formation response plateaus (39). This is well-illustrated in animal studies using axial compression on the ulna or tibia showing that this relationship is not linear but rather more sigmoid in nature (40), and in human athletes. In the latter instance, those who perform triple jumps generating 20 G of force have higher limb BMD than gymnasts (10 G) and runners (3 G) (41). In a recent study, volumetric BMD of the proximal tibia

increased with running distance to reach a plateau over 30 km/week, and was positively associated with peak acceleration (vertical acceleration expressed in G) in athletes running over 30 km/week (42). The second major observation is that extending the duration of skeletal loading does not yield proportional increases in bone mass as it does with the frequency of exercise. As loading duration is increased, the bone formation response tends to fade as the osteocytes and/or osteoblasts sensing the signals become desensitized. As a corollary, adaptive bone responses would be improved with brief, intermittent exercise. The third major observation is that the orientation and magnitude of stress on the skeleton during exercise must differ from the usual pattern of bone loading in order to produce a maximal stimulus on bone formation. This has been well-described by Lanyon *et al.* who write that “[t]he mechanically adaptive response is dominated not by the numerous cycles of ‘normal’ strain change engendered during the predominant activity but rather by far fewer cycles of relatively ‘abnormal’ strain changes produced during unusual loading situations” (43).

Mechanisms of mechanotransduction

Osteocytes, which represent up to 90% of all bone cells, seem to be the most appropriate cells to sense the magnitude and direction of mechanical strain: they are broadly distributed through both trabecular and cortical compartments, embedded in the calcified tissue and well-interconnected to other osteocytes, osteoblasts and osteoclasts through a network of canaliculi. Although it is widely accepted that the osteocyte is the cell responsible for sensing mechanical strain, there is a debate within the field as to whether or not this is detected at the cellular level and how this mechanical transduction is carried out. Osteocytes probably do not respond directly to mechanical strains, but rather respond to extracellular fluid waves in the osteocytic lacunae and canalicular system generated by the loading stimuli (44). Mechanoreceptors (strain-generated potential) present on osteocyte dendrites

would in turn generate small changes in electrical charges, activate calcium channels, and/or stimulate the release of molecular mediators such as prostaglandin (through the activation of COX2) and nitric oxide (NO) that further communicate the mechanical signal to osteoblasts at the periosteum, and to osteoblasts and osteoclasts intracortically and at endocortical surfaces (45). Another possibility is that osteocytes, mesenchymal stem cells and/or osteoblast lining cells sense strain at their plasma membranes through stretch-activated ion channels that permit calcium flux, potentially initiating other intracellular responses (46) (Fig. 3). These biochemical signals have two primary functions: activation of transcriptional mechanisms regulating bone modeling/remodeling, and propagation of mechanical information to other osteocytes through paracrine effects (by NO and PGE2 secretion, for instance) and transmission of other chemical signals by gap junctions (2;47). Recent evidence in mice and from *in vitro* studies indicates that Wnt-LRP (48-50) signaling plays a major role in mediating mechanical loading (51). Mechanical strains in osteoblasts induce a rapid, transient accumulation of active β -catenin in the cytoplasm and its translocation to the nucleus inducing target genes such as COX2 (52). Interestingly, the Wnt- β -catenin pathway also seems important in the mechanotransduction performed by the osteocyte. Bonewald *et al.* have demonstrated that fluid flow shear stress treatment of MLO-Y4 osteocytes results in increased phosphorylation of GSK-3 β , β -catenin nuclear translocation, and changes in the expression of β -catenin target genes (such as those for sclerostin and dickkopf 1) (53). Other signaling pathways activated in response to mechanical loading, such as those involving Akt, may also crosstalk with the Wnt- β -catenin pathway (54). Consistent with these *in vitro* studies, *in vivo* loading of the ulna of the TOPGAL mouse results in activation of β -catenin signaling in osteocytes 1 hour after a single loading, whereas it was detectable on the bone surface only after 24 hours, suggesting an




Fig. 3. Mechanotransduction. Osteocytes, osteoblasts, mesenchymal stem cells and many other cells have the capacity to respond to mechanostimulators (fluid shear stress, pressure, electric fields and tissue strain). Several biological components have been proposed to act as mechanosensors: **a**. Stretch-activated ion channels in the plasma membrane open in response to membrane strain and allow the influx of calcium and other ions. **b**. Single cilia or the glycocalyx, a layer of carbohydrate-rich proteins on the cell surface, can mediate mechanotransduction signaling in response to fluid shear stress by inducing the influx of calcium or increased intracellular cAMP, inducing COX2 and osteopontin signaling. **c**, **d**. Cell-cell junctional receptors or extracellular matrix (ECM)-cell focal adhesions allow cells to probe their environments, as is the case for the complex integrin-actin filaments that can activate several signaling pathways such as mitogen-activated protein kinase (MAPK)-focal adhesion kinase (FAK). **e**. Force-induced unfolding of ECM proteins, such as fibronectin, can initiate mechanotransduction signaling outside the cell. **f**. Intracellular strain can induce conformational changes in cytoskeletal elements such as filaments, crosslinkers or motor proteins, thereby changing binding affinities to specific molecules and activating signaling pathways such as the caveolin known to activate G protein signaling. **g**. The nucleus itself has been proposed to act as a mechanosensor. Intracellular deformations can alter chromatin conformation and modulate access to transcription factors or transcriptional machinery. **h**. Compression of the intercellular space can alter the effective concentration of autocrine and paracrine signaling molecules, such as nitric oxide (NO) and PGE2. In addition, changes in seven transmembrane domain G protein-coupled receptor (GPCR), lipid fluidity and even mitochondrial activity have been proposed as mechanosensors. These effects are often mediated through multiple, overlapping and crosstalking signaling pathways. The highlights of such cascades include the three modules of the MAPK family underscored by the activation of Ras, the Janus-activated kinase (JAK)/signal transducer and activator of transcription (STAT) pathway, Rac activation, calcium (Ca^{2+}) and NO signaling. The convergence of these pathways results in the activation of select transcription factors such as β -catenin implicated in the Wnt/LRP pathway, the most important pathway both in osteoblasts and osteocytes. Adapted from Jaalouk DE, Lammerding J. Mechanotransduction gone awry. *Nat Rev Mol Cell Biol.* 2009 Jan;10(1):63-73, with permission from Macmillan Publishers Ltd.

earlier stimulation of canonical Wnt signaling in the osteocyte rather than in the osteoblast. It should also be noted that osteocyte-specific β -catenin-deficient mice are characterized by low bone mass (55). Fig. 3 illustrates the many mechanotransduction mechanisms possibly activated in response to exercise.

Skeletal effects of exercise through muscle

The strong association between muscle mass/strength and bone anabolism and catabolism during growth, development and aging highlights the important interaction of these two systems to optimize locomotion (56). The fact that peak rates of bone mineral acquisition are preceded by peak rates of muscle mass gain, whereas sarcopenia later in life is accompanied by bone loss, supports the idea that exercise programs aiming to improve muscle mass and strength would also stimulate bone formation (57). Several exercise programs, therefore, have been developed to stimulate both muscle and bone anabolism in postmenopausal women (58). It appears that high-intensity resistance training that improves strength and muscle mass also improves bone mass (59). Muscle contraction can activate bone mechanoreceptors (Fig. 2), particularly in the periosteum where tendons are attached. In fact, 70% of the tension exerted on bone depends on muscle contraction, which is therefore more important than body weight itself as a mechanical stimulus (as further demonstrated by bone loss during immobilization or paralysis) (33). This property is well-illustrated by the phenotype of myostatin-deficient mice, which present myofiber hypertrophy, high BMD in the spine and limbs as well as increased bone strength, mainly due to changes in bone geometry primarily at sites of muscle insertion. In contrast there were no changes in shape or cross-sectional area of the midshaft femur (60;61). Interestingly, the frequency of skeletal muscle contraction is higher (15-30 Hz) than the frequency of direct mechanical stimuli on bone (0-15 Hz), raising the possibility of synergistic effects on bone of exercises combining impact and

muscle contraction (62;63). Again, this has been well-illustrated in mice lacking myostatin and performing exercise on a treadmill, in which case bone strength (represented by ultimate force, toughness and ultimate strain) was 25% higher than in normal mice (60). However the bone-muscle unit paradigm has several exceptions, such as in cycling or swimming where the bone response to exercise is lower compared to other activities, suggesting that there is less bone formation when gravitational forces are lower. For that reason no study has yet demonstrated unequivocally that either gravitational or muscle forces provide the dominant anabolic and anti-catabolic stimuli of exercise.

A potentially interesting new mechanism regulating the relationship between muscle and bone involves the nuclear receptor protein peroxisome proliferator-activated receptors (PPARs) β/δ . It has been shown that a PPAR β/δ agonist mimics the effects of exercise on muscle by increasing its oxidative metabolism and the switch of type II fast fibers responsible for rapid movements to type I slow fibers responsible for sustained activity (64-66). Transgenic mice over-expressing PPAR β/δ run faster and longer without myofiber hypertrophy and have been called 'marathon mice.' Preliminary data from our laboratory suggest that PPAR β/δ -deficient mice exhibit a low bone mass phenotype in relation to their low lean mass. Recently it was also shown that muscle can secrete growth factors such as IGF-1 and FGFs that would exert a paracrine role on periosteal cells (67). Muscle has also been suggested to contain a reserve of potential osteoprogenitor cells (68).

Exercise effects on bone through hormones

Several studies have shown that an acute bout of exercise can increase concentrations of anabolic hormones, such as GH/IGF-1, and FSH/LH/estrogen, across a wide age range (69). In particular, studies of estrogen by Lance Lanyon's group have provided evidence for a critical role of the estrogen

receptor in the mechanotransduction pathway (70;71). These hormonal changes are amplified by exercise patterns and intensity (see above).

Exercise modifies liver production of IGF-1 by modifying the flow of GH secreted by the pituitary gland. The rise in the amplitude and frequency of GH/IGF-1 production is explained by the stressing effects of exercise on serum glucose levels (hypoglycemia) (72;73). IGF-1 is also locally secreted by bone and muscle cells in response to the activation of mechanoreceptors (74) and IGF-I expression is increased in osteocytes and bone-lining cells within 6 hours of mechanical loading (75). Conversely, the low bone formation seen with disuse is due in part to a resistance to the effects of IGF-I on bone formation (76).

Parathyroid hormone (PTH) levels are also increased after maximal exercise independently of sex, age and training status (77-79). During physical exercise, the increase in PTH is due mainly to the calciuric effects of exercise, but other factors such as catecholamines and acidosis can also modify the secretion of PTH (80). One hypothesis is that the intermittent release of PTH produced by exercise is a systemic mediator of the anabolic actions on bone tissue (81). It was reported recently that variations in the circulating level of calciotropic hormones (PTH, vitamin D metabolites, and calcitonin) related to physical activity may modulate the bone tissue response to exercise (82). This is consistent with experimental evidence that administration of PTH before mechanical stimulation significantly increases the osteogenic response (83-85). This could be mediated by a PTH-stimulated production of PGE2 (86), and an increase in the activation of intracellular calcium, an important second messenger in the mechanotransduction cascade (see above) (83).

Adipokines, such as leptin, and enterokines, such as ghrelin, are also regulated by exercise (87;88). The role of the sympathetic nervous system, primarily β 2 adrenergic receptor-mediated signaling, in the

regulation of bone turnover has also been reported (89-91), as has the role of other neuroendocrine factors (glutamate, NPY, AgRp) (92;93). Our preliminary observations in mice lacking the β 2 adrenergic receptor indicate that their skeletal response to mechanical loading is unaltered, whereas the lack of the β 1 adrenergic receptor attenuates the trabecular and cortical response to loading (94). The influence of adrenergic tone on bone metabolism in response to exercise, however, will be complex in consideration of adrenergic effects on fat tissue and metabolism (lipolysis) (89;95;96). An extensive review of β -adrenergic effects on bone and response to exercise has been published in recent years (89;95).

Bone vascularization has also been implicated in exercise effects on bone structure. Moreover, the skeletal response to treadmill exercise was blocked by an anti-VEGF neutralizing antibody (97), suggesting that neo-angiogenesis is indispensable for bone gain in these conditions.

In summary, exercise can effectively improve bone mass, structure and strength, provided it is administered early in life, and with sufficient loading intensity and frequency. Its ultimate effects on the prevention of osteoporosis, however, could be quite limited. A better understanding of the complex pathways transducing mechanical stimulation into bone formation, particularly at the muscle-bone functional unit, is leading to the development of novel pharmacological approaches, such as SARMs, myostatin antagonists and sclerostin antibodies, which in turn could mimic the effects of exercise on the skeleton.

Conflict of Interest: Dr. Ferrari reports that he receives research support from Amgen and Merck Sharp & Dohme, and is an advisory committee member and lectures occasionally at conference symposia for the Alliance for Better Bone Health (sanofi aventis/P&G), Amgen, Merck Sharp & Dohme, Eli Lilly, Servier, and Novartis. Dr. Bonnet: none reported.

Peer Review: This article has been peer-reviewed.

References

1. Lee TC, Taylor D. Bone remodelling: should we cry Wolff? *Ir J Med Sci.* 1999 Apr-Jun;168(2):102-5.
2. Rubin J, Rubin C, Jacobs CR. Molecular pathways mediating mechanical signaling in bone. *Gene.* 2006 Feb 15;367:1-16.
3. Pearson OM, Lieberman DE. The aging of Wolff's "law": ontogeny and responses to mechanical loading in cortical bone. *Am J Phys Anthropol.* 2004;Suppl 39:63-99.
4. Lieberman DE, Pearson OM, Polk JD, Demes B, Crompton AW. Optimization of bone growth and remodeling in response to loading in tapered mammalian limbs. *J Exp Biol.* 2003 Sep;206(Pt 18):3125-38.
5. Bourrin S, Palle S, Genty C, Alexandre C. Physical exercise during remobilization restores a normal bone trabecular network after tail suspension-induced osteopenia in young rats. *J Bone Miner Res.* 1995 May;10(5):820-8.
6. Joo YI, Sone T, Fukunaga M, Lim SG, Onodera S. Effects of endurance exercise on three-dimensional trabecular bone microarchitecture in young growing rats. *Bone.* 2003 Oct;33(4):485-93.
7. Bonnet N, Beaupied H, Vico L, Dolleans E, Laroche N, Courteix D, Benhamou CL. Combined effects of exercise and propranolol on bone tissue in ovariectomized rats. *J Bone Miner Res.* 2007 Apr;22(4):578-88.
8. MacIntyre NJ, Adachi JD, Webber CE. In vivo detection of structural differences between dominant and nondominant radii using peripheral quantitative computed tomography. *J Clin Densitom.* 1999 Winter;2(4):413-22.
9. Ducher G, Tournaire N, Meddahi-Pellé A, Benhamou CL, Courteix D. Short-term and long-term site-specific effects of tennis playing on trabecular and cortical bone at the distal radius. *J Bone Miner Metab.* 2006;24(6):484-90.
10. Bass SL, Saxon L, Daly RM, Turner CH, Robling AG, Seeman E, Stuckey S. The effect of mechanical loading on the size and shape of bone in pre-, peri-, and postpubertal girls: a study in tennis players. *J Bone Miner Res.* 2002 Dec;17(12):2274-80.
11. Bass SL. The prepubertal years: a uniquely opportune stage of growth when the skeleton is most responsive to exercise? *Sports Med.* 2000 Aug;30(2):73-8.
12. Bourrin S, Genty C, Palle S, Gharib C, Alexandre C. Adverse effects of strenuous exercise: a densitometric and histomorphometric study in the rat. *J Appl Physiol.* 1994 May;76(5):1999-2005.
13. Cardoso L, Herman BC, Verborgt O, Laudier D, Majeska RJ, Schaffler MB. Osteocyte apoptosis controls activation of intracortical resorption in response to bone fatigue. *J Bone Miner Res.* 2009 Apr;24(4):597-605.
14. Noble BS, Stevens H, Loveridge N, Reeve J. Identification of apoptotic changes in osteocytes in normal and pathological human bone. *Bone.* 1997 Mar;20(3):273-82.
15. Nybo L, Sundstrup E, Jakobsen MD, Mohr M, Hornstrup T, Simonsen L, Bülow J, Randers MB, Nielsen JJ, Aagaard P, Krstrup P. High-intensity training vs. traditional exercise interventions for promoting health. *Med Sci Sports Exerc.* 2010 Feb 26. [Epub ahead of print]
16. Hind K, Burrows M. Weight-bearing exercise and bone mineral accrual in children and adolescents: a review of controlled trials. *Bone.* 2007 Jan;40(1):14-27.

17. Borer KT. Physical activity in the prevention and amelioration of osteoporosis in women: interaction of mechanical, hormonal and dietary factors. *Sports Med.* 2005;35(9):779-830.
18. Kannus P, Haapasalo H, Sankelo M, Sievänen H, Pasanen M, Heinonen A, Oja P, Vuori I. Effect of starting age of physical activity on bone mass in the dominant arm of tennis and squash players. *Ann Intern Med.* 1995 Jul 1;123(1):27-31.
19. Nordström P, Pettersson U, Lorentzon R. Type of physical activity, muscle strength, and pubertal stage as determinants of bone mineral density and bone area in adolescent boys. *J Bone Miner Res.* 1998 Jul;13(7):1141-8.
20. Fuchs RK, Bauer JJ, Snow CM. Jumping improves hip and lumbar spine bone mass in prepubescent children: a randomized controlled trial. *J Bone Miner Res.* 2001 Jan;16(1):148-56.
21. Weeks BK, Young CM, Beck BR. Eight months of regular in-school jumping improves indices of bone strength in adolescent boys and girls: the POWER PE study. *J Bone Miner Res.* 2008 Jul;23(7):1002-11.
22. Pettersson U, Nordström P, Alfredson H, Henriksson-Larsén K, Lorentzon R. Effect of high impact activity on bone mass and size in adolescent females: A comparative study between two different types of sports. *Calcif Tissue Int.* 2000 Sep;67(3):207-14.
23. Rizzoli R, Bianchi ML, Garabédian M, McKay HA, Moreno LA. Maximizing bone mineral mass gain during growth for the prevention of fractures in the adolescents and the elderly. *Bone.* 2010 Feb;46(2):294-305.
24. Eser P, Hill B, Ducher G, Bass S. Skeletal benefits after long-term retirement in former elite female gymnasts. *J Bone Miner Res.* 2009 Dec;24(12):1981-8.
25. Gunter K, Baxter-Jones AD, Mirwald RL, Almstedt H, Fuchs RK, Durski S, Snow C. Impact exercise increases BMC during growth: an 8-year longitudinal study. *J Bone Miner Res.* 2008 Jul;23(7):986-93.
26. Ferrari S, Rizzoli R, Slosman D, Bonjour JP. Familial resemblance for bone mineral mass is expressed before puberty. *J Clin Endocrinol Metab.* 1998 Feb;83(2):358-61.
27. Karasik D, Ferrari SL. Contribution of gender-specific genetic factors to osteoporosis risk. *Ann Hum Genet.* 2008 Sep;72(Pt 5):696-714.
28. Chevalley T, Rizzoli R, Hans D, Ferrari S, Bonjour JP. Interaction between calcium intake and menarcheal age on bone mass gain: an eight-year follow-up study from prepuberty to postmenarche. *J Clin Endocrinol Metab.* 2005 Jan;90(1):44-51.
29. Rowlands AV, Ingledew DK, Powell SM, Eston RG. Interactive effects of habitual physical activity and calcium intake on bone density in boys and girls. *J Appl Physiol.* 2004 Oct;97(4):1203-8.
30. Chevalley T, Bonjour JP, Ferrari S, Rizzoli R. High-protein intake enhances the positive impact of physical activity on BMC in prepubertal boys. *J Bone Miner Res.* 2008 Jan;23(1):131-42.
31. Martyn-St James M, Carroll S. Effects of different impact exercise modalities on bone mineral density in premenopausal women: a meta-analysis. *J Bone Miner Metab.* 2010 May;28(3):251-67.
32. American College of Sports Medicine position stand. Osteoporosis and exercise. *Med Sci Sports Exerc.* 1995 Apr;27(4):i-vii.
33. Kohrt WM, Bloomfield SA, Little KD, Nelson ME, Yingling VR; American

- College of Sports Medicine. American College of Sports Medicine Position Stand: physical activity and bone health. *Med Sci Sports Exerc.* 2004 Nov;36(11):1985-96.
34. Ozcivici E, Luu YK, Adler B, Qin YX, Rubin J, Judex S, Rubin CT. Mechanical signals as anabolic agents in bone. *Nat Rev Rheumatol.* 2010 Jan;6(1):50-9.
35. Huang SC, Wu TC, Yu HC, Chen MR, Liu CM, Chiang WS, Lin KM. Mechanical strain modulates age-related changes in the proliferation and differentiation of mouse adipose-derived stromal cells. *BMC Cell Biol.* 2010 Mar 10;11:18.
36. Korpelainen R, Keinänen-Kiukaanniemi S, Heikkinen J, Väänänen K, Korpelainen J. Effect of impact exercise on bone mineral density in elderly women with low BMD: a population-based randomized controlled 30-month intervention. *Osteoporos Int.* 2006 Jan;17(1):109-18.
37. Karlsson MK, Nordqvist A, Karlsson C. Physical activity, muscle function, falls and fractures. *Food Nutr Res.* 2008;52.
38. Sherrington C, Whitney JC, Lord SR, Herbert RD, Cumming RG, Close JC. Effective exercise for the prevention of falls: a systematic review and meta-analysis. *J Am Geriatr Soc.* 2008 Dec;56(12):2234-43.
39. Lanyon LE, Rubin CT. Static vs dynamic loads as an influence on bone remodelling. *J Biomech.* 1984;17(12):897-905.
40. Turner CH, Forwood MR, Otter MW. Mechanotransduction in bone: do bone cells act as sensors of fluid flow? *FASEB J.* 1994 Aug;8(11):875-8.
41. Benhamou CL, Ducher G. Activité physique et architecture osseuse. In: Hérisson C, Fardellone P, eds. *Os, Activité Physique et Ostéoporose*. Paris: Masson, Collection de Pathologie
- Locomotrice et de Médecine Orthopédique; 2005:60-68.
42. Deriaz O, Najafi B, Ballabeni P, Crettenand A, Gobelet C, Aminian K, Rizzoli R, Gremion G. Proximal tibia volumetric bone mineral density is correlated to the magnitude of local acceleration in male long distance runners. *J Appl Physiol.* 2010 Feb 4. [Epub ahead of print]
43. Lanyon LE. The success and failure of the adaptive response to functional loading-bearing in averting bone fracture. *Bone.* 1992;13:S17-S21.
44. Turner CH. Bone strength: current concepts. *Ann N Y Acad Sci.* 2006 Apr;1068:429-46.
45. Cowin SC, Moss ML. Mechanosensory mechanism in bone. In: Cowin SC, ed. *Bone Mechanics Handbook*. 2nd ed. Boca Raton, FL: CRC Press; 2001:29-1-29-17.
46. Davidson RM, Lingenbrink PA, Norton LA. Continuous mechanical loading alters properties of mechanosensitive channels in G292 osteoblastic cells. *Calcif Tissue Int.* 1996 Dec;59(6):500-4.
47. Jaalouk DE, Lammerding J. Mechanotransduction gone awry. *Nat Rev Mol Cell Biol.* 2009 Jan;10(1):63-73.
48. Choi HY, Dieckmann M, Herz J, Niemeier A. Lrp4, a novel receptor for Dickkopf 1 and sclerostin, is expressed by osteoblasts and regulates bone growth and turnover in vivo. *PLoS One.* 2009 Nov 20;4(11):e7930.
49. Balemans W, Van Hul W. The genetics of low-density lipoprotein receptor-related protein 5 in bone: a story of extremes. *Endocrinology.* 2007 Jun;148(6):2622-9.
50. Williams BO, Insogna KL. Where Wnts went: the exploding field of Lrp5 and

- Lrp6 signaling in bone. *J Bone Miner Res.* 2009 Feb;24(2):171-8.
51. Robinson JA, Chatterjee-Kishore M, Yaworsky PJ, Cullen DM, Zhao W, Li C, Kharode Y, Sauter L, Babij P, Brown EL, Hill AA, Akhter MP, Johnson ML, Recker RR, Komm BS, Bex FJ. Wnt/beta-catenin signaling is a normal physiological response to mechanical loading in bone. *J Biol Chem.* 2006 Oct 20;281(42):31720-8.
52. Case N, Ma M, Sen B, Xie Z, Gross TS, Rubin J. Beta-catenin levels influence rapid mechanical responses in osteoblasts. *J Biol Chem.* 2008 Oct 24;283(43):29196-205.
53. Bonewald LF, Johnson ML. Osteocytes, mechanosensing and Wnt signaling. *Bone.* 2008 Apr;42(4):606-15.
54. Fukumoto S, Hsieh CM, Maemura K, Layne MD, Yet SF, Lee KH, Matsui T, Rosenzweig A, Taylor WG, Rubin JS, Perrella MA, Lee ME. Akt participation in the Wnt signaling pathway through Dishevelled. *J Biol Chem.* 2001 May 18;276(20):17479-83.
55. Kramer I, Halleux C, Keller H, Pegurri M, Gooi JH, Weber PB, Feng JQ, Bonewald LF, Kneissel M. Osteocyte Wnt/beta-catenin signaling is required for normal bone homeostasis. *Mol Cell Biol.* 2010 Jun;30(12):3071-85.
56. Rittweger J. Ten years muscle-bone hypothesis: what have we learned so far? Almost a festschrift. *J Musculoskelet Neuronal Interact.* 2008 Apr-Jun;8(2):174-8.
57. Robling AG. Is bone's response to mechanical signals dominated by muscle forces? *Med Sci Sports Exerc.* 2009 Nov;41(11):2044-9.
58. Marini M, Sarchielli E, Brogi L, Lazzeri R, Salerno R, Sgambati E, Monaci M. Role of adapted physical activity to prevent the adverse effects of the sarcopenia. A pilot study. *Ital J Anat Embryol.* 2008 Oct-Dec;113(4):217-25.
59. Layne JE, Nelson ME. The effects of progressive resistance training on bone density: a review. *Med Sci Sports Exerc.* 1999 Jan;31(1):25-30.
60. Hamrick MW, Samaddar T, Pennington C, McCormick J. Increased muscle mass with myostatin deficiency improves gains in bone strength with exercise. *J Bone Miner Res.* 2006 Mar;21(3):477-83.
61. Hamrick MW, McPherron AC, Lovejoy CO, Hudson J. Femoral morphology and cross-sectional geometry of adult myostatin-deficient mice. *Bone.* 2000 Sep;27(3):343-9.
62. Burger EH, Klein-Nulend J, Cowin SC. Mechanotransduction in bone. *Advances in Organ Biology.* 1998;5:123-36.
63. Cowin SC. Mechanosensation and fluid transport in living bone. *J Musculoskelet Neuronal Interact.* 2002 Mar;2(3):256-60.
64. Narkar VA, Downes M, Yu RT, Emblar E, Wang YX, Banayo E, Mihaylova MM, Nelson MC, Zou Y, Juguilon H, Kang H, Shaw RJ, Evans RM. AMPK and PPARdelta agonists are exercise mimetics. *Cell.* 2008 Aug 8;134(3):405-15.
65. Wang YX, Zhang CL, Yu RT, Cho HK, Nelson MC, Bayuga-Ocampo CR, Ham J, Kang H, Evans RM. Regulation of muscle fiber type and running endurance by PPARdelta. *PLoS Biol.* 2004 Oct;2(10):e294.
66. Ehrenborg E, Krook A. Regulation of skeletal muscle physiology and metabolism by peroxisome proliferator-activated receptor delta. *Pharmacol Rev.* 2009 Sep;61(3):373-93.
67. Hamrick MW, McNeil PL, Patterson SL. Role of muscle-derived growth factors in

- bone formation. *J Musculoskeletal Neuronal Interact.* 2010 Mar;10(1):64-70.
68. Liu R, Peacock L, Schindeler A, Little D. Myogenic progenitors contribute to ectopic bone formation and open fracture repair. *J Bone Miner Res.* 2009;24(Suppl 1):SA0228.
69. Copeland JL, Consitt LA, Tremblay MS. Hormonal responses to endurance and resistance exercise in females aged 19-69 years. *J Gerontol A Biol Sci Med Sci.* 2002 Apr;57(4):B158-65.
70. Lanyon LE. Using functional loading to influence bone mass and architecture: objectives, mechanisms, and relationship with estrogen of the mechanically adaptive process in bone. *Bone.* 1996 Jan;18(1 Suppl):37S-43S.
71. Armstrong VJ, Muzylak M, Sinters A, Zaman G, Saxon LK, Price JS, Lanyon LE. Wnt/beta-catenin signaling is a component of osteoblastic bone cell early responses to load-bearing and requires estrogen receptor alpha. *J Biol Chem.* 2007 Jul 13;282(28):20715-27.
72. Nindl BC, Hymer WC, Deaver DR, Kraemer WJ. Growth hormone pulsatility profile characteristics following acute heavy resistance exercise. *J Appl Physiol.* 2001 Jul;91(1):163-72.
73. Wideman L, Weltman JY, Hartman ML, Veldhuis JD, Weltman A. Growth hormone release during acute and chronic aerobic and resistance exercise: recent findings. *Sports Med.* 2002;32(15):987-1004.
74. Adams GR. Autocrine and/or paracrine insulin-like growth factor-I activity in skeletal muscle. *Clin Orthop Relat Res.* 2002 Oct;(403 Suppl):S188-96.
75. Lean JM, Jagger CJ, Chambers TJ, Chow JW. Increased insulin-like growth factor I mRNA expression in rat osteocytes in response to mechanical stimulation. *Am J Physiol.* 1995 Feb;268(2 Pt 1):E318-27.
76. Sakata T, Wang Y, Halloran BP, Elalieh HZ, Cao J, Bikle DD. Skeletal unloading induces resistance to insulin-like growth factor-I (IGF-I) by inhibiting activation of the IGF-I signaling pathways. *J Bone Miner Res.* 2004 Mar;19(3):436-46.
77. Iwamoto J, Shimamura C, Takeda T, Abe H, Ichimura S, Sato Y, Toyama Y. Effects of treadmill exercise on bone mass, bone metabolism, and calciotropic hormones in young growing rats. *J Bone Miner Metab.* 2004;22(1):26-31.
78. Maimoun L, Simar D, Caillaud C, Coste O, Barbotte E, Peruchon E, Rossi M, Mariano-Goulart D. Response of calciotropic hormones and bone turnover to brisk walking according to age and fitness level. *J Sci Med Sport.* 2009 Jul;12(4):463-7.
79. Scott JP, Sale C, Greeves JP, Casey A, Dutton J, Fraser WD. The effect of training status on the metabolic response of bone to an acute bout of exhaustive treadmill running. *J Clin Endocrinol Metab.* 2010 Jun 2. [Epub ahead of print]
80. Bouassida A, Latiri I, Bouassida S, Zalleg D, Zaouali M, Feki Y, Gharbi N, Zbidi A, Tabka Z. Parathyroid hormone and physical exercise: a brief review. *J Sports Sci Med.* 2006;5:367-74.
81. Rudberg A, Magnusson P, Larsson L, Joborn H. Serum isoforms of bone alkaline phosphatase increase during physical exercise in women. *Calcif Tissue Int.* 2000 May;66(5):342-7.
82. Maïmoun L, Sultan C. Effect of physical activity on calcium homeostasis and calciotropic hormones: a review. *Calcif Tissue Int.* 2009 Oct;85(4):277-86.
83. Zhang J, Ryder KD, Bethel JA, Ramirez R, Duncan RL. PTH-induced actin depolymerization increases

- mechanosensitive channel activity to enhance mechanically stimulated Ca²⁺ signaling in osteoblasts. *J Bone Miner Res.* 2006 Nov;21(11):1729-37.
84. Kim CH, Takai E, Zhou H, von Stechow D, Müller R, Dempster DW, Guo XE. Trabecular bone response to mechanical and parathyroid hormone stimulation: the role of mechanical microenvironment. *J Bone Miner Res.* 2003 Dec;18(12):2116-25.
85. Sugiyama T, Saxon LK, Zaman G, Moustafa A, Sunters A, Price JS, Lanyon LE. Mechanical loading enhances the anabolic effects of intermittent parathyroid hormone (1-34) on trabecular and cortical bone in mice. *Bone.* 2008 Aug;43(2):238-48.
86. Bakker AD, Joldersma M, Klein-Nulend J, Burger EH. Interactive effects of PTH and mechanical stress on nitric oxide and PGE2 production by primary mouse osteoblastic cells. *Am J Physiol Endocrinol Metab.* 2003 Sep;285(3):E608-13.
87. Jürimäe J, Kums T, Jürimäe T. Adipocytokine and ghrelin levels in relation to bone mineral density in physically active older women: longitudinal associations. *Eur J Endocrinol.* 2009 Mar;160(3):381-5.
88. Prouteau S, Benhamou L, Courteix D. Relationships between serum leptin and bone markers during stable weight, weight reduction and weight regain in male and female judoists. *Eur J Endocrinol.* 2006 Mar;154(3):389-95.
89. Bonnet N, Courteix D, Benhamou CL. Leptin, central nervous system, and bone: influence of physical activity. *Joint Bone Spine.* 2005 Dec;72(6):477-80.
90. Kondo A, Togari A. In vivo stimulation of sympathetic nervous system modulates osteoblastic activity in mouse calvaria. *Am J Physiol Endocrinol Metab.* 2003 Sep;285(3):E661-7.
91. Levasseur R, Sabatier JP, Potrel-Burgot C, Lecoq B, Creveuil C, Marcelli C. Sympathetic nervous system as transmitter of mechanical loading in bone. *Joint Bone Spine.* 2003 Dec;70(6):515-9.
92. Mason DJ. Glutamate signalling and its potential application to tissue engineering of bone. *Eur Cell Mater.* 2004 Apr 7;7:12-25; discussion 25-6.
93. Szczesniak AM, Gilbert RW, Mukhida M, Anderson GI. Mechanical loading modulates glutamate receptor subunit expression in bone. *Bone.* 2005 Jul;37(1):63-73.
94. Bonnet N, Pierroz D, Ferrari S. Low bone mass and decreased biomechanical response in beta 1 adrenergic receptor KO but not in beta 2 adrenergic receptor KO Mice. *J Bone Miner Res.* 2008;23(Suppl 1):S41.
95. Bonnet N, Pierroz DD, Ferrari SL. Adrenergic control of bone remodeling and its implications for the treatment of osteoporosis. *J Musculoskelet Neuronal Interact.* 2008 Apr-Jun;8(2):94-104.
96. Blaak EE. Adrenergically stimulated fat utilization and ageing. *Ann Med.* 2000 Sep;32(6):380-2.
97. Yao Z, Lafage-Proust MH, Plouët J, Bloomfield S, Alexandre C, Vico L. Increase of both angiogenesis and bone mass in response to exercise depends on VEGF. *J Bone Miner Res.* 2004 Sep;19(9):1471-80.