



Article scientifique

Article

2021

Accepted version

Open Access

This is an author manuscript post-peer-reviewing (accepted version) of the original publication. The layout of the published version may differ .

---

A new mutation in *mgrb* mediating polymyxin resistance in *Klebsiella variicola*

---

Lenzi, Michael H; Martins, Willames Mbs; Roch, Mélanie; Ramos, Patrícia L; Sands, Kirsty; Cayô, Rodrigo; Walsh, Timothy R; Andrey, Diego Olivier; Gales, Ana C

**How to cite**

LENZI, Michael H et al. A new mutation in *mgrb* mediating polymyxin resistance in *Klebsiella variicola*. In: International journal of antimicrobial agents, 2021, vol. 58, n° 5, p. 106424. doi: 10.1016/j.ijantimicag.2021.106424

This publication URL: <https://archive-ouverte.unige.ch//unige:169899>

Publication DOI: [10.1016/j.ijantimicag.2021.106424](https://doi.org/10.1016/j.ijantimicag.2021.106424)

1  
2 **A New Mutation in *mgrB* Mediating Polymyxin Resistance in *Klebsiella variicola***

3  
4 Michael H. Lenzi<sup>1</sup>, Willames M. B. S. Martins<sup>1,2\*</sup>, Mélanie Roch<sup>3</sup>, Patrícia L. Ramos<sup>4</sup>,  
5 Kirsty Sands<sup>2,5</sup>, Rodrigo Cayô<sup>1</sup>, Timothy R. Walsh<sup>6</sup>, Diego O. Andrey<sup>3</sup>, Ana C. Gales<sup>1</sup>

6  
7  
8 <sup>1</sup>Division of Infectious Diseases, Department of Internal Medicine. Escola Paulista de  
9 Medicina/Universidade Federal de São Paulo (UNIFESP), São Paulo, Brazil.

10 <sup>2</sup>Department of Medical Microbiology, Division of Infection and Immunity, Cardiff University,  
11 Cardiff, United Kingdom.

12 <sup>3</sup>Service of Infectious Diseases, Geneva University Hospitals and Faculty of Medicine, Geneva,  
13 Switzerland.

14 <sup>4</sup>Departamento de Pesquisas Aplicadas, Fundação Parque Zoológico de São Paulo – FPZSP, São  
15 Paulo, Brazil.

16 <sup>5</sup>Department of Zoology, University of Oxford, United Kingdom.

17 <sup>6</sup>Ineos Oxford Institute of Antimicrobial Research, Department of Zoology, University of Oxford,  
18 United Kingdom.

19 **Short running title:** Polymyxin-resistant *Klebsiella variicola*.

20  
21  
22  
23 \*Corresponding Author: **Willames M. B. S. Martins**  
24 Rua Pedro de Toledo, 781, 6<sup>th</sup> Floor  
25 São Paulo - SP  
26 Brazil  
27 ZIP code: 04039-032  
28 Phone/Fax.: +55 11 55715180  
29 E-mail: [willamesbrasileiro@hotmail.com](mailto:willamesbrasileiro@hotmail.com)

**Abstract**

Polymyxin resistance is a public health concern, present in humans, animals and the environment, occasioned by chromosomal- or plasmid-encoding mechanisms. Chromosomal alterations in MgrB are frequently detected in *Klebsiella* spp., but not yet reported and characterised in *K. variicola*. Here, we performed microbiological and genomic characterisation of three polymyxin-resistant *K. variicola* isolates (M14, M15, and M50) recovered from the microbiota of migratory birds in Brazil. The isolates were submitted to SpeI-PFGE, broth microdilution, and Whole Genome Sequencing using Illumina MiSeq for analysis of genetic relatedness, sequence typing, and detection of antimicrobial resistance genes. *K. variicola* isolates belonged to two clones, and susceptibility tests showed resistance only for polymyxins. Sequences of chromosomal two-component systems (PmrAB, PhoPQ, RstAB, CrrAB) and MgrB were evaluated by blastN and blastP, against a polymyxin-susceptible *K. variicola* (A58243), and mutations biological effect was checked by the PROVEAN tool. In M14 and M15, *phoQ* mutations (D90N, I122S, and G385S) were identified, while in M50 a *mgrB* variant containing a single deletion (C deletion on position 93) leading to the production of a non-functional protein was detected. *mgrB* complementation studies showed restoration of polymyxin susceptibility (64 to  $\leq 0.25$  mg/L) when a WT *mgrB* was inserted into the *mgrB* deficient M50. This data confirmed the role of a non-functional *mgrB* variant in conferring polymyxin resistance, stressing the role of this regulator in *K. variicola* and drawing attention to novel polymyxin resistance mechanisms emerging in wildlife.

**Keywords:** Enterobacterales, Two-component systems, migratory birds, Antimicrobial Resistance

## 51        **1. Introduction**

52            Polymyxins are polypeptide antibiotics broadly used in human and veterinary medicine due  
53 to their great activity against Gram-negative bacteria, mainly against those pathogens with a  
54 multidrug-resistant (MDR) profile [1]. There have been reports of polymyxin-resistant  
55 *Enterobacteriales* recovered from distinct settings in recent years, either by chromosomal or plasmid-  
56 mediated mechanisms [1]. Plasmid-mediated mechanisms, represented by mobile colistin resistance  
57 (*mcr*) genes, are often reported in *Enterobacteriales*; however, they are more prevalent in food-animal  
58 samples than in humans and food products [2].

59            Chromosomal mechanisms leading to polymyxin resistance are closely associated with lipid  
60 A modification after a series of genetic and biochemical events coordinated by Two-Component  
61 Systems (TCS) such as PmrAB, PhoPQ, RstAB and CrrAB [3,4]. In 2014, the role of MgrB, a small  
62 protein responsible for negatively regulating PhoPQ activity, was described as the main mechanism  
63 driven polymyxin resistance in *Klebsiella pneumoniae* [5]. Since then, this mechanism has been  
64 worldwide reported [6-8] in *K. pneumoniae* and *K. oxytoca* [1,9], and more recently in *Enterobacter*  
65 spp. [10]. However, to date, no data associating polymyxin resistance to *mgrB* disruptions in *K.*  
66 *variicola* has been published. Herein, we have characterised three polymyxin-resistant *K. variicola*  
67 isolates recovered from the microbiota of migratory birds, ultimately mediated by chromosomal  
68 mutations.

69

## 70        **2. Material and Methods**

71

72        **2.1. Bacterial strains.** A surveillance study in 2012 aimed to identify antimicrobial-resistant  
73 bacteria carried by migratory and zoo resident birds in São Paulo, Brazil [11]. For this purpose, birds'  
74 choanal and cloacal swabs were collected and further plated on agar plates containing antimicrobial  
75 agents to favor the selection of resistant microorganisms. *K. variicola* isolates were recovered from

76 the choana of distinct *Dendrocygna viduata* birds following cultivation onto MacConkey agar  
77 containing 2 mg/L of polymyxin and further characterized in this study. The recovered isolates were  
78 identified by MALDI-TOF MS (Bruker Daltonics, Massachusetts, EUA) and 16S rRNA DNA-  
79 sequencing.

80

81

82 **2.2. Antimicrobial susceptibility testing and molecular typing.** Susceptibility profiles were  
83 determined by broth microdilution and interpreted according to EUCAST guidelines [12]. In addition,  
84 to check for the presence of polymyxin resistance mechanism-dependent of divalent ions such as  
85 MCR-like proteins, polymyxin B MICs were also determined in the presence of 10 mM of EDTA.  
86 Genetic relatedness was established by SpeI pulsed-field gel electrophoresis (PFGE) and interpreted  
87 using Tenover criteria [13].

88

89

90 **2.3. Whole Genome Sequence Analysis.** For genomic purposes, the isolates were whole genome  
91 sequenced using Illumina MiSeq (NexteraXTv2 and MiSeqReagent V3 Kit; 2×300cycles). Besides,  
92 a polymyxin-susceptible *K. variicola* strain (A58243) recovered from human-infection was also  
93 sequenced and used as a control. Raw sequences (fastq) were trimmed using Trim Galore (v0.5.0)  
94 and assembled into contigs using SPAdes (v3.9.0) [14]. The genomes were submitted to RAST  
95 (<http://rast.nmpdr.org/>) for automatic annotation followed by analysis on ResFinder  
96 (<https://cge.cbs.dtu.dk/services/ResFinder-3.0/>), PlasmidFinder  
97 (<https://cge.cbs.dtu.dk/services/PlasmidFinder-2.0/>), and *K. variicola* MLST website [15]. Mutation  
98 analysis of TCS (PmrAB, PmrD, PhoPQ, RstAB, CpxAR) and *mgrB* were conducted using as  
99 reference four genomes of *K. variicola* described as polymyxin-susceptible on NCBI (CP028555.1,  
100 CP016344.1, CP017289.1, and NZ\_CEGG01000025.1) and the strain *K. variicola* A58243

101 (polymyxin MIC 0.25 µg/mL). Mutation impact in proteins was determined by PROVEAN  
102 (<http://provean.jcvi.org/index.php>), using threshold values below to -2.5 as biologically significant.

103  
104  
105 **2.4. Expression level of TCS.** To ascertain the expression levels of *pmrA*, *pmrB*, *phoP*, *phoQ* and  
106 *mgrB* we performed qRT-PCR. Briefly, total RNA was isolated from *K. variicola* isolates using  
107 RNeasy Mini Kit (Qiagen, Hilden, Germany) with addition of RNase-free DNase (Qiagen, Hilden,  
108 Germany). Reverse transcription of the extracted RNA was performed using High Capacity cDNA  
109 Reverse Transcription Kit (Life Technologies, Carlsbad, CA, USA), followed by transcripts  
110 quantification, performed in triplicate using SYBR® Green PCR Master Mix (Life Technologies,  
111 Carlsbad, CA, USA) and the 7500 Real Time (Life Technologies, Carlsbad, CA, USA). For this  
112 purpose, 16S rRNA was used as endogenous control, and the levels of TCS and regulator gene  
113 expression were compared to *K. variicola* A58243.

114  
115 **2.5. mgrB complementation.** To evaluate the relationship between this new *mgrB* mutation and  
116 polymyxin resistance in *K. variicola*, we performed the *mgrB* complementation on M50 isolate.  
117 Briefly, an apramycin resistance gene was amplified and cloned into pBluescript SK+ XceI site  
118 (Stratagene Inc.) to build *pskA* vector. The gene *mgrB* was amplified from *Klebsiella variicola*  
119 A58243 (*mgrB*-WT) and M50 (*mgrB*-mutant) using the primers *mgrB*-BamHI-ext-R 5'-  
120 CGGGATCCCGAAGGCGTTCATTCTACCACC-3' EcoRI-*mgrB*-var-F 5'-  
121 GGAATTCCTTAAGAAGGCCGTGTTATCC-3' and cloned into *pskA*. Clones were selected on LB  
122 agar plates supplemented with apramycin (50 mg/L) and the constructions verified by sanger  
123 sequencing (Fasteris, Geneva, Switzerland). Polymyxin B broth microdilution was performed to  
124 quantify any modification on the MIC after *mgrB* complementation.

125

### 3. Results and Discussion

Three *K. variicola* (M14, M15, and M50) were recovered from the choana of distinct *Dendrocygna viduata* birds following cultivation onto MacConkey agar containing polymyxin B. The three isolates were subject to SpeI-PFGE and this analysis suggested that the isolates M14 and M15 were genetically related and classified as subtypes of the pattern A, A1 and A2. In contrast, M50 isolate belonged to the pattern B (Table 1). Interestingly, the isolates were susceptible to all antimicrobial agents tested except for polymyxin B (Table 1).

Genomic and microbiological features of all isolates were displayed in table 1. Isolates M14 and M15 belonged to ST137, while the isolate M50 was assigned to a new sequencing type denominated ST167. According to MLST database, isolates belonging to ST137 were previously identified in Germany causing human infections, highlighting this bacteria species' ability to exert symbiotic and pathogenic roles (<http://mlstkv.insp.mx/>). No plasmids were detected in these isolates, and only chromosomal resistance genes were detected. The gene encoding the intrinsic  $\beta$ -lactamase LEN was detected, *bla*<sub>LEN-24</sub> in M14 and M15 isolates, and *bla*<sub>LEN-13</sub> in M50 isolate. LEN-enzymes are chromosomally encoded and show a high level of similarity to SHV  $\beta$ -lactamases, which are also chromosomally encoded by *K. pneumoniae* [16].

Mutations in PmrB, PhoQ, and RstB were identified in two isolates; however, only mutations in PhoQ were considered deleterious by PROVEAN (Table 1). The deleterious mutation D90N was detected in M14 and M15 isolates. Interestingly, genetic modifications in the same TCS leading to polymyxin resistance were described previously in a *K. variicola* isolate from China, albeit not at the same amino-acid position [17]. However, the D90N substitution has not been described as causing polymyxin resistance in *K. pneumoniae*, although this highly conserved amino acid among Enterobacterales has been proposed as crucial for PhoQ function [18]. While the PhoQ G385S substitution has been described (although not functionally demonstrated) in *K. pneumoniae*, the I122 has not been described so far and its role remains to be demonstrated [19].

152 The level of transcription of TCS corroborated with the data obtained from genomic analysis,  
153 with increased transcriptional levels to *pmrB*, *phoP*, and *phoQ* in the M14 isolate, while in the M15  
154 isolate the transcriptional levels of *pmrB* and *phoQ* rose (Figure 1).

155 Although the isolate M50 did not show mutation on these TCS, a single nucleotide deletion  
156 (C deletion on position 93) was observed in *mgrB*. This deletion changed the amino acid residues  
157 sequence downstream of D31 (aspartic acid 31), thus encoding a non-functional MgrB protein of 52  
158 amino acids versus 47 amino acids for the wild-type MgrB (Figure 1). qRT-PCR experiments showed  
159 that the transcription level of the mutated *mgrB* was almost 5 times higher (4.8 times) than the  
160 susceptible isolate (A58243) and 4.6 times higher than other polymyxin-resistant isolates studied  
161 without mutation in *mgrB* (Figure 1a). To the best of our knowledge, this is the first description of  
162 polymyxin resistance being mediated by deletion of a single nucleotide in *mgrB*, resulting in an  
163 unfunctional protein. To date, polymyxin resistance mediated by *mgrB* has been frequently associated  
164 with incorporation of insertion sequences within the gene or its promoter or point mutations  
165 generating premature stop codons [1,3].

166 The complementation of M50 by a WT *mgrB* resulted in a drop of polymyxin B MIC from 64  
167 to  $\leq 0.25$  mg/L (Table 2), corroborating our initial suspicion on this mutation's role for conferring  
168 polymyxin resistance in *K. variicola*. The complementation with the deficient *mgrB* amplified from  
169 M50 isolate, did not show any impact in polymyxin B MIC, proving that this mutation caused the  
170 production of a non-functional MgrB protein (Table 2). Recently, six polymyxin-resistant *K.*  
171 *pneumoniae* strains were described carrying a duplication of 79 nucleotides in *mgrB*, resulting in an  
172 unfunctional MgrB, being 26 amino acids longer than expected [20]. These isolates were also  
173 polymyxin-resistant, and no other mechanism causing resistance to polymyxins was detected by WGS  
174 analysis [20]. This data further supports our findings since the production of an unfunctional MgrB  
175 protein is also detected in *K. pneumoniae*.



#### 177 **4. Conclusion**

178 *K. variicola* is an emerging human pathogen that should be monitored, especially regarding  
179 antimicrobial resistance and virulence determinants. Our results show that although *K. variicola*  
180 isolates are likely to be very susceptible to many antimicrobial agents, the susceptibility to  
181 polymyxins cannot be indirectly predicted. The identification of polymyxin-resistant *K. variicola* in  
182 migratory birds, also reinforces the need of constant effort on One Health surveillance programs since  
183 the close relationship between human and animals can facilitate the spread of such resistance  
184 determinants to hospital settings.

#### 185 186 187 188 189 190 **Data availability**

191 Whole genomic sequences of three studied polymyxin-resistant *K. variicola* isolates have  
192 been deposited in GenBank under accession numbers JAAGEJ000000000, JAAGEK000000000,  
193 JAAGEL000000000.

#### 194 195 **Conflict of interests**

196 A.C.G. has recently received research funding and/or consultation fees from Cristália,  
197 Enthasis Therapeutics, InfectoPharm, Eurofarma, Pfizer, MSD, and Zambon. Other authors have  
198 nothing to declare.

#### 199 200 **Funding**

201 This work received funds of The São Paulo Research Foundation (FAPESP), process number:  
202 2017/02258-6. M.H.L and A.C.G received are supported by The Brazilian National Council for  
203 Scientific and Technological Development (A.C.G. - 312066/2019-8), W. M. B. S. Martins is the  
204 recipient of Postdoctoral fellowship of The São Paulo Research Foundation (FAPESP 2018/24431-  
205 4), and D. O. A. was the recipient of a Swiss National Science Foundation Fellowship (APM

206 P300PB\_171601), a Sir Julius Thorn Foundation (Switzerland) research grant and a Geneva  
207 University Hospital research funding.

208

209

210 **Acknowledgements**

211 We thank the platform *Klebsiella variicola* Multi locus Sequence Typing (Instituto Nacional  
212 de Salud Pública) for coding MLST alleles. We also would like to thanks Ana Clara Narciso, Carolina  
213 Nodari, Carolina Yshihama, and Ye Liu Ou for their excellent technical contribution to this study.

214

215

216

**5. References**

- [1] Poirel L, Jayol A, Nordmann P. Polymyxins: Antibacterial Activity, Susceptibility Testing, and Resistance Mechanisms Encoded by Plasmids or Chromosomes. *Clin Microbiol Rev.* 2017;30(2):557–596. doi:10.1128/CMR.00064-16
- [2] Elbediwi M, Li Y, Paudyal N, Pan H, Li X, Xie S, et al. Global Burden of Colistin-Resistant Bacteria: Mobilized Colistin Resistance Genes Study (1980-2018). *Microorganisms.* 2019 Oct 16;7(10):461. doi: 10.3390/microorganisms7100461
- [3] Olaitan AO, Morand S, Rolain JM. Mechanisms of polymyxin resistance: acquired and intrinsic resistance in bacteria. *Front Microbiol.* 2014 Nov 26;5:643. doi: 10.3389/fmicb.2014.00643
- [4] Bhagirath AY, Li Y, Patidar R, Yerex K, Ma X, Kumar A, et al. Two Component Regulatory Systems and Antibiotic Resistance in Gram-Negative Pathogens. *Int J Mol Sci.* 2019 Apr 10;20(7):1781. doi: 10.3390/ijms20071781
- [5] Cannatelli A, Giani T, D'Andrea MM, Di Pilato V, Arena F, Conte V, et al. MgrB inactivation is a common mechanism of colistin resistance in KPC-producing *Klebsiella pneumoniae* of clinical origin. *Antimicrob Agents Chemother.* 2014 Oct;58(10):5696-703. doi: 10.1128/AAC.03110-14
- [6] Poirel L, Jayol A, Bontron S, Villegas MV, Ozdamar M, Türkoglu S, et al. The mgrB gene as a key target for acquired resistance to colistin in *Klebsiella pneumoniae*. *J Antimicrob Chemother.* 2015 Jan;70(1):75-80. doi: 10.1093/jac/dku323
- [7] Aires CA, Pereira PS, Asensi MD, Carvalho-Assef AP. mgrB Mutations Mediating Polymyxin B Resistance in *Klebsiella pneumoniae* Isolates from Rectal Surveillance Swabs in Brazil. *Antimicrob Agents Chemother.* 2016 Oct 21;60(11):6969-6972. doi: 10.1128/AAC.01456-16
- [8] Giordano C, Barnini S, Tsioutis C, Chlebowicz MA, Scoulica EV, Gikas A, et al. Expansion of KPC-producing *Klebsiella pneumoniae* with various mgrB mutations giving rise to colistin resistance: the role of ISL3 on plasmids. *Int J Antimicrob Agents.* 2018 Feb;51(2):260-265. doi: 10.1016/j.ijantimicag.2017.10.011
- [9] Jayol A, Nordmann P, Brink A, Poirel L. Heteroresistance to colistin in *Klebsiella pneumoniae* associated with alterations in the PhoPQ regulatory system. *Antimicrob Agents Chemother.* 2015 May;59(5):2780-4. doi: 10.1128/AAC.05055-14
- [10] Mhaya A, Bégu D, Tounsi S, Arpin C. MgrB Inactivation Is Responsible for Acquired Resistance to Colistin in *Enterobacter hormaechei* subsp. *steigerwaltii*. *Antimicrob Agents Chemother.* 2020 May 21;64(6):e00128-20. doi: 10.1128/AAC.00128-20
- [11] Martins W, Narciso AC, Cayô R, Santos SV, Fehlberg L, Ramos PL, et al. SPM-1-producing *Pseudomonas aeruginosa* ST277 clone recovered from microbiota of migratory birds. *Diagn Microbiol Infect Dis.* 2018;90(3):221–227. doi:10.1016/j.diagmicrobio.2017.11.003
- [12] European Committee on Antimicrobial Susceptibility Testing (EUCAST). Breakpoint tables for interpretation of MICs and zone diameters. Version 11, 2021. Available at: <http://www.eucast.org>. Published 2021. Access January, 2021.
- [13] Tenover FC, Arbeit RD, Goering RV, Mickelsen PA, Murray BE, Persing DH, et al. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol.* 1995 Sep;33(9):2233-9. doi: 10.1128/JCM.33.9.2233-2239.1995
- [14] Yang QE, Tansawai U, Andrey DO, Wang S, Wang Y, Sands K, et al. Environmental dissemination of mcr-1 positive *Enterobacteriaceae* by *Chrysomya* spp. (common blowfly): An increasing public health risk. *Environ Int.* 2019;122:281–290. doi:10.1016/j.envint.2018.11.021

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

[15] Barrios-Camacho H, Aguilar-Vera A, Beltran-Rojel M, Aguilar-Vera E, Duran-Bedolla J, Rodriguez-Medina N., et al. Molecular epidemiology of *Klebsiella variicola* obtained from different sources. *Sci Rep*. 2019 Jul 23;9(1):10610. doi: 10.1038/s41598-019-46998-9

[16] Fonseca EL, Ramos ND, Andrade BG, Morais LL, Marin MF, Vicente AC. A one-step multiplex PCR to identify *Klebsiella pneumoniae*, *Klebsiella variicola*, and *Klebsiella quasipneumoniae* in the clinical routine. *Diagn Microbiol Infect Dis*. 2017 Apr;87(4):315-317. doi: 10.1016/j.diagmicrobio.2017.01.005

[17] Lu Y, Feng Y, McNally A, Zong Z. Occurrence of colistin-resistant hypervirulent *Klebsiella variicola*. *J Antimicrob Chemother*. 2018;73(11):3001–3004. doi:10.1093/jac/dky301

[18] Minagawa S, Okura R, Tsuchitani H, Hirao K, Yamamoto K, Utsumi R. Isolation and molecular characterization of the locked-on mutant of Mg<sup>2+</sup> sensor PhoQ in *Escherichia coli*. *Biosci Biotechnol Biochem*. 2005 Jul;69(7):1281-7. doi: 10.1271/bbb.69.1281

[19] Olaitan AO, Diene SM, Kempf M, Berrazeg M, Bakour S, Gupta SK, et al. Worldwide emergence of colistin resistance in *Klebsiella pneumoniae* from healthy humans and patients in Lao PDR, Thailand, Israel, Nigeria and France owing to inactivation of the PhoP/PhoQ regulator mgrB: an epidemiological and molecular study. *Int J Antimicrob Agents*. 2014 Dec;44(6):500-7. doi: 10.1016/j.ijantimicag.2014.07.020

[20] Silva D, Faria-Junior C, Nery DR, Oliveira PM, Silva L, Alves EG, et al. Insertion sequences disrupting mgrB in carbapenem-resistant *Klebsiella pneumoniae* strains in Brazil. *J Glob Antimicrob Resist*. 2020 Nov 24;24:53-57. doi: 10.1016/j.jgar.2020.11.003

295

**Table 1.** Microbiologic and genetic features of *K. variicola* isolates evaluated in this study.

296

Microbiological and Genetic Features	Bacteria Isolates			
	M14	M15	M50	A58243
Bacteria species identification by MALDI-TOF	<i>K. variicola</i>	<i>K. variicola</i>	<i>K. variicola</i>	<i>K. variicola</i>
<b>Antimicrobial susceptibility testing<sup>a</sup></b>				
Colistin	128	64	64	0.25
Polymyxin B	128	128	64	0.25
Polymyxin B + EDTA (10 mM)	128	128	64	ND
Ceftazidime	0.25	0.25	0.25	≤0.25
Ceftriaxone	≤0.125	≤0.125	≤0.125	ND
Aztreonam	≤0.06	≤0.06	≤0.06	≤0.25
Levofloxacin	≤0.06	≤0.06	≤0.06	ND
Gentamicin	2	2	2	ND
Tobramicin	0.5	0.5	0.5	ND
Amikacin	2	2	2	2
Cefepime	≤0.125	≤0.125	≤0.125	≤0.25
Meropenem	≤0.06	≤0.06	≤0.06	≤0.06
Imipenem	≤0.125	≤0.125	≤0.125	0.25
Piperacillin-Tazobactam	2	2	4	8
Fosfomicin	16	16	4	8
<b>Genomic Features</b>				
PFGE	A1	A2	B1	ND

MLST	ST137	ST137	ST167	ND
Genome size (bp)	5,880,384	5,882,887	5,403,894	5,711,702
GC (%)	56.9	56.9	57.5	57.2
ORFs	5860	5855	5235	5566
RNAs	94	94	91	92
Contigs	105	104	77	83
Chromosomal beta-lactamase	<i>bla</i> <sub>LEN-24</sub>	<i>bla</i> <sub>LEN-24</sub>	<i>bla</i> <sub>LEN-13</sub>	<i>bla</i> <sub>LEN-9</sub>
<b>Alteration analysis in TCS components and regulators</b>				
PmrB	N13H E272K	N13H E272K	Not found	-
PhoQ	<b>D90N (deleterious)</b> D101N R116H	<b>D90N (deleterious)</b> D101N	Not found	-
RstB	<b>G385S (deleterious)</b> M82I	<b>I122S (deleterious)</b> M82I	Not found	-
MrgB	Not found	Not found	Deletion on 93 nct position	-

297

298

<sup>a</sup>The antimicrobial susceptibility profile of *K. variicola* isolates was determined by agar dilution, except for polymyxin B, which was tested by broth microdilution according to the

299

EUCAST/BrCAST guidelines. MICs were expressed in mg/L.

300 **Table 2.** Polymyxin B MICs of the laboratory derivative strains

Strains <sup>a</sup>	Mean Polymyxin B MIC (mg/L) <sup>b</sup>	Mean Colistin MIC (mg/L) <sup>b</sup>
M50	64	64
M50 + pskA	64	16
M50 pskA- <i>mgrB</i> -WT <sup>c</sup>	≤0.25	0.25
M50 pskA- <i>mgrB</i> -mutant <sup>c</sup>	64	64

301

302 <sup>a</sup>To prevent plasmid loss, MICs of completed strains were performed in presence of apramycin 50 mg/L.303 <sup>b</sup>Experiments were performed in biological triplicates.304 <sup>c</sup>*mgrB* from A58243 was used as wild-type (WT) while *mgrB* from M50 was used as *mgrB*-mutant.

305 **Figure 1.** (A) Relative transcriptional levels of TCS evaluated in polymyxin-resistant *K. variicola* isolates. (B) Multiple nucleotide alignment of *mgrB*  
 306 variants detected in this study. (C) Multiple protein alignment of MgrB proteins. Red lines highlight the change in nucleotide/aminoacid sequences.  
 307

