



Chapitre de livre

2000

Published version

Open Access

This is the published version of the publication, made available in accordance with the publisher's policy.

The flowering process : a prolonged incompletion

Greppin, Hubert; Simon, Patrice

How to cite

GREPPIN, Hubert, SIMON, Patrice. The flowering process : a prolonged incompletion. In: Integrated plant systems. Hubert Greppin, Claude Penel, William J. Broughton and Reto Strasser (Ed.). Genève : Université de Genève, 2000.

This publication URL: <https://archive-ouverte.unige.ch/unige:97994>

The flowering process: a prolonged incompleteness

H. GREPPIN & P. SIMON*

** Laboratory of Plant Physiology and Biochemistry, University of Geneva, Place de l'Université 3, CH-1211 Genève 4, Switzerland. E-mail: hubert.greppin@bota.unige.ch*

Introduction

Flowering, the sexual reproductive development, is a multistep integrated and dynamical networking process, oriented in time (physiological clocks) and space (root, leaves, stem, shoot apex territories), producing as finality a new organ: the flower. Since the cretaceous time ($\sim 1,3 \cdot 10^8$ years), it ensures the reproduction (species survival) and the space expansion of higher plants ($\sim 3,5 \cdot 10^5$ different species). This phase of the plant life cycle is the most important, together with the biomass production by photosynthesis (vegetative growth and development: $2 \cdot 10^{11}$ t dry weight per year).

The dynamic stability of ecosystems and biosphere, as well as environmental adaptation, are widely dependent on the evolution of flowering, in connection with the response capacity to ecospace changes (light, temperature, water, soil, etc.) and to biospace specific structure (mutation, recombination; population genetics; ecological web) [4,14,21,68,140].

The flowering process, putting aside the inconvenience of complexity, is a good model to study and understand the co-action and adaptation between the genetic biospace and the physico-generic activities on one side and the ecospace properties changes on the other side. These properties are, for example, temperature, light and water that are important agents for growth and development, and also signalling by photo-nyctotemperature alternation, photo-nyctoperiodic induction, water stress, etc. [58-60,112].

Despite the numerous publications that have appeared in this field during the last century, the flowering mechanism is still not clearly known. It is only recently, after a long period of morphological and ultrastructural description, followed by a chemical and biochemical analysis and a classical genetic approach, that we are progressively entering into a molecular genetic phase and a biophysical and physiological systemic approach. This new trend, with the help of a lot of new techniques (nucleic acids amplification and analysis, macro- and microarrays, bioinformatics, genomics, proteomics; atomic force microscopy, nuclear magnetic resonance, fluorescence analysis, network and far from equilibrium thermodynamics, metabolic compartmental

analysis, modeling, physiomatics, etc.) will probably allow to solve this always-open question during the new century (the *Arabidopsis* genome will be totally deciphered in 2001) [35,63,64,90,110,112,133,145,171].

The different components of the flower (sepal, petal, stamen, pistil) are the end-result of floral morphogenesis, determined by the action of some specific genes (floral meristem and organ identity genes) and the consecutive physico-biochemical changes in the apical shoot meristem: a few hundred cells constitute three territories, the initial ring for leaves production (plastochrone), the medullary and the prosporogenic territories for respectively stem and flowers formation. Significant knowledge has been acquired concerning the molecular nature of floral morphogenesis in the apex. Less is known about floral evocation, the phase following the vegetative functioning (leaves production). This period, lying from the onset of induction to the second wave of apical mitosis (onset of the floral pieces construction), is characterized by a dedifferentiation (hydrolytic activities) of the prosporogenic territory and then a stimulation of mitosis to produce, later, the flower. Evocation is coupled to a space-time change in the mitotic distribution and control of the different territories in the shoot apex (change in the phyllotaxis and Fibonacci series) [36]. Good indicators of evocation, as we have proposed, are the localization and amount of glucose-6-phosphate dehydrogenase activity, the mitochondrial and plastid DNA synthesis and the plasma membrane proliferation [5-7,13,39,57,87,151,157,194].

Another important progress has been made by the progressive identification of the genes responsible for the orientation in time (physiological clocks) and those encoding the pigments (phytochrome, cryptochrome, etc.) that enable to cope with environmental parameters triggering floral development (reproduction) or maintaining vegetative mass production [37,41,59,104,108,114,122,150,160].

The environmental conditions may act as optimizing agents which are monitored by the plant to determine the best time for reproduction or for biomass production. Each ecotype has a specific set of different genetic modules interfacing with the environment and allowing various adaptation patterns.

If the shoot apex has all the genetic information to produce flowers, the fitting of the reproductive program within some environmental and obliged internal parameters is an absolute necessity for the survival of the plant species, because they can not escape from the ecospace constraints. Plants are immobile, strictly depending on light and water for their autonomy and, contrarily to animals, have a low degree of global homeostasy against environmental pressure. They are however adapted by other specific ways, such as continuous embryogenesis, cellular totipotency, low degree of tissue and organ differentiation, leaf abscission, cross-resistance and stress resistance patterns, etc., affording a multifactorial solution leading to the same finality (equifinality). That "extraspection" feature offers multiple ways of responding to the fluctuating environment; some aspects of these modular responses are detected by the shoot apex to induce or not flower formation (integration with environmental and internal possibilities) [63,112,133].

Some signals (floral stimuli or inhibitors) are produced in the plant (leaves, root) by the transduction of eco- and internal sensors information (environment perception; memory, Pavlov-like reactions, learning) and, under the control of biological clocks, are then transferred (communication) to the apex where they trigger or block the

flowering process at the prosporogenic meristem level. The real nature of these stimuli and of the implicated genes (flowering genes) are for the moment not sufficiently known. But when the leaf has been induced to produce the floral stimuli, it is possible by grafting it to a vegetative plant, to induce the flowering of the receptor. This transition of vegetative leaf to floral state, characterized by specific proteins, is irreversible. The graft effect is linked to the plasma membrane symplastic reconnection with the stock. The phloem is an important conduit for the circulation of the floral stimuli. Senescence is accelerated when flowering is launched.

By the dynamic integration of the environmental possibilities it becomes possible to achieve with security (environmental risk) the reproductive development and to guarantee the future of the species. A varied networking of genes under activation or repression control (methylation, etc.) [22,34,38,45,72,146] associated with different functional modules and receptors in space and time (physiological, genetic and thermo-compensated clocks) is in operation (co-action) in the whole plant, integrating the influence of environmental stimuli to decide (at the apex) whether to enter the flowering stage or to remain at the vegetative state, on the wait. This particularity is important for ecological and geographical reasons and for optimization of the biosphere heterogenic and viable structure. Outcomes and applications from a better and perfect understanding of this process are manifold and relate to several fields: industry, agriculture, sylviculture, territory management, etc...[25,51,70, 170,171,175].

Genetics of flowering

During the last decade, molecular genetic studies have increased our understanding of flower development from the vegetative-to-floral transition to the specification of floral organ identity by homeotic genes. The genetic control of flowering time in *Arabidopsis* has been the subject of intensive research that has been reviewed in details [71,89,101,102,136,144,157] and we will describe here only some aspects and recent findings. The genetics of organ specification in the flower structure will not be considered (for reviews and updating data of the famous ABC model, see [123,129,136,188]). Genetic analysis has also been conducted with other plants such as pea and wheat [159,189] but has not been yet complemented by a molecular approach as in *Arabidopsis*.

Flowering is under the control of environmental signals such as light and temperature. In many plant species, the initiation of flowering is controlled by daylength, a phenomenon known as photoperiodism. The measurement of daylength by a daily timer, the circadian clock, is now generally accepted [171]. The control of flowering by temperature is operative through vernalization, a response of the plant to low temperature that abolish the repression of developmental processes, such as the initiation of flowering. This phenomenon is most apparent in plants growing at high latitudes, and the repression of flowering occurs even if the plant is grown under favorable photoperiods. The vernalization response can be considered in a way as a biological timer measuring the length of the winter period. Whereas the inductive daylength is perceived in the leaves and generates a signal that travels to the shoot apex, vernalization acts in the mitotically active cells that will form the inflorescence [45,72]. Flowering is also influenced by environmental nutrient deprivation and by endogenous developmental factors associated with the age of the plant.

Arabidopsis is a facultative long-day plant that flowers faster under long days than short days conditions. It will also flower in complete darkness in specific laboratory conditions such as growth on vertical plates which provide sucrose directly to the developing cotyledons [142]. In the mutant *embryonic flower (emf)* the vegetative phase will even be skipped and an inflorescence meristem will be produced upon germination [168]. The commonly used laboratory varieties Columbia, Landsberg *erecta* and Wassilewskija are rapid cycling ecotypes in contrast to other ecotypes that show a vernalization response.

Mutagenesis and molecular genetic studies has permitted the identification and cloning of many genes implicated in the control of transition to flowering. There are currently about 80 genes and loci known to affect flowering time in *Arabidopsis* [101]. In addition to the laboratory-induced mutants, the naturally occurring variation among accessions is becoming an alternative complementary source for mapping of loci and the cloning of large-effect genes (discussed in [2]).

The function of these genes, their distribution in specific genetic pathways, the interplay of these pathways in a complex genetic network has been reviewed recently [16,89,101,136,157]. Mutants affected in flowering time are either late or early, mostly late. A collection of late-flowering mutations has been identified by the pioneering work of Koornneef and coworkers [88-91]. These late flowering mutants are derived from the early flowering parent Landsberg *erecta* and have been distributed in two groups, according to their response to daylength and vernalization. One group is unaffected by vernalization and short days and is constituted by the mutants *fe*, *ft*, *fd*, *fwa*, *constans (co)* and *gigantea (gi)*. These mutants define genes required to promote flowering specifically in response to long day and belong therefore to a photoperiod pathway, photoperiod-dependent but temperature-insensitive. The other group is accelerated by vernalization and delayed in short days and is constituted by the mutants *fca*, *fve*, *fy*, *fpa* and *luminidependens (ld)*. This group of mutants are recessive and still able to respond to environmental cues, and they are therefore considered to be disrupted in an autonomous promotion pathway, photoperiod-independent but temperature-sensitive. A third pathway consists of a class of mutants reducing the biosynthesis or the response to the growth regulator gibberellin, with most severe effects in short days. A fourth pathway has been defined with mutants that are specifically impaired in their response to a cold treatment and that may be defective either in the perception of cold temperature or in the transduction of the cold signal. No single mutation has been yet described in *Arabidopsis* that abolish flowering completely, because the promotive pathways are functionally redundant. Due to this redundancy, inactivation of two pathways results however in a more severe phenotype than inactivation of one. The inductive pathways will lead to activation of the meristem identity genes that will change the fate of the meristem from vegetative to floral [135,186,187].

Control by temperature

In *Arabidopsis*, two major loci act together to delay flowering, *FRIGIDA (FRI)* and *FLOWERING LOCUS C (FLC)*. These loci have been characterized by comparing the naturally occurring varieties having or not a vernalization response. The vernalization-suppressible late-flowering phenotype of winter-annual ecotypes of *Arabidopsis*

results from the presence of dominant late-flowering alleles at these two loci. Both genes have recently been cloned. The *FLC* gene encodes a putative transcription factor of the MADS-box class acting as a dosage-dependent repressor of the floral transition [111,152]. The *FRI* gene is found as a single copy gene in the Arabidopsis genome and encodes an open reading frame for a protein of 609 amino acids, which shows no significant match to any known protein or domain or protein domain in the available databases [76]. The levels of *FLC* transcripts are regulated positively by *FRI* and negatively by vernalization and genome demethylation. A quantitative relationship has been found between the duration of the cold treatment and the extent of down-regulation of *FLC* transcriptional activity [154]. DNA demethylation is assumed to mediate the vernalization response [34,43,45,153]. It has been shown indeed that a reduction in genome methylation by the demethylating agent 5-azacytidine or by a methyltransferase [MET+] antisense substitutes for a vernalization treatment in promoting flowering [22,42,44]. A correlation was found between *FLC* transcription and decreased genomic methylation in the early-flowering *MET1* antisense plants [152]. A third gene, *VRN2* which has not been cloned yet, acts during vernalization and reduces *FLC* expression. From these observations, *FLC* is now considered to play a pivotal role in controlling the vernalization response. *FLC* is also regulated negatively by genes of the autonomous pathway (*FCA*, *FVE*, *FPA*, *FD*, *FLD* and *LD*), as indicated by the elevated levels of *FLC* and the vernalization-responsiveness of the late flowering mutants obtained from early ecotypes [111,154]. *LD* and *FCA* have been cloned and encode respectively a putative transcription factor [95] and an RNA binding protein [106], presuming a control of *FLC* at both the transcriptional and posttranscriptional levels. Taken together, *FLC* appears as integrating the vernalization and the autonomous pathways. It has also been proposed that *FLC* may act by antagonizing the promotive effect of gibberellins on flowering, since the *flf-1* mutant, overexpressing the *FLC* transcripts, requires higher amount of exogenous GA to decrease flowering time [152]. Another MADS-box gene, *SHORT VEGETATIVE PHASE* [*SVP*], has been shown recently to repress flowering and the possibility of forming a *FLC/SVP* heterodimer has been suggested [69].

It is worth mentioning that contrasting phenotypes have been related to altering methylation, such as the late flowering phenotype of *fwa* mutants, which is caused by gain-of-function epigenetic alleles of a homeodomain gene induced by hypomethylation of two direct repeats in the 5' region of the gene [162]. The function of *FWA* in flowering of wild-type plants is questioned by the authors.

Control by daylength

The molecular relationships between the regulation of flowering time by daylength and the circadian clock has gained evidence from the recent description of mutants in which both circadian rhythms and flowering time are disrupted. The clock is entrained by light input signals that are perceived by the photoreceptors phytochromes (red light) and cryptochromes (blue light), which also regulate flowering [67,103,114,166,171]. The molecular basis of circadian oscillators has been established as oscillating transcription/translation loops of positive and negative elements more or less conserved between *Drosophila*, *Neurospora* and mammals [37]. In plants, candidates for a molecular circadian oscillator have been identified recently, but a specific role for

these candidates remains to be asserted. The plant oscillator would apparently be different to the ones established in other organisms, since components are not homologues. One of the best candidate as an oscillator component is *TIMING OF CAB (TOC1)*, which encodes a nuclear protein containing an atypical response regulator receiver domain, and two motifs that suggest a role in transcriptional regulation [165]. The next two candidates are the *CIRCADIAN CLOCK ASSOCIATED 1 (CCA1)* and *LATE ELONGATED HYPOCOTYL (LHY)* genes encoding both MYB-related transcription factors [147,185]. Modulation of CCA1 by protein kinase CK2 has been shown and suggested to play a role in vivo [167]. Two other candidates are the *ZEITLUPE (ZTL)* and *FKF1* genes and its related homologue *LKPI* [85,113,121,161]. These genes encode novel proteins with a PAS domain similar to the flavin-binding domain Arabidopsis blue-light photoreceptor NPH1, an F-box characteristics of proteins that direct ubiquitin-mediated protein degradation, and 6 kelch repeats that could serve as the protein-protein interaction domain recruiting specific protein for degradation. Disruption of any of these putative clock gene results in an altered flowering time. Mutations in *FKF1* is also responsive to vernalization. Its role might therefore be more complex than just acting in the pathway that promotes flowering in response to long days. The well known photoperiod mutant *gi* was characterized recently at the molecular level [46,73,128]. From computer-based prediction, GI was first proposed to be an integral plasmamembrane-localized protein, but recent data have shown that it is a nucleoplasmically localized protein [73]. GI was shown to be required for maintaining circadian amplitude and appropriate period length in the expression of its own gene as well as *CCA1* and *LHY*, and also to affect light signalling to the clock, which suggests a role in a feed back, loop in the circadian clock. The photoperiod information is further relayed to *CONSTANS (CO)*, a gene that encodes a zinc finger transcription factor and that activates downstream floral meristem identity genes such as *LEAFY (LFY)*, *TERMINAL FLOWER 1 (TFL1)* and *APETALATA 1 (API)* [15,135,138,141,156]. The *FLOWERING LOCUS T (FT)* is also acting downstream of *CO*, but its promotive action is antagonized by the homologous gene *TFL* [79,86]. The *LFY* promoter integrates the floral inductive signals of both the GA and the photoperiod pathways at different cis elements [17] and the protein LFY participate in cell signalling by moving from cell to cell [151].

Using a suppressor mutagenesis approach [127,145], CO has been shown recently to play a pivotal role in promoting flowering by activating at least four early target genes: *SUPPRESSOR OF CONSTANS 1 (SOC1)*, previously known as *AGAMOUS-LIKE 20* [AGL20], *FLOWERING TIME (FT)*, *ACS10* and *AtP5C52*, encoding respectively a MADS-box transcription factor a putative phosphatidylethanolamine binding protein, an enzyme involved in ethylene synthesis and an enzyme involved in proline biosynthesis. Floral meristem identity genes within the shoot apex are subsequently activated by *SOC1* and *FT*. A reduction of the expression of *SOC1* and *FT* has been observed in a low *FLC* mRNA background and it has been proposed that levels of *SOC1* and *FT* expression may be determined by a balance of CO and FLC, and represents therefore common components of distinct flowering-time pathways. The role of SOC1/AGL20 as an integrator of the autonomous, vernalization and photoperiod pathways was also shown by another group [94].

Despite all the knowledge acquired from studies on the genetic control of flowering, there is no clear evidence about the identity of genes implicated in the signal generation in the leaves, its transmission and its perception in the shoot apical meristem (discussed in [25,105]). *CO* for example is expressed uniformly in the leaves and in the shoot apical meristem. In maize, *ID1* which encodes a zinc finger protein (like *CO*) acts non-cell-autonomously and has been proposed to regulate the production of a floral stimulus or the repression of a floral inhibitor [25,26]. It has been hypothesized [25] that a mechanism for the long distance transmission of the flowering signal could be posttranscriptional gene silencing [40,77,117]. Indeed, the epigenetic state of genes can be transferred from one part of a plant to the other via plasmodesmata and phloem channels and could implicate double-stranded RNA [134,169,178,192]. The epigenetic silencing at the transcriptional level can also be considered as a way of controlling developmental gene expression. It is for example regulated by the polycomb proteins and several of them have been identified in Arabidopsis, namely *CURLY LEAF* which is required to regulate expression of floral homeotic genes [52,84,126,137]. Chromatin remodelling through methylation-dependent or -independent mechanisms has been observed lately in Arabidopsis and represent a basis for gene activation and repression during development [3,38,74,75,78,163,164,191].

Flowering induction and ecospace

In the case of photoperiodically controlled plants (LDP, SDP, etc...), it is possible to demonstrate that floral induction proceeds in a networking three-step process, as soon as the critical photo- or nyctoperiod is reached (physiological and molecular clocks control). These three steps are the following (Fig. 1):

- 1) Following a juvenile phase of the leaf (emergence of sensitivity), light on and light off detection of photo-nyctoperiod duration by several specific pigments like phytochromes, cryptochromes and other photoreceptors [41,103,114,166]. Depending on external and internal coincidences of this event with the metabolic effects of clock genes [89,121,128,144] and of biophysical oscillators, photo-nyctoperiodic induction and structuro-functional transduction (flowering genes) of this processing to all the cells of the leaf. Then, production of the floral stimuli of unknown nature and, with some delay, acquisition by the leaves of the floral state, characterized by a few new specific proteins [9,17,96,97,109,141,157]. The induction property is transmissible to vegetative plants by grafting of an induced leaf.

- 2) Migration-transmission of the floral stimuli through the petiole and stem towards the apical meristems which have become competent. The extraction and characterization of the floral stimuli are the subject of a lot of uncertainty despite the quality of the research [14,24,66,77,105,124,134,184,193].

- 3) Commitment of the meristem cells and territories to floral evocation, and topological modulation of the mitotic activity; then, after two general waves of mitosis, start of floral morphogenesis as a consequence of the activation of floral meristem and organ identity genes. This last step is well known for the moment [5-7,13,14,156].

Ecospace-Biospace (Higher Plants)

The Flowering Process (Reproduction)
Unsolved problem since > 100 years of studies
(Autonomous , photoperiodic, vernalization pathways)




Figure 1. The Plant System and the Flowering Process. P.M.: prosporogenic meristem territory; M.M.: medullary meristem territory; I.R.: initial ring, meristem territory.

Table 1. Some physical, chemical and biochemical parameters correlated with the induction process in leaves of spinach (LDP). Trends to a maximum or minimum value at the onset of induction [10-12,19,20,29,31,32,99,107,133,148,149,176].

Parameter	Critical Photoperiod 11-12 hrs light	Floral induction Effect
1. <u>Energy-signals</u> a) Energetic charge b) Redox charge c) Chloroplastic MDH.NADPH dep. d) Cytosolic MDH. NADH dep. e) Chloroplastic A.K., C1 f) Cytosolic Adenylate Kinase g) New pool of free glucose	max. max. max. max. min. max. → max.	+ 50 % + 100 % + 200 % + 100 % - 70 % + 50 % + 400 %
2. <u>Proton pumps, AT~P dep.</u> a) App. K_m b) Activity c) I A A desensitization	→ min. → max. transition	- 68 % $10^{-11}M \rightarrow 10^{-6}M$ -30% + 30 %
3. <u>Calcium Pumps. AT~P dep.</u> a) App. K_m b) Calcium desensitization	→ max	+ 50 %
4. <u>Water organization</u> a) 1H NMR; T1	min	- 50 %
5. <u>Plasma membrane</u> a) Oleic/Linoleic acids in phospholipids ($C_{18:1} / C_{18:2}$) b) ^{31}P NMR c) Thickness d) C_6 ^{19}F NMR (lyophilized); T1	max. Shoulder max. min	+ 60 % Hexagonal II ? + 20 % - 40 %

Many hypotheses have been put forward during the last 100 years to explain the flowering process and the real nature of the floral stimuli, from J. Sachs to G. Klebs proposals (specific floral substances versus ratio between substances); for example, specific hormones and inhibitors [24,93,195], sink-source chemical interaction [142,143,158], fragments of cell-wall specific oligosaccharides [124], volatile substances [193], plasmamembrane modification [53,56], electrochemical and hydraulic signalling and coded frequency [184], etc...

Different ways and modules of interactions, feed-backs and communications could be activated during the induction of flowering in leaves and roots, implicating immediate and short-term regulation as the slower one. Quantitative increase or decrease and qualitative effects could be observed, depending on the evolution of environmental constraints (light, temperature, water, stress and shock) and on the species and genetic dotation of the strain: different ways could be activated or inhibited to tend, by equifinality, to the same final result, the production of the floral stimuli. We will present here some aspects we have prospected with spinach, a qualitative long-day plant, *Arabidopsis thaliana*, a quantitative long-day plant and *Chenopodium rubrum*, a short-day plant.

Energy-signal and brownian motion

Energy and its control are playing a central and important role during development. Adaptation to new environmental conditions (day, night duration) might be immediately reflected in the alteration of energy transduction. Therefore emphasis should be put on the time-dependent interconversion in adenine and pyridine nucleotide pools. The Energy Status [18-20,65,112,179] has been determined by different approaches and evaluations: chemical analysis of free glucose or saccharose, biochemical analysis of nucleotides and calculation of the energy and redox charges (the energy charge $(\text{ATP} + \frac{1}{2}\text{ADP})/(\text{ATP} + \text{ADP} + \text{AMP})$; the redox anabolic charge $\text{NADPH}/(\text{NADP} + \text{NADPH})$; the redox catabolic charge $\text{NADH}/(\text{NAD} + \text{NADH})$), spectroscopic analysis, as for example ^1H -NMR or C_6^{19}F -NMR (nuclear magnetic resonance), a non-invasive method [54,99]. By this technique, it is possible to get a precise identification of chemicals and to appreciate intramolecular exchange, intermolecular dynamics and statistical entropy variation. The photoperiodic induction time in spinach leaves (Table 1 and Figs 2,3,4) corresponds to the circadian period of the maximum levels in energy and redox charges and of the minimum levels of statistical entropy (frequency of correlation f_c). At this time, an important increase of free glucose is observed, which can also be detected when floral induction is provoked by a displaced-short-day treatment. This rise corresponds to a photoperiodic and clock control associated to the processing of floral induction, as it is usual for LDP. The comparison with SDP (Fig. 5) suggests that the photoperiodic free glucose control relates to the cell filling in LDP and the cell emptying in SDP (differential threshold). Another aspect of the sensitivity of this photoperiodic free glucose pool is illustrated by the inhibition of induction by a stress or a pricking (Fig. 6), that can abolish this effect for about 48 hrs [31-33,47,133,158].

The plant cells contain 80 to 90 % of water and a great number of molecules are essentially constituted by hydrogen. Biodynamics attempts to investigate the physical status of living cells by the measurement of random movements, as for example the




Figure 2. Time-dependent evolution in short days (vegetative state) of the relative correlation frequencies f_c ($^1\text{H-NMR}$: 90 MHz) and free glucose concentration in spinach leaves (8 hrs light, 16 hrs night, 4 weeks-old); f_c rhythm: 22,8 hrs [31,32,99].




Figure 3. Idem fig. 2, but after an inductive transfer to continuous light (flowering induction). Arrow: onset of the critical photoperiod (C. Ph) for the induction of flowering; f_c rhythm: 24,4 hrs




Figure 4. Time-dependent evolution under permanent continuous light (3 weeks) of f_c in fully induced plants (floral state).




Figure 5. Free glucose content evolution in petioles of spinach leaves (LDP) and *Chenopodium rubrum* (SDP) during flowering induction, respectively by transfer to continuous light or to night. Arrows: critical photo- or nyctoperiod [133].




Figure 6. Free glucose content in spinach leaves and petioles at the vegetative state (short days) and after a light inductive transfer (23 hrs). Comparison with pricked plants (stress) [33,133].

brownian motion [63,98]. This particular motion is in relation with the general thermic motion of water (heat reservoir) through the dissipation of energy by the metabolism. NMR spectra and other characteristics such as spin lattice relaxation time (T_1) give us some information about the status of the Brownian system, from random organization to different degree of order and liberty. Macromolecules and membranes are playing an important role in controlling the degree of order of water molecules and the exchange between molecules, at long distance too. Ordered water could in turn organize other molecules and transmit some signals. With a correlation time ranging from 10^{-12} sec to 10^{-6} sec, the motion gives essentially a thermic contribution to internal energy, entropy and pressure ($E = \frac{3}{2} kT$); but from 10^{-5} sec to 10^{-1} sec, it can

provide signals (slow co-operative movement of polymer chains in membranes and chromosomes as momentaneous water molecules alignment).

Brownian dynamics is quantitatively determined by the correlation time τ_c , the activation energy E_a and the diffusion coefficient D . The dynamic entropy S_d is related with the correlation time and the frequency of correlation: $S_d \propto \frac{1}{\tau_c} \propto f_c$. The correlation time may be determined by NMR spectra (bandwidth Δ) and the measurement of the spin lattice relaxation time T_1 : $\Delta \propto \tau_c \propto \frac{1}{\tau_c} \propto \frac{1}{f_c} \propto \frac{1}{S_d}$. The ordered molecular systems are present in membranes with lipid bilayers and in some part of water, in the vicinity of membranes, chromosomes, macromolecules and ions.




Figure 7. ^1H -NMR spectra of *Chenopodium rubrum* plantlets (SDP) in continuous darkness (induction of flowering). C.N.: critical nyctoperiod. Measurement, in a flux calorimeter, of heat exchanges during the same time (continuous curve). Endo-exothermic rhythm of 22 hrs [54].

Proton NMR in spinach leaves and *Chenopodium* plantlets (Figs 2,3,4,7; Table 2) gives us an information about the biodynamics of water molecules under the control and modulation by macromolecules, membranes and cell energetization. In both cases, it is when the photo- or nyctoperiod are critical that we observe a maximum of restriction of the water motion; it means a change in brownian versus semi-brownian dynamic order. The frequency of correlation f_c , and also T_1 and S_d present a minimal value. A maximum is measured for the bandwidth line Δ , and the correlation time τ_c (free water : τ_c , 10^{-12} sec; T_1 , 2,6 sec; proton exchange time, $2 \cdot 10^{-4}$ sec; the situation in leaves is different because the presence of various populations of water molecules : τ_c from 10^{-12} sec to 10^{-6} sec, as 10^{-5} sec, 10^{-4} sec, etc.; T_1 from 150 msec to 800 msec).

The dynamically ordered water, because of its specific physical and chemical exchange capacity, could transduce signals reflecting the energetic state in the cell, and through the physico-generic interaction with plasmalemma could operate at distance from leaves to apex (chromosomes could be another target of that water signal). So we hypothesize that the water brownian motion control and modulation could play a role as *primum movens* in the flowering induction process in leaves by its interaction with plasmamembrane ($T_1 \text{ Fl} < T_1 \text{ Veg}$; $\tau c \text{ Fl} > \tau c \text{ Veg}$; $fc \text{ Fl} < fc \text{ Veg}$; $\Delta \text{ Fl} > \Delta \text{ Veg}$; $Sd \text{ Fl} < Sd \text{ Veg}$ ratio of negative entropy variation ~ 2)[11,12,19,20,28,29,32,64,99,107,148,149,167,176].

Table 2. Plasmamembrane thickness evolution in apex or leaves of spinach (LDP) or *Chenopodium rubrum* (SDP) correlated with the induction process. [27,59,107]

Spinach : LDP	Plasma membrane Thickness
Short Days (8h. L, 16h. N): 4 weeks Vegetative plants	Leaves: $8,6 \pm 0,3 \text{ nm}$
	Apex: $8,9 \pm 0,5 \text{ nm}$
Transfer to continuous light (24hrs) Floral induction	Leaves: $10,2 \pm 0,2 \text{ nm}$
	Apex: $10,4 \pm 0,4 \text{ nm}$
Short days + GA_3 (10^{-3} M , 3 days) Floral induction	Leaves: $10,2 \pm 0,3 \text{ nm}$
	Apex: $10,6 \pm 0,4 \text{ nm}$
Chenopodium rubrum: SDP	Plasma membrane thickness
Continuous light (3 weeks) Vegetative plants	Leaves: $8,2 \pm 0,3 \text{ nm}$
Transfer to 12 hrs night Floral induction	Leaves: $9,6 \pm 0,2 \text{ nm}$
Transfer + red light break of night. Vegetative plants	Leaves: $8,3 \pm 0,5 \text{ nm}$

Plasmamembrane

During the critical period of leaf induction by light in spinach or Arabidopsis, by night for *Chenopodium*, some structural, biochemical and physiological modifications are very rapidly occurring in plasmalemma both in leaves and apex. For example, the thickness of the membrane is increasing ($\sim 2\text{nm}$) and the peripheral aspect is more clotted or lumpy after photoperiodic induction (osmic fixation). With NMR spectroscopy, we observe, at this time, a decrease of diffusivity of small molecules such as fluor nuclei (probe) or proton with about the same ratio of bandwidth line or T_1 variation. This restriction of molecular motion corresponds to a higher crosslinking with the macromolecular cellular system after induction, and with a blocking of the molecular dynamics, an increase in structuration and order, a decrease of the entropy of fluctuation. The ^{31}P -NMR suggest the appearance of hexagonal II phase in the plasmalemma (Fig. 8)[8,27-30,48,55,56,99,100,133].




Figure 8. ^{31}P -NMR (200 MHz) of spinach leaves (plasmamembranes) at the vegetative state (T) and after transfer in continuous light (I: induction of flowering. Arrow: shoulder (hexagonal II ?) [28, 133].




Figure 9. IAA sensitivity evolution of the H^+ pumps activity, ATPase dependent, at the vegetative state (spinach plasmamembranes) and during the transition to floral state (transfer from SD to continuous light: CL: 13 hrs, critical photoperiod). Three weeks-old spinach [11,12].

Usually the lamellar alpha phase is typical of membranes of young functioning cells, and the beta phase gel concerns the membranes of senescing cells. Under certain circumstances (temperature transition, increase of intercellular ions, hydration degree of tissue) a non-bilayer configuration could occur in the membrane mosaic. Phospholipids molecules could form cylinders with the polar head groups facing a water-filled core. Such inverted micelles could be distributed like hydrophobic buttons strewn the lumpy floral membrane and could serve as endogenous calcium ionophore or Ca channels. A rise of calcium is appearing during the photoperiodic induction and calcium pumps sensitivity is diminishing during this phase. All these modifications could influence the conformation of enzymes and their interactions (calcium and proton pumps, auxin receptors, etc.). The apparent K_m of proton pumps ATPase dependent is decreasing and a desensitization to IAA occurs during floral induction (Table 1, Fig. 9). In spinach, the induction is associated with a high level in the endogenous auxin concentration and with a low level of peroxidase activity.




Figure 10. Cybernetical, structural and thermodynamical model. S: statistical entropy. P.M.: plasmamembrane. E.C.: energetic charge. R.C.: redox charge. Vegetative homeostasy and rheostasy (plastochrone): global master negative feedback (specific energetic and structural envelope). Transition: appearance of some positive feedbacks. Floral state: new homeo- and rheostasy in a new envelope, stabilised with new proteins pattern.




Figure 11. Effect of 2 min R (660 nm; 0,045 W/m²) or 2 min FR (730 nm; 0,015 W/m²) on the activity of free basic peroxidase extracted from leaves of SD-grown spinach (vegetative state). Data are expressed as % variation of the irradiated plants versus non-irradiated ones, extracted at the same time [80-83,131-133].




Figure 12. Idem fig. 11, but after induction of flowering with transfer from SD to continuous light (24 hrs). The response is inverted.

During the floral transition the chemical analysis points out a change in plasmamembrane sterols and an increase in the ratio of fatty acids C 18:1 (saturated) to fatty acids C 18:2 (unsaturated). This could influence the structural conformation and the exchange capacity of membranes. For example, the tails of lipids exhibit rotational motion (τ_c 10^{-9} sec to 10^{-7} sec), translational movement (τ_c 10^{-7} sec to 10^{-6} sec), exchanges among polar heads (τ_c 10^{-5} sec), slow flips along whole lipids (τ_c 10^{-1} sec), with some consequences in the bilayer lipids transmission along the membrane and the interactions with various membrane ligands (receptors, ions, secondary messengers, etc.). This physico-generic aspect and the interaction with the water brownian control and modulation could be the *primum movens* of the induction in leaves that lead to floral information, after the perception of specific light duration and temperature in conjunction with a clock control. Externalities of these primary effects are ionic and electrical messengers and hormonal-receptors modifications in co-action with quantitative gene activation, in a first time, followed by qualitative activation to stabilize the leaf floral state (new specific proteins). The Figure 10 summarizes the regulatory relation between energy, plasmamembrane state and the induction of flowering. We attach importance to a systemic physico-chemical effect during the first step of leaf induction [46,49,58,63,64,101,112,130,155,177,180-183].

Peroxidases and phytochrome

Peroxidases have been implicated with various physiological process and in free radical scavenging and auxin catabolism: control of endogenous free auxin level, cell wall biogenesis, H_2O_2 consumers or producers. The number of genes and isozymes in a plant is very high (60 to 90 genes for *Arabidopsis thaliana*).

Total peroxidase activity in leaves exhibits a clear correlation with the developmental stage of spinach : a minimum value is observed at the time of induction and floral evocation, after this period a progressive increase occurs until the floral morphogenesis and senescence. The balance between anodic and cathodic peroxidases is very rapidly modified during the leaf induction, with a great decrease in the activity of basic peroxidases.

Basic peroxidasic activity is under the control of phytochrome as shown by the 660 versus 730 nm light effect on the activity. The peroxidase activity reacts rapidly to 1 or 2 minutes of *in vivo* irradiation of the leaves (Fig. 11) and this photocontrol is modified at the moment of floral induction in leaves (Fig. 12). The effect of this local photoconversion, was very rapidly propagated to all the parts of the plant (Figs 13,14). This signal transmission was inhibited or delayed by several substances including LiCl, EGTA and the Ca ionophore A23187. Figure 15 gives a temptative model explaining the propagation of the signal and the modification of the kinetics after leaf induction that could be linked to the plasmamembrane modification [23,50,80,81,130-132].

We have demonstrated that it was possible to modulate, both *in vivo* and *in vitro*, a specific binding or release of the basic peroxidases (glycoproteins) on leaf plasmamembrane or tonoplast by far-red and red light (phytochrome), or by Mn^{2+} , Ca^{2+} , and EGTA. This control is modified during the induction process in the LDP spinach as well in the SDP Pharbitis. This association of some peroxidases with membranes is specific to sites in membranes. The plasmamembrane of Pharbitis leaves

present a hemicircadian rhythm of the basic peroxidase association capacity, with a maximum at the moment when the sensitivity to red light in flowering inhibition is at its maximum (Fig. 16). In spinach a similar observation could be associated with red or far-red light effect (Fig. 17). A correlation is observed with the light on, light off control of the plasmamembrane electrical potential [82,83]




Figure 13. Idem Fig. 11; test with the irradiated leaf (continuous line) as with non-irradiated leaf (broken line) of the same plant (communication at distance). Vegetative state (spinach).




Figure 14. Idem Fig. 12 and same test as Fig. 13, after flowering induction.




Figure 15. Model of the transmission in the whole plant of the "peroxidase signal": interaction between light, phytochromes and other pigments, ions, plasmamembrane (timing mechanism, receptors, structural organisation, ion pumps, secondary signals, etc.) and peroxidases. The "flowering signals" are transmitted through plasmamembrane and phloem cells (symplast). The transmission of both signals implicates Ca^{++} and could be blocked by LiCl , whereas ouabain blocks only the peroxidase signals.




Figure 16. Basic peroxidase binding on purified plasmamembrane (*Pharbitis nil*, SDP) extracted at different times of the day (hemicircadian rhythm). Vegetative state. Red light break (660 nm) during inductive long night has the maximum effect (inhibition of flowering) at 20 hrs. It corresponds to the period when the maximum peroxidase binding could be observed [83,133].




Figure 17. Correlation between the light on effect (Red or Infrared: 660, 730 nm) on the binding or release of basic peroxidases with the membrane, and the light on (left) or light off (right) effect on the bioelectrical potential (micro-electrodes) of leaf plasmamembranes (spinach). Electrochemical and ionic mediation [133].

Bioelectricity

Active cells and organs are surrounded by electrochemical activities essentially stemming from photosynthetic metabolism during daylight and respiratory pathway during the night. About half of the generated energy is spent in electrochemical activities. As a result, because of biological activity far from equilibrium and thermodynamically open....,dynamic electrical potentials (span: ± 150 mV) are generated at different compartmental levels (plasmamembrane, chloroplasts, mitochondria, nucleus, vacuole). The value and evolution of these potentials give us an insight about the intra- and inter- compartmental relation in cells and between the tissues and organs, the interaction with environmental constraints and the stage of growth and development of the plant. The coupling of bioelectrical tests with the use of different LDP or SDP, and mutants or transgenic plants, could be a way of identifying, in different environmental background, what are the primary invariants of leaf induction producing the flowering stimuli. A pattern of gene activity upstream could be determined by microarray.




Figure 18. Fluctuation of the bioelectrical potential of plasmamembrane (spinach leaves) during one day, at the vegetative state (4 weeks-old plants. SD: 8 hrs light, 16 hrs night) [125].

In continuous light (floral state) spinach leaves show a rest-potential at the plasmamembrane modulated by a circadian rhythm of $25,2 \pm 0,9$ hrs, supporting an alternate train of small electrical fluctuations (5 to 10 mV) with a frequency of about $1,4 \cdot 10^{-3}$ Hz, for a first set and $8,9 \cdot 10^{-4}$ Hz for another set (hemicircadian alternation). Under short days (vegetative state), a fluctuation of $8,4 \cdot 10^{-4}$ Hz is observed during the light period and $2,2 \cdot 10^{-3}$ Hz during the night (Fig. 18). If we induce flowering by transferring the plant from short days to 30 hrs continuous light, and then back-transferring the plant in short days, the frequency of electrical oscillation is $2,2 \cdot 10^{-3}$ Hz at the night, as before, and respectively $4,9 \cdot 10^{-4}$ Hz and $6,7 \cdot 10^{-4}$ Hz during the light period of the first and second short-day. This result suggests that flowering induction in leaf has modified the electrical response and regulation in light, which indicated that some irreversible transformation has occurred [125].








Figure 19. Detail of surface bioelectrical potential (leaf of bean) evolution: light on, light off effect (polarization, depolarization) during ~20 minutes (12 hrs light, 12 hrs night) [62].

In photoperiodic plants the orientation in time is dependent upon the coincidence of light on and light off signals with the internal clock system oscillations. The light-dark breaks produce specific fast variation of the plasmamembrane electrical potential and of the surface potential (Fig. 19). These signals present circadian property in the electrical signature with two series of fluctuations, 0,016 Hz and 0,0041 Hz, which are closely fitted with the photonyctoperiod length (enslaving). Some modifications of the signal structure seem to be correlated with flowering induction (Fig. 20) [66].

These results and the existence of a clock adjustment between the different parts of the plant (multiplicity of rhythmic phenomena) suggest that the periodic organization in electrochemical oscillation could be the basis for communication between leaves and apex via frequency coded signals. Apical cells would be activated to modulate mitosis in the different apical territories, as a consequence of brownian water

modulation, and because of changes in plasmamembrane that provoke coded electrochemical fluctuations and transduction to secondary messengers, hormones and receptors (Figs 21,22) [1,61,62,92,115,116,174,184].

Bioelectrical Potential

Parameter	Amplitude A		Period τ	
	Fl.	Veg.	Fl.	Veg.
ON				
OFF	 30%		 50%	

CHENOPODIUM: S.D.P

Figure 20. Light on, light off global analysis at the vegetative state and after flowering induction in *Chenopodium rubrum* leaves (SDP): modification of the amplitude of the signal as the duration of this one [66].

Physiomatics

Macrofunctions, plasmamembranes and clocks are playing an important role in this vision of the primary events of flowering induction in leaves (Figs 21 to 23). Concerning the time measurement, NADH oxidase associated to membranes, could be implicated as an ultradian and thermocompensated rhythm of redox activity [57,58,118-120,139,172,173,177,190].

During these last years, important progress has been made in computing registration and sensors development (macro, micro, nano-sensors), and in mathematical and system analysis. It is possible to fit out a plant with a lot of various probes for continuous measuring and registration of the different physical, chemical and biochemical parameters. Depending on the program, it is possible to stimulate or inhibit the plant with the action of electrodes and other devices. So we can prospect the capacity and pattern of cybernetic accommodation and adaptation of the plant under different and global environmental treatment (light, temperature, chemicals) in relation with genetic properties (mutants, transgenics, ecotypes). The set up of a phytofeedbacks between the measured parameters and the imposed program of stimulation or inhibition, under different global environmental constraints, allows by this self-enslaving the determination of the different self-equilibrium level, by various genes activation or inhibition, and also the nature of the various transition




Figure 21. General sketch of the basic events during the flowering induction in leaves. Veg: vegetative; Fl: floral. dS : entropy variation. P.M.: prosporogenic meristem. M.M.: medullary meristem. I.R.: initial ring. E.C.: energetic charge. R.C.: redox charge. R.M.: root meristem. L.D.: long day. S.D.: short day.




Figure 22. Coded electrochemical model of communication (hypothesis). N: nucleus. Chl.: chloroplast, photosynthesis. Mito: mitochondria, respiration. Vac: vacuole. Cytoplasm [66,183,184].




Figure 23. Cellular working programs supervised by a few macrofunctions controlling at distance a few number of tissues and organs [59].

possibilities and the damageable situations. An expected result, in particular when the experimentation is made with the extreme environmental constraints that lead to flowering induction, is the progressive determination of the operators which associate the genetic matrix of information with the dynamical phenotypic expression (non-linear equation) (Fig. 24). It should represent a progress in the discovery of the information grammar of the interacting genes in plant, and also in the dialectic of biospace versus ecospace [51,58, 60,70,134,155,160,170,175,181].

References

1. Adamec L, Machackova I, Novakova-Prospikova M, Krekule J (1989) Electric current inhibits flowering in the short-day plant *Chenopodium rubrum* L. J Plant Physiol 134: 43-46
2. Alonso-Blanco C, Koornneef M (2000) Naturally occurring variation in Arabidopsis: an underexploited resource for plant genetics. Trends Plant Sci 5: 22-29
3. Amedeo P, Habu Y, Afsar K, Scheid OM, Paszkowski J (2000) Disruption of the plant gene MOM releases transcriptional silencing of methylated genes. Nature 405: 203-206

PHYSIOMATICS

10^{11} - 10^{13} cells




Figure 24. Physiomatics, a way to prospect the genes networking in relation with global effects on the plant territories [59].

4. **Atherton JG** (1987) Manipulation of Flowering. Butterworths, London
5. **Auderset G, Gahan P, Dawson A, Greppin H** (1980) Glucose-6-phosphate dehydrogenase as an early marker of floral induction in shoot apex of *Spinacia oleracea*, Nobel. Plant Sci Lett 20: 109-113
6. **Auderset G, Greppin H** (1976) Etude de l'apex caulinaire d'épinard avant et après l'induction florale. Saussurea, Genève 7: 65-72
7. **Auderset G, Greppin H** (1977) Effet de l'induction florale sur l'évolution ultrastructurale de l'apex caulinaire de *Spinacia oleracea*, Nobel. Protoplasma 91: 281-301
8. **Auderset G, Sandelius AS, Penel C, Brightman A, Greppin H, Morré DJ** (1986) Isolation of plasma membrane and tonoplast fractions from spinach leaves by preparative free-flow electrophoresis and effect of photoinduction. Physiol Plant 68: 1-12
9. **Balet-Buron A, Greppin H** (1977) Etude immunochimique de l'induction florale chez *Spinacia oleracea*. Saussurea, Genève 8: 57-64
10. **Bellamine J, Greppin H, Arrizabalaga PH** (1993) Etude par HATR/FTIR de la membrane plasmique des feuilles d'épinard avant et après induction florale. Analysis 21: 31-33
11. **Bellamine J, Penel C, Greppin H** (1993) Proton pump and IAA sensitivity changes in spinach leaves during the flowering induction. Plant Physiol Biochem 31: 197-203
12. **Bellamine J, Penel C, Greppin H** (1994) Effect of photoperiod and gibberellic acid on the plasma membrane H⁺ATPase activity of spinach leaves. Biol Plant 36: 565-573
13. **Bernier G** (1988) The control of floral evocation and morphogenesis. Annu Rev Plant Physiol Plant Mol Biol 39: 175-219
14. **Bernier G, Kinet JM, Sachs RM** (1981) The Physiology of Flowering. CRC Press, Boca Raton, Florida
15. **Blazquez MA** (1997) Illuminating flowers: CONSTANS induces LEAFY expression. Bioessays 19: 277-279
16. **Blazquez MA** (2000) Flower development pathways. J Cell Sci 113: 3547-3548
17. **Blazquez MA, Weigel D** (2000) Integration of floral inductive signals in Arabidopsis. Nature 404: 889-892
18. **Bonzon M, Greppin H, De Greef J** (1977) Contenu en ATP des feuilles de l'épinard sous différents régimes photopériodiques inducteurs ou non de la floraison. Arch Int Physiol Biochim 85: 953-955
19. **Bonzon M, Hug M, Wagner E, Greppin H** (1981) Adenine nucleotides and energy charge evolution during the induction of flowering in spinach leaves. Planta 152: 189-194
20. **Bonzon M, Simon P, Greppin H, Wagner E** (1983) Pyridine nucleotides and redox charge evolution during the induction of flowering in spinach leaves. Planta 159: 254-260
21. **Bünning E** (1973) The Physiological Clocks. Springer Verlag, Berlin
22. **Burn JE, Bagnall DJ, Metzger J, Dennis ES, Peacock WJ** (1993) DNA methylation, vernalization, and the initiation of flowering. Proc Natl Acad Sci USA 90: 287-291

23. **Bush SD** (1995) Calcium regulation in plant cells and its role in signalling. *Annu Rev Plant Physiol Plant Mol Biol* 46: 95-122
24. **Chailakhyan MKH, Loznikova VN** (1985) The florigen hypothesis and its substantiation by extraction of substances which induce flowering in plants. *Fiziol Rast* 32: 1172-1181
25. **Colasanti J, Sundaresan V** (2000) 'Florigen' enters the molecular age: long-distance signals that cause plants to flower. *Trends Biochem Sci* 25: 236-240
26. **Colasanti J, Yuan Z, Sundaresan V** (1998) The indeterminate gene encodes a zinc finger protein and regulates a leaf-generated signal required for the transition to flowering in maize. *Cell* 93: 593-603
27. **Crespi P, Crèvecoeur M, Lefort F, Greppin H** (1989) Changes in plasmalemma after gibberellic acid treatment. *Plant Sci* 62: 63-71
28. **Crespi P, Greppin H** (1994) Investigations par résonance magnétique nucléaire (RMN) sur la membrane plasmique de l'épinard pendant l'induction florale. *Archs Sci, Genève* 47: 273-286.
29. **Crespi P, Penel C, Crèvecoeur M, Greppin H** (1993) Plasma membrane sterols and flowering induction. *Plant Sci* 89: 153-160
30. **Crèvecoeur M, Lesniewska ME, Vié V, Goudonnet JP, Greppin H, Le Grimellec C** (2000) Atomic-force microscopy imaging of plasma membrane purified from spinach leaves. *Protoplasma* 212: 46-55
31. **Degli Agosti R, Bonzon M, Greppin H** (1989) Photoperiodic floral induction and glucose content changes in spinach, mustard and *Chenopodium rubrum* plants. *Bot Helv* 99: 73-79
32. **Degli Agosti R, Greppin H** (1987) Mise en évidence de deux types de contrôle du glucose chez l'épinard (*Spinacia oleracea*): analyse mathématique et modèle en jour court et lors de la variation de photopériode inductrice de la floraison. *Saussurea, Genève* 18: 45-54
33. **Degli Agosti R, Greppin H** (1998) Systemic stress effect on the sugar metabolism under photoperiodic constraint. *Archs Sci, Genève* 51: 337-346
34. **Dennis ES, Bilodeau P, Burn J, Finnegan EJ, Genger R, Helliwell C, Kang BJ, Sheldon CC, Peacock WJ** (1998) Methylation controls the low temperature induction of flowering in Arabidopsis. *Symp Soc Exp Biol* 51: 97-103
35. **Devlin PF, Kay SA** (2000) Flower arranging in Arabidopsis. *Science* 288: 1600-1602
36. **Douady S, Couder Y** (1992) Phyllotaxis as a physical self-organized growth process. *Phys Rev Lett* 68: 2098-2101
37. **Dunlap JC** (1999) Molecular bases for circadian clocks. *Cell* 96: 271-290
38. **Eden S, Hashimshony T, Keshet I, Cedar H, Thorne AW** (1998) DNA methylation models histone acetylation. *Nature* 394: 842
39. **El Toukhi Fouda M, Greppin H** (1988) G-6-PDH et évocation florale. *Saussurea, Genève* 19: 141-146
40. **Fagard M, Vaucheret H** (2000) Systemic silencing signal(s). *Plant Mol Biol* 43: 285-293
41. **Fankhauser C, Chory J** (1997) Light control of plant development. *Annu Rev Cell Dev Biol* 13: 203-229

42. **Finnegan EJ, Genger RK, Kovac K, Peacock WJ, Dennis ES** (1998) DNA methylation and the promotion of flowering by vernalization. *Proc Natl Acad Sci USA* 95: 5824-5829
43. **Finnegan EJ, Kovac KA** (2000) Plant DNA methyltransferases. *Plant Mol Biol* 43: 189-201
44. **Finnegan EJ, Peacock WJ, Dennis ES** (1996) Reduced DNA methylation in *Arabidopsis thaliana* results in abnormal plant development. *Proc Natl Acad Sci USA* 93: 8449-8454
45. **Finnegan EJ, Peacock WJ, Dennis ES** (2000) DNA methylation, a key regulator of plant development and other processes. *Curr Opin Genet Dev* 10: 217-223
46. **Fowler S, Lee K, Onouchi H, Samach A, J Putterill** (1999) GIGANTEA : a circadian clock-controlled gene that regulates photoperiodic flowering in *Arabidopsis* and encodes a protein with several possible membrane-spanning domains. *EMBO J* 18: 4679-4688
47. **Frosch S, Tsala G, Degli Agosti R, Greppin H** (1986) Effect of photoperiod length on leaf ontogenesis in spinach plants (*Spinacia oleracea*) as related to photosynthetic O₂ production, rubisco capacity and total sugar content. *Archs Sci, Genève* 39: 263-275
48. **Garab G, Lohner K, Laggner P, Farkas T** (2000) Self-regulation of the lipid content of membranes by non-bilayer lipids: a hypothesis. *Trends Plant Sci* 5: 489-494
49. **Gaspar T, Penel C, Degli Agosti R, Greppin H** (1987) Relations inter-organes, croissance, dominance apicale et floraison. *L'année biologique* 26: 81-89
50. **Gaspar T, Penel C, Greppin H** (1975) Peroxidase and isoperoxidases in relation to root and flower formation. *Plant Biochem J* 2: 33-47
51. **Genoud T, Métraux JP** (1999) Crosstalk in plant cell signaling : structure and function of the genetic network. *Trends Plant Sci* 4: 503-507
52. **Goodrich J, Puangsomlee P, Martin M, Long D, Meyerowitz EM, Coupland G** (1997) A Polycomb-group gene regulates homeotic gene expression in *Arabidopsis*. *Nature* 386: 44-51
53. **Greppin H** (1975) La floraison: ébauche d'une nouvelle stratégie. *Saussurea, Genève* 6: 252-254
54. **Greppin H** (1994) Des protéines comme organisatrices de la fréquence spécifique des rythmes circadiens ? In: B Millet, ed, *Mouvements, rythmes et irritabilité chez les végétaux*, Université de Franche-Comté, Besançon, pp 29-35
55. **Greppin H, Auderset G, Bonzon M, Degli Agosti R, Lenk R, Penel C** (1986) Le mécanisme de l'induction florale. *Saussurea, Genève* 17: 71-84
56. **Greppin H, Auderset G, Bonzon M, Penel C** (1978) Changement d'état membranaire et mécanisme de la floraison. *Saussurea, Genève* 9: 83-101
57. **Greppin H, Bonzon M, Crespi P, Crèvecoeur M, Degli Agosti R, Penel C** (1990) Physiological macrofunctions and indicators of the flowering process. In: B Millet, H Greppin, eds, *Intra- and Intercellular Communications in Plants*, INRA Paris, pp 107-123

58. **Greppin H, Degli Agosti R** (1997) Physiomatics and macrofunctions as a tool for development analysis. In: Greppin H, Penel C, Simon P, eds, Travelling Shot on Plant Development, Univ Geneva, Switzerland, pp 335-344
59. **Greppin H, Degli Agosti R, Bonzon M** (1996) Vistas on Biorhythmicity. Univ Geneva, Switzerland
60. **Greppin H, Degli Agosti R, Penel C** (1998) The Co-Action between Living Systems and the Planet, Univ Geneva, Switzerland
61. **Greppin H, Horwitz BA, Horwitz LP** (1973) Light stimulated bioelectric response of spinach leaves and photoperiodic induction. *Z Pflanzenphysiol* 68: 336-345
62. **Greppin H, Kayali S, Montavon M** (1996) Electrical signature and nyctophotoperiodic transition. In: Greppin H, Degli Agosti R & Bonzon M, eds, Vistas on Biorhythmicity, Univ Geneva, Switzerland, pp 269-278
63. **Greppin H, Penel C, Simon P** (1997) Travelling Shot on Plant Development. Univ Geneva, Switzerland
64. **Greppin H, Penel C, Tacchini P** (1995) Some indicators of flowering and floral stage. *Flowering News Lett* 19: 20-27
65. **Greppin H, Strasser R** (1983) Complexity and entropy changes in the photosynthetic apparatus during floral induction of spinach plants. In: Sybesma C, ed, *Advances in Photosynthesis Research*, Nijhoff M & Junk W Publ, pp III.1.59-62
66. **Greppin H, Wagner E, Degli Agosti R, Favre P** (1999) Activité électrique et floraison. *Archs Sci, Genève* 52: 29-40
67. **Guo H, Yang H, Mockler TC, Lin C** (1998) Regulation of flowering time by *Arabidopsis* photoreceptors. *Science* 279: 1360-1363
68. **Halevy AH** (1985) *Handbook of Flowering*. CRC Press, Boca Raton, Florida
69. **Hartmann U, Hohmann S, Nettesheim K, Wisman E, Saedler H, Huijser P** (2000) Molecular cloning of SVP: a negative regulator of the floral transition in *Arabidopsis*. *Plant J* 21: 351-360
70. **Hartwell LH, Hopfield JJ, Leibler S, Murray AW** (1999) From molecular to modular cell biology. *Nature* 402 (supp): C47-C52
71. **Haughn GW, Schultz EA, Martinez-Zapater JM** (1995) The regulation of flowering in *Arabidopsis thaliana*: meristems, morphogenesis and mutants. *Can J Bot* 73: 959-981
72. **Hempel FD, Welch DR, Feldman LJ** (2000) Floral induction and determination: where is flowering controlled ? *Trends Plant Sci* 5: 17-21
73. **Huq E, Tepperman JM, Quail PH** (2000) GIGANTEA is a nuclear protein involved in phytochrome signaling in *Arabidopsis*. *Proc Natl Acad Sci USA* 97: 9789-9794
74. **Jacobsen SE, Sakai H, Finnegan EJ, Cao X, Meyerowitz EM** (2000) Ectopic hypermethylation of flower-specific genes in *Arabidopsis*. *Curr Biol* 10: 179-186
75. **Jeddeloh JA, Stokes TL, Richards EJ** (1999) Maintenance of genomic methylation requires a SWI2/SNF2-like protein. *Nat Genet* 22: 94-97
76. **Johanson U, West J, Lister C, Michaels S, Amasino R, Dean C** (2000) Molecular analysis of FRIGIDA, a major determinant of natural variation in *Arabidopsis* flowering time. *Science* 290: 344-347

77. Jorgensen RA, Atkinson RG, Forster RL, Lucas WJ (1998) An RNA-based information superhighway in plants. *Science* 279: 1486-1487
78. Kakutani T, Jeddeloh JA, Richards EJ (1995) Characterization of an *Arabidopsis thaliana* DNA hypomethylation mutant. *Nucleic Acids Res* 23: 130-137
79. Kardailsky I, Shukla VK, Ahn JH, Dagenais N, Christensen SK, Nguyen JT, Chory J, Harison MJ, Weigel D (1999) Activation tagging of the floral inducer FT. *Science* 286: 1962-1965
80. Karege F, Penel C, Greppin H (1979) Reaction of a peroxidase activity to red and far red light in relation to the floral induction in spinach. *Plant Sci Lett* 17: 37-42
81. Karege F, Penel C, Greppin H (1982) A rapid correlation between the leaves of spinach and the photocontrol of peroxidase activity. *Plant Physiol* 69: 437-441
82. Karege F, Penel C, Greppin H (1982) Détection de l'état végétatif et floral de la feuille de l'épinard: emploi d'un indicateur biochimique. *Archs Sci, Genève* 35: 331-340
83. Kiefer S, Penel C, Greppin H, Wagner E (1987) Association de peroxydases aux membranes de courgette, de raifort et de *Pharbitis nil*. *Archs Sci, Genève* 40: 369-378
84. Kiyosue T, Ohad N, Yadegari R, Hannon M, Dinneny J, Wells D, Katz A, Margossian L, Harada JJ, Goldberg RB, Fischer RL (1999) Control of fertilization-independent endosperm development by the MEDEA polycomb gene in Arabidopsis. *Proc Natl Acad Sci USA* 96: 4186-4191
85. Kiyosue T, Wada M (2000) LKP1 (LOV kelch protein 1): a factor involved in the regulation of flowering time in Arabidopsis. *Plant J* 23: 807-815
86. Kobayashi Y, Kaya H, Goto K, Iwabuchi M, Araki T (1999) A pair of related genes with antagonistic roles in mediating flowering signals. *Science* 286: 1960-1962
87. Kondo T, Masahiro I (1999) The circadian clocks of plants and cyanobacteria. *Trends Plant Sci* 4: 171-176
88. Koornneef M, Alonso-Blanco C, Blankestijn-de Vries H, Hanhart CJ, Peeters AJ (1998) Genetic interactions among late-flowering mutants of Arabidopsis. *Genetics* 148: 885-892
89. Koornneef M, Alonso-Blanco C, Peeters AJM, Soppe W (1998) Genetic control of flowering time in Arabidopsis. *Annu Rev Plant Physiol Plant Mol Biol* 49: 345-370
90. Koornneef M, Hanhart CJ, Van Der Veen JH (1990) The genetic analysis of flowering time in *Arabidopsis thaliana*. *Flowering Newslett* 10: 9-12
91. Koornneef M, Hanhart CJ, Van der Veen JH (1991) A genetic and physiological analysis of late flowering mutants in *Arabidopsis thaliana*. *Mol Gen Genet* 229: 57-66
92. Krol E, Trebacz K (2000) Ways of ion channel gating in plant cells. *Ann Bot* 86: 449-469
93. Lang A (1984) Die photoperiodische Regulation von Förderung und Hemmung Blütenbildung. *Ber Dtsch Bot Ges* 97: 293-314

94. Lee H, Suh SS, Park E, Cho E, Ahn JH, Kim SG, Lee JS, Kwon YM, Lee I (2000) The AGAMOUS-LIKE 20 MADS domain protein integrates floral inductive pathways in Arabidopsis. *Genes Dev* 14: 2366-2376
95. Lee I, Aukerman MJ, Gore SL, Lohman KN, Michaels SD, Weaver LM, John MC, Feldmann KA, Amasino RM (1994) Isolation of LUMINIDEPENDENS: a gene involved in the control of flowering time in Arabidopsis. *Plant Cell* 6: 75-83
96. Lefort F, Greppin H (1992) Modifications du profil protéique dans les protoplastes d'apex d'épinard (*Spinacia oleracea*) lors de l'induction florale. *Archs Sci, Genève* 45: 69-84
97. Lefort F, Thiellement H, Greppin H (1991) Polypeptide pattern modifications during floral induction in leaves of *Spinacia oleracea*. *Archs Sci, Genève* 44: 227-234
98. Lenk R, Bonzon M, Greppin H (1980) Dynamically oriented biological water as studied by NMR. *Chem Phys Lett* 76: 175-177
99. Lenk R, Bonzon M, Greppin H (1981) Irreversible thermodynamics and biological evolution in spinach leaves as studied by NMR. *Z Pflanzenphys* 101: 107-118
100. Leshem Y (1992) *Plant Membranes*. Kluwer, Dordrecht
101. Levy YY, Dean C (1998) The transition to flowering. *Plant Cell* 10: 1973-1990
102. Levy YY, Dean C (1998) Control of flowering time. *Curr Opin Plant Biol* 1: 49-54
103. Lin C (2000) Plant blue-light receptors. *Trends Plant Sci* 5: 337-342
104. Lumsden PJ, Millar AJ (1998) *Biological Rhythms and Photoperiodism in Plants*. Bios Scientific Pub, Oxford
105. Ma H (1998) Flowering time: from photoperiodism to florigen. *Curr Biol* 8: R690-692
106. Macknight R, Bancroft I, Page T, Lister C, Schmidt R, Love K, Westphal L, Murphy G, Sherson S, Cobbett C, Dean C (1997) FCA, a gene controlling flowering time in Arabidopsis, encodes a protein containing RNA-binding domains. *Cell* 89: 737-745
107. Martinec J, Crèvecoeur M, Crespi P, Greppin H (1993) Plasma membrane thickness and flowering induction in *Chenopodium rubrum* and in *Spinacia oleracea*. *Archs Sci, Genève* 46: 233-248
108. Mc Clung RC (1998) It's about time: putative component of an Arabidopsis circadian clock. *Trends Plant Sci* 3: 454-456
109. McDaniel CN, Singer SR, Smith ME (1992) Developmental states associated with the floral transition. *Dev Biol* 153: 59-69
110. Meyerowitz EM (1989) Arabidopsis, a useful weed. *Cell* 56: 263-269
111. Michaels SD, Amasino RM (1999) FLOWERING LOCUS C encodes a novel MADS domain protein that acts as a repressor of flowering. *Plant Cell* 11: 949-956
112. Millet B, Greppin H (1990) *Intra-and Intercellular Communications in Plants*. INRA, Paris
113. Mizoguchi T, Coupland G (2000) ZEITLUPE and FKF1: novel connections between flowering time and circadian clock control. *Trends Plant Sci* 5: 409-411

114. Mockler TC, Guo H, Yang H, Duong H, Lin C (1999) Antagonistic actions of Arabidopsis cryptochromes and phytochrome B in the regulation of floral induction. *Development* 126: 2073-2082
115. Montavon M, Greppin H (1983) Effet sur le développement de l'épinard de l'application d'un potentiel électrique sur le pétiole d'une feuille. *Saussurea*, Genève 14: 79-85
116. Montavon M, Greppin H (1986) Développement apical de l'épinard et application d'un potentiel électrique de contrainte. *Saussurea*, Genève 17: 85-91
117. Morel JB, Vaucheret H (2000) Post-transcriptional gene silencing mutants. *Plant Mol Biol* 43: 275-284
118. Morré DJ, Morré D (1998) NADH oxidase activity of soybean plasma membranes oscillates with a temperature compensated period of 24 min. *Plant J* 16: 277-284
119. Morré DJ, Morré D, Penel C, Greppin H (1998) Auxin modulated protein disulfide thiol interchange activity from plasma membranes of spinach leaves responds to photoperiod and NADH. *Int J Plant Sci* 159: 105-109
120. Morré DJ, Morré D, Penel C, Greppin H (1999) NADH oxidase periodicity of spinach leaves synchronized by light. *Int J Plant Sci* 160: 855-860
121. Nelson DC, Lasswell J, Rogg LE, Cohen MA, Bartel B (2000) FKF1, a clock-controlled gene that regulates the transition to flowering in Arabidopsis. *Cell* 101: 331-340
122. Neuhaus G, Bowler C, Hiratsuka K, Yamagata H, Chua NH (1997) Phytochrome-regulated repression of gene expression requires calcium and cGMP. *EMBO J* 16: 2554-2564
123. Ng M, Yanofsky MF (2000) Three ways to learn the ABCs. *Curr Opin Plant Biol* 3: 47-52
124. Nikiforova VY, Sidorkina EV, Milyaeva EL (1999) The control of plant transition to flowering by cell-wall active compounds. *Russ J Plant Physiol* 46: 661-665
125. Novak B, Greppin H (1979) High-frequency oscillation and circadian rhythm of the membrane potential in spinach leaves. *Planta* 144: 235-240
126. Ohad N, Yadegari R, Margossian L, Hannon M, Michaeli D, Harada JJ, Goldberg RB, Fischer RL (1999) Mutations in FIE, a WD polycomb group gene, allow endosperm development without fertilization. *Plant Cell* 11: 407-416
127. Onouchi H, Igeno MI, Perilleux C, Graves K, Coupland G (2000) Mutagenesis of plants overexpressing CONSTANS demonstrates novel interactions among Arabidopsis flowering-time genes. *Plant Cell* 12: 885-900
128. Park DH, Somers DE, Kim YS, Choy YH, Lim HK, Soh MS, Kim HJ, Kay SA, Nam HG (1999) Control of circadian rhythms and photoperiodic flowering by the Arabidopsis GIGANTEA gene. *Science* 285: 1579-1582
129. Pelaz S, Ditta GS, Baumann E, Wisman E, Yanofsky MF (2000) B and C floral organ identity functions require SEPALLATA MADS-box genes. *Nature* 405: 200-203
130. Penel C, Gaspar T, Greppin H (1985) Rapid interorgan communications in higher plants with special reference to flowering. *Biol Plant* 27: 334-338

131. **Penel C, Greppin H** (1975) Photocontrôle immédiat de l'activité peroxydasique et corrélation entre les feuilles de l'épinard. *Saussurea*, Genève 6: 287-291
132. **Penel C, Greppin H** (1975) The balance between acid and basic peroxidases and its photoperiodic control in spinach leaves. *Plant Sci Lett* 5: 41-48
133. **Penel C, Greppin H** (1991) *Plant Signalling, Plasma Membrane and Change of State*. Univ Geneva, Switzerland
134. **Pickard BG, Beachy RN** (1999) Intercellular connections are developmentally controlled to help move molecules through the plant. *Cell* 98: 5-8
135. **Pidkowich MS, Klenz JE, Haughn GW** (1999) The making of a flower: control of floral meristem identity in *Arabidopsis*. *Trends Plant Sci* 4: 64-70
136. **Pineiro M, Coupland G** (1998) The control of flowering time and floral identity in *Arabidopsis*. *Plant Physiol* 117: 1-8
137. **Preuss D** (1999) Chromatin silencing and *Arabidopsis* development: A role for polycomb proteins. *Plant Cell* 11: 765-768
138. **Putterill J, Robson F, Lee K, Simon R, Coupland G** (1995) The *CONSTANS* gene of *Arabidopsis* promotes flowering and encodes a protein showing similarities to zinc finger transcription factors. *Cell* 80: 847-857
139. **Queiroz-Claret C, Lenk R, Queiroz O, Greppin H** (1988) NMR studies of *in vitro* slow oscillations in enzyme properties and dissipative structures. *Plant Physiol Biochem* 26: 333-338
140. **Ramade F** (1989) *Éléments d'écologie*. Mc Graw-Hill, Paris
141. **Reeves PH, Coupland G** (2000) Response of plant development to environment: control of flowering by daylength and temperature. *Curr Opin Plant Biol* 3: 37-42
142. **Roldan M, Gomez-Mena C, Ruiz-Garcia L, Salinas J, Martinez-Zapater JM** (1999) Sucrose availability on the aerial part of the plant promotes morphogenesis and flowering of *Arabidopsis* in the dark. *Plant J* 20: 581-590
143. **Sachs RM, Hackett WP** (1983) Source-sink relationships and flowering. In: WJ Mendt, ed, *Strategies of Plant Reproduction*, Allanheld, Osmun, N.Y
144. **Samach A, Coupland G** (2000) Time measurement and the control of flowering in plants. *Bioessays* 22: 38-47
145. **Samach A, Onouchi H, Gold SE, Ditta GS, Schwarz-Sommer Z, Yanofsky MF, Coupland G** (2000) Distinct roles of *CONSTANS* target genes in reproductive development of *Arabidopsis*. *Science* 288: 1613-1616
146. **Sarazin B, Chané-Favre L, Greppin H** (2000) Effet de la 5-azacytidine sur la floraison d'*Arabidopsis thaliana*. *Archs Sci*, Genève 53: 233-237
147. **Schaffer R, Ramsay N, Samach A, Corden S, Putterill J, Carre IA, Coupland G** (1998) The late elongated hypocotyl mutation of *Arabidopsis* disrupts circadian rhythms and the photoperiodic control of flowering. *Cell* 93: 1219-1229
148. **Schlattner U** (1996) L'adénylate kinase du tabac (*Nicotiana tabacum* L. cv Havana 425). Thèse No 2806, Université de Genève
149. **Schlattner U, Wagner E, Greppin H, Bonzon M** (1996) Change in distribution of chloroplast adenylate kinase isoforms during floral induction. *Physiol Plant* 96: 319-323

150. Schultz TF, Raman R, Somers DE, Mas P, Panda S, Kreps JA, Kay SA (2000) Cloning of the Arabidopsis clock gene TOC1, an autoregulatory response regulator homolog. *Science* 289: 768-771
151. Sessions A, Yanofsky MF, Weigel D (2000) Cell-cell signaling and movement by the floral transcription factors LEAFY and APETALA1. *Science* 289: 779-782
152. Sheldon CC, Burn JE, Perez PP, Metzger J, Edwards JA, Peacock WJ, Dennis ES (1999) The FLF MADS box gene: a repressor of flowering in Arabidopsis regulated by vernalization and methylation. *Plant Cell* 11: 445-458
153. Sheldon CC, Finnegan EJ, Rouse DT, Tadege M, Bagnall DJ, Helliwell CA, Peacock WJ, Dennis ES (2000) The control of flowering by vernalization. *Curr Opin Plant Biol* 3: 418-422
154. Sheldon CC, Rouse DT, Finnegan EJ, Peacock WJ, Dennis ES (2000) The molecular basis of vernalization: the central role of FLOWERING LOCUS C (FLC). *Proc Natl Acad Sci USA* 97: 3753-3758
155. Shepherd GM (1991) Sensory transduction: entering the mainstream of membrane signaling. *Cell* 67: 845-851
156. Simon R, Igeno MI, Coupland G (1996) Activation of floral meristem identity genes in Arabidopsis. *Nature* 384: 59-62
157. Simpson G, Gendall T, Dean C (1999) When to switch to flowering. *Annu Rev Cell Dev Biol* 15: 519-550
158. Smeekens S (2000) Sugar-induced signal transduction in plants. *Annu Rev Plant Physiol Plant Mol Biol* 51: 49-81
159. Snape JW, Quarrie SA, Laurie DA (1996) Comparative mapping and its use for the genomic analysis of agronomic characters in wheat. *Euphytica* 89: 27-31
160. Somers DE (1999) The physiology and molecular bases of the plant circadian clock. *plant physiology* 121: 9-19
161. Somers DE, Schultz TF, Milnamow M, Kay SA (2000) ZEITLUPE encodes a novel clock-associated PAS protein from Arabidopsis. *Cell* 101: 319-329
162. Soppe WJ et al. (2000) The late flowering phenotype of *fwa* mutants is caused by gain-of- function epigenetic alleles of a homeodomain gene. *Mol Cell* 6: 791-802
163. Steimer A, Amedeo P, Afsar K, Fransz P, Scheid OM, Paszkowski J (2000) Endogenous targets of transcriptional gene silencing in Arabidopsis. *Plant Cell* 12: 1165-1178
164. Stokes TL, Richards EJ (2000) Mum's the word. Mom and modifiers of transcriptional gene silencing. *Plant Cell* 12: 1003-1006
165. Strayer C, Oyama T, Schultz TF, Raman R, Somers DE, Mas P, Panda S, Kreps JA, Kay SA (2000) Cloning of the Arabidopsis clock gene TOC1, an autoregulatory response regulator homolog. *Science* 289: 768-771
166. Suarez-Lopez P, Coupland G (1998) Plants see the blue light. *Science* 279: 1323-1324
167. Sugano S, Andronis C, Green RM, Wang ZY, Tobin EM (1998) Protein kinase CK2 interacts with and phosphorylates the Arabidopsis circadian clock-associated 1 protein. *Proc Natl Acad Sci USA* 95: 11020-11025

168. Sung ZR, Belachew A, Shunong B, Bertrand-Garcia R (1992) EMF, an Arabidopsis gene required for vegetative shoot development. *Science* 258
169. Terry N, Rouze P (2000) The sense of naturally transcribed antisense RNAs in plants. *Trends Plant Sci* 5: 394-396
170. Thellier M, Le Sceller L, Norris V, Verdu MC, Ripoll C (2000) Long-distance transport, storage and recall of morphogenetic information in plants. The existence of a sort of primitive plant "memory". *CR Acad Sci Paris* 323: 81-91
171. Thomas B, Vince-Prue D (1997) Photoperiodism in Plants. Acad Press London
172. Tokarska M, Greppin H (1992). Fluctuation périodique de l'activité peroxydasique chez *Pseudomonas fluorescens* Mig. *Saussurea*, Genève 23: 83-87
173. Tokarska M, Greppin H, Penel C, Bonzon M (1993) In vitro slow fluctuation in horseradish peroxidase activity. *Chronobiol Int* 10: 238-243
174. Trebacz K, Stolarz H, Dziubinska H, Zawadski T (1997) Electrical control of plant development. In: H Greppin, C Penel & P Simon, eds, Travelling Shot on Plant Development, Univ Geneva, Switzerland, pp 165-181
175. Trewavas T (1999) How plants learn. *Proc Natl Acad Sci USA* 96: 4216-4218
176. Valat M, Greppin H (1992) Activité des différentes isozymes de la malate deshydrogénase dans la feuille de l'épinard (*Spinacia oleracea*): état végétatif et induction florale. *Archs Sci*, Genève 45: 191-206
177. Van Den Driessche T, Guisset JL, Petiau-De Vries GM (1996) Membranes and Circadian Rhythms. Springer Verlag, Berlin
178. Voinnet O, Baulcombe DC (1997) Systemic signalling in gene silencing. *Nature* 389: 553
179. Wagner E, Albrechtova JTP, Normann J, Greppin H (2000) Redox state and phosphorylation potential as macro-parameters in rhythmic control of metabolism-a molecular basis for seasonal adaptation of development. In: Vanden Driessche T, Guisset JL & Petiau de Vries GM, eds, The Redox state and Circadian Rhythms, Kluwer Acad Pub, The Netherlands, pp 257-277
180. Wagner E, Bonzon M, Greppin H (1985) Membrane-oscillator hypothesis of metabolic control in photoperiodic time measurement and the temporal organization of development and behaviour in plants. In: Packer L, ed, Recent Advances in Biological Membrane Studies, Plenum Pub, London, pp 525-546
181. Wagner E, Bonzon M, Normann J, Albrechtova JTP, Machackova I, Greppin H (1996) Signal transduction and metabolic control of timing in photoperiodism. The case of flower initiation. In: Greppin H, Degli Agosti R & Bonzon M, eds, Vistas on Biorhythmicity, Univ Geneva, Switzerland, pp 3-23
182. Wagner E, Fernandez SR, Normann J, Bonzon M, Greppin H (1993) Chronobiology: spatio-temporal organization of living systems. In: Greppin H, Bonzon M, Degli Agosti R, eds, Some Physico-Chemical and Mathematical Tools for Understanding of Living Systems, Univ Geneva, Switzerland, pp 109-126
183. Wagner E, Greppin H, Millet B (1987) The Cell Surface in Signal Transduction, Springer Verlag, Berlin
184. Wagner E, Normann J, Albrechtova JTP, Walczysko P, Bonzon M, Greppin H (1998) Electrochemical hydraulic signalling in photoperiodic control of

- flowering: is "florigen" a frequency-coded electric signal ? Flowering Newslett 26: 62-74
185. **Wang ZY, Tobin EM** (1998) Constitutive expression of the CIRCADIAN CLOCK ASSOCIATED 1 (CCA1) gene disrupts circadian rhythms and suppresses its own expression. *Cell* 93: 1207-1217
 186. **Weigel D** (1998) From floral induction to floral shape. *Curr Opin Plant Biol* 1: 55-59
 187. **Weigel D, Alvarez J, Smyth DR, Yanofsky MF, Meyerowitz EM** (1992) LEAFY controls floral meristem identity in *Arabidopsis*. *Cell* 69: 843-859
 188. **Weigel D, Meyerowitz EM** (1994) The ABCs of floral homeotic genes. *Cell* 78: 203-209
 189. **Weller JL, Reid JB, Taylor SA, C. MI** (1997) The genetic control of flowering in pea. *Trends Plant Sci* 2: 412-418
 190. **Woodward C** (1993) Is the slow-exchange core the protein folding core? *Trends Biochem Sci* 18: 359-360
 191. **Wu K, Malik K, Tian L, Brown D, Miki B** (2000) Functional analysis of a RPD3 histone deacetylase homologue in *Arabidopsis thaliana*. *Plant Mol Biol* 44:167-176
 192. **Xoconostle-Cazares B, Xiang Y, Ruiz-Medrano R, Wang HL, Monzer J, Yoo BC, McFarland KC, Franceschi VR, Lucas VJ** (1999) Plant paralog to viral movement protein that potentiates transport of mRNA into the phloem. *Science* 283: 94-98
 193. **Xue GX, Greppin H** (1994) Interaction par voie gazeuse et floraison. *Archs Sci, Genève* 47: 267-272
 194. **Yanofsky MF** (1995) Floral meristems to floral organs: genes controlling early events in *Arabidopsis* flower development. *Annu Rev Plant Physiol Plant Mol Biol* 46: 167-188
 195. **Zeevaart JAD** (1976) Physiology of flower formation. *Annu Rev Plant Physiol* 27: 321-348