



Article  
scientifique

Revue de la  
littérature

2023

Published  
version

Open  
Access

This is the published version of the publication, made available in accordance with the publisher's policy.

---

## SARS-CoV-2 viral load and shedding kinetics

---

Puhach, Olha; Meyer, Benjamin; Eckerle, Isabella Anne

### How to cite

PUHACH, Olha, MEYER, Benjamin, ECKERLE, Isabella Anne. SARS-CoV-2 viral load and shedding kinetics. In: Nature reviews. Microbiology, 2023, vol. 21, n° 3, p. 147–161. doi: 10.1038/s41579-022-00822-w

This publication URL: <https://archive-ouverte.unige.ch//unige:168711>

Publication DOI: [10.1038/s41579-022-00822-w](https://doi.org/10.1038/s41579-022-00822-w)

# SARS-CoV-2 viral load and shedding kinetics

Olha Puhach<sup>1</sup>, Benjamin Meyer<sup>2</sup> & Isabella Eckerle<sup>1,3,4</sup>✉

## Abstract

SARS-CoV-2 viral load and detection of infectious virus in the respiratory tract are the two key parameters for estimating infectiousness. As shedding of infectious virus is required for onward transmission, understanding shedding characteristics is relevant for public health interventions. Viral shedding is influenced by biological characteristics of the virus, host factors and pre-existing immunity (previous infection or vaccination) of the infected individual. Although the process of human-to-human transmission is multifactorial, viral load substantially contributed to human-to-human transmission, with higher viral load posing a greater risk for onward transmission. Emerging SARS-CoV-2 variants of concern have further complicated the picture of virus shedding. As underlying immunity in the population through previous infection, vaccination or a combination of both has rapidly increased on a global scale after almost 3 years of the pandemic, viral shedding patterns have become more distinct from those of ancestral SARS-CoV-2. Understanding the factors and mechanisms that influence infectious virus shedding and the period during which individuals infected with SARS-CoV-2 are contagious is crucial to guide public health measures and limit transmission. Furthermore, diagnostic tools to demonstrate the presence of infectious virus from routine diagnostic specimens are needed.

## Sections

Introduction

Measuring SARS-CoV-2 viral load

Viral load and shedding dynamics

SARS-CoV-2 transmission

SARS-CoV-2 diagnostics in public health

Conclusions

<sup>1</sup>Department of Medicine, Faculty of Medicine, University of Geneva, Geneva, Switzerland. <sup>2</sup>Centre for Vaccinology, Department of Pathology and Immunology, University of Geneva, Geneva, Switzerland. <sup>3</sup>Geneva Centre for Emerging Viral Diseases, Geneva University Hospitals, Geneva, Switzerland. <sup>4</sup>Division of Infectious Diseases, Geneva University Hospitals, Geneva, Switzerland. ✉e-mail: [Isabella.Eckerle@unige.ch](mailto:Isabella.Eckerle@unige.ch)

## Introduction

At the end of 2019, a novel coronavirus emerged, later termed severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which is the causative agent of coronavirus disease 2019 (COVID-19). SARS-CoV-2 primarily targets multiciliated cells in the upper respiratory tract (URT), but was also reported to infect cells outside the URT<sup>1</sup>. It can spread to the lower respiratory tract (LRT), where it infects alveoli, leading to reduced gas exchange, inflammation and pulmonary pathologies that are typical of COVID-19 (ref.<sup>2</sup>). Individuals who are infected shed the virus through the URT, with emission of infectious virus leading to secondary transmission and thus further spread of the virus.

Because of their nonspecific clinical presentation, precise diagnostic tools are needed to identify SARS-CoV-2 infections. Specific real-time PCR (RT-PCR) assays were quickly available after the emergence of the virus, later followed by antigen-detecting (rapid) diagnostic tests (Ag-RDTs) and serological assays. Although detection of viral RNA in respiratory specimens by RT-PCR is highly sensitive and specific, it does not distinguish between replication-competent virus and residual RNA. In the absence of a diagnostic test, infectiousness is often established using one of two proxies: the presence of viral RNA above a defined cycle threshold (Ct) value, or a positive Ag-RDT. RT-PCR is a useful tool for initial diagnosis, whereas Ag-RDTs can serve as an indicator for ending the isolation period. This is because viral RNA (which would be picked up by RT-PCR) remains detectable in the absence of infectious virus, whereas positivity of Ag-RDTs better correlates with the presence of infectious virus.

Aside from the respiratory tract, SARS-CoV-2 RNA has been detected in peripheral blood, stool, urine and ocular secretions<sup>3–7</sup>. Virus isolation from non-respiratory specimens was unsuccessful in most studies<sup>4,8,9</sup>, with very few reported cases of infectious virus presence in non-respiratory specimens<sup>10–13</sup>. Furthermore, viral loads from respiratory tract samples were found to be much higher than from other materials, the latter often with RNA viral loads that are incompatible with the presence of infectious virus. Such specimens are not considered relevant for transmission and therefore, we concentrate on SARS-CoV-2 virus shedding only through the respiratory tract.

Here, we elucidate the relationship between SARS-CoV-2 viral load and infectious virus presence, the biological and host factors that determine infectious virus shedding, measurement of infectious virus and the role diagnostics can have as a proxy for infectious virus shedding.

## Measuring SARS-CoV-2 viral load

The gold standard for laboratory diagnosis of a respiratory tract infection is demonstration of viral RNA with a virus-specific (semi-)quantitative RT-PCR from material collected from the respiratory tract. The most commonly used materials are swab specimens from the nasopharynx or oropharynx, but swabs of the nasal cavity, saliva or gargled liquid solution have also been suggested as alternative materials, with the advantage of being a less uncomfortable procedure for the participant. Viral load as determined by RT-PCR is either expressed as the number of viral RNA copies per millilitre of viral transport medium or per swab, or by the arbitrary test-specific Ct value. By contrast, infectiousness is determined by qualitative or quantitative assessment of infectious virus in a clinical specimen by replication of virus in cell culture. The limitations to measuring viral shedding are described in Box 1. In this Review, we refer to viral particles that can cause infection as infectious virus, and to viral RNA levels (which are widely used as surrogates for infectious virus) as viral load.

## Detection of infectious virus

The gold standard for determining the presence of infectious (that is, replication competent) virus in respiratory specimens is the recovery of virus in cell culture, a procedure that is commonly termed virus isolation (Fig. 1).

In the case of SARS-CoV-2, various cell lines and primary cells can be used for virus isolation, including those that express angiotensin-converting enzyme 2 (ACE2; the receptor required for virus entry) or transmembrane protease 2 (TMPRSS2; which is also important for virus entry)<sup>14</sup>. A cell line derived from African green monkey kidney cells, Vero E6, is commonly used for virus isolation, propagation and titration<sup>15</sup>. Other human cell lines that have been successfully used for SARS-CoV-2 isolation are a colorectal adenocarcinoma cell line (Caco-2), a lung adenocarcinoma cell line (Calu-3), a lung adenocarcinoma cell line ectopically overexpressing ACE2 (A549) and a human hepatocellular carcinoma cell line (Huh7)<sup>16,17</sup>.

The presence of infectious virus in the cell culture is qualitatively assessed using light microscopy, which can be used to identify cells undergoing the cytopathic effects (and death) caused by SARS-CoV-2 infection, consisting of syncytium formation, cell rounding, detachment and degeneration<sup>17</sup>. Infection is usually confirmed by a second method, either by a specific RT-PCR for viral RNA from the supernatant of infected cells, indicating virus replication by an increase of viral load over time in comparison to the baseline sample, or by immunostaining for viral proteins<sup>15,18</sup>.

This qualitative measurement of virus presence cannot, however, quantify the infectious virions in the inoculated specimens, although samples with lower viral load commonly show delayed development of a cytopathic effect<sup>19</sup>. Instead, methods such as plaque assays, focus-forming assays or 50% tissue culture infectious dose (TCID<sub>50</sub>) can be used to quantify infectious virus in a patient sample.

The above methodologies are reliable tools to detect infectious virus in clinical specimens of individuals who are infected with SARS-CoV-2, although there are limitations. Detection of viable virus particles is highly influenced by the quality of the sample, and infectious viral particles can quickly lose their infectiousness in unsuitable storage conditions. To preserve infectious virus in specimens, swab samples from patients infected with SARS-CoV-2 should be immediately submerged in a viral transport medium suitable for cell culture and stored at  $-80^{\circ}\text{C}$  as early as possible after collection. Prolonged exposure to higher temperatures or repeated freeze–thaw cycles can drastically influence the quality of the sample, leading to potentially complete loss of infectious viral particles. Therefore, many factors can influence the reproducibility of the results between different laboratories. Furthermore, cell lines used for isolation can show a high variability between laboratories even when they are presumably the same. Consumables used during cell culture, such as culture medium or additives such as fetal bovine serum and antibiotics, could potentially also impact virus isolation success. In human primary airway epithelial cells, which mimic the primary site of entry in the human respiratory tract, the probability of isolating infectious virus was reduced compared with that of Vero E6 cells, indicating that infectious virus determined using Vero E6 cells might be overestimated for assessing transmission risks *in vivo*<sup>20</sup>.

Importantly, all cell culture work with SARS-CoV-2 is done under biosafety level 3 conditions, so only specially trained personnel in laboratories with advanced infrastructure can perform these experiments. Thus, detection of viable virus through virus isolation is not suitable for diagnostics and is restricted to research only.

## Box 1

### Limitations to measuring viral load

#### Specimen selection site

The anatomical site chosen to collect the swab specimen for detection of SARS-CoV-2 might influence viral load detection. Higher RNA viral load was reported from nasopharyngeal than oropharyngeal swabs<sup>28,124,181</sup>. As a result, nasopharyngeal samples show the highest diagnostic accuracy compared with other upper respiratory tract samples<sup>182</sup>. Similarly, higher virus isolation success was reported from nasopharyngeal swabs than from saliva, nasal or sublingual swabs<sup>124</sup>. However, another study found higher RNA viral loads in the throat and sputum than from nasal swabs<sup>30</sup>. Two studies indicate that virus can be detected earlier in the throat<sup>29</sup> or saliva<sup>33</sup>, but reaches significantly higher levels and remains detectable for longer in the nose<sup>29,33</sup>. A meta-analysis, which evaluated different clinical sampling methods using nasopharyngeal swab as a reference, demonstrated that pooled nasal and throat swabs showed the best diagnostic performance<sup>183</sup>. Notably, this analysis revealed higher heterogeneity of results in studies using nasal or saliva specimens than using pooled nasal and throat swabs<sup>183</sup>.

The effect of the swabbing method (self-administered or performed by trained person) on measured viral loads cannot be overlooked: the sensitivity of antigen-detecting (rapid) diagnostic tests achieved by health-care professionals was higher than for self-tests<sup>57,184</sup>.

#### Impact of individual infection kinetics

To date, there is a limited number of studies that describe the longitudinal dynamics of SARS-CoV-2 shedding<sup>29,33</sup>. Most of the studies used only a single time point to collect respiratory swabs from individuals who were infected for measurement of viral load.

As a result, different times from symptom onset can be a confounding factor when comparing viral load between different patients, which might also explain the variation of available data on viral load.

#### Influence of epidemic period

RNA viral loads across the specimens collected at single time points were found to indicate the trajectory of the epidemic, as a high proportion of individuals who were recently infected with low cycle threshold values correlates with a higher reproduction number, indicative of a growing epidemic<sup>185</sup>. Similarly, the rise and fall of RNA viral load correlated with the number of COVID-19 cases and hospital admissions across the population as it was identified using SARS-CoV-2 RNA measurement in wastewater samples<sup>186</sup>. Moreover, variation in estimates of the mean incubation period was shorter before the epidemic peak in China than after the peak<sup>187</sup>. Therefore, sampling at single time points can be biased by the epidemic period and might reflect more epidemiological dynamics than individual shedding kinetics.

#### Influence of SARS-CoV-2 variant of concern

Available data on SARS-CoV-2 variants of concern demonstrated that, although the overall pattern of viral load dynamics is conserved between the variants, infection with different SARS-CoV-2 variants of concern led to highly distinct infectious virus amounts and RNA viral loads<sup>25,86,87,90,92,94</sup> and variations in SARS-CoV-2 incubation period<sup>113</sup>. Therefore, extrapolation of our understanding from shedding of current or earlier SARS-CoV-2 variants to newly emerged variants may be of only limited value.

#### Detection of RNA viral load


Techniques for detecting viral RNA by RT-PCR were quickly established at the beginning of the pandemic<sup>21,22</sup> (Fig. 1). The high specificity and sensitivity of RT-PCR make it the gold standard for diagnosing SARS-CoV-2 infections. Quantitative RT-PCR assays provide a Ct value, which is inversely correlated with the concentration of the target viral RNA in the clinical sample (that is, the higher the value, the lower the target RNA in the sample). By using an external standard with a defined number of RNA copies, Ct values can be transformed into absolute viral RNA copy numbers or international units per millilitre of viral transport medium or per total swab.

Although RT-PCR cannot directly determine infectiousness owing to its inability to differentiate between replication-competent (infectious) virus and residual (non-infectious) viral RNA, a correlation between RNA viral load and the presence of infectious virus has been sought. Several studies have attempted to correlate the quantity of viral RNA with infectiousness by isolating virus across a range of Ct values. Indeed, there was a stepwise decrease in the probability of virus isolation with increasing Ct values in samples collected during the first 8 days post-onset of symptoms (dpos)<sup>18,23</sup>. However, other studies have found that the correlation between infectious virus and RNA viral load was low and that viral load (or Ct values as a proxy) is only

a weak predictor of infectious virus presence in the first 5 dpos<sup>4,20,24,25</sup>. Furthermore, when taking a certain Ct value or RNA copy number as a threshold, it is not possible to determine whether the RNA viral load is increasing or already decreasing; therefore, a low viral load could be measured at the end of infection or in the early (pre-)symptomatic phase before reaching peak viral load.

In a routine diagnostic context, analytical sensitivity and limits of detection may vary between the tests and laboratories where they are applied. An analytical performance comparison between different RT-PCR assays showed variation between the measured Ct values and the detection rate<sup>26</sup>. Therefore, application of RNA standards and calculation of RNA genome copy number based on a standard curve can improve comparability between laboratories and assays. To facilitate easier calibration and control of nucleic acid amplification techniques, an international standard with assigned potency in the form of an inactivated SARS-CoV-2 isolate was introduced by the World Health Organization (WHO)<sup>27</sup>.

As with the detection of infectious virus, several other parameters can influence whether viral load can be detected. The site of specimen collection can impact the findings on viral load; although some studies report higher RNA viral load in nasal or nasopharyngeal swabs<sup>28,29</sup>, others show higher RNA viral load in throat samples<sup>30</sup>. Moreover, the transport



**Fig. 1 | Methods to measure infectious virus and RNA viral load.** Swab specimens from the nasopharynx or oropharynx are used for detection of SARS-CoV-2 viral loads. Detection of viral nucleic acids (RNA) is performed by quantitative real-time PCR (qRT-PCR). Viral RNA is extracted from lysed virus, reverse transcribed and amplified by qPCR using primers specific for one or more target regions in the viral genome. The amplification cycle at which samples cross the threshold (cycle threshold) defines the amount of viral RNA. RNA viral load can be expressed as the number of viral RNA copies per millilitre, or by the arbitrary test-specific cycle threshold value. Lateral flow assays detect the presence of specific viral proteins in the lysed viral particles. SARS-CoV-2 nucleocapsid is used in most antigen-detecting (rapid) diagnostic tests. The presence of infectious (replication-competent) virus in respiratory specimens can only be determined by the recovery of virus in cell culture by isolation or by quantification of infectious virus titres using 50% tissue culture infectious

dose (TCID<sub>50</sub>), focus-forming assays or plaque-forming assays. Virus isolation is performed by applying infectious medium on the monolayer of cells; isolation success is determined by the presence of a cytopathic effect approximately 3–5 days post-infection. White colour indicates the presence of a cytopathic effect in cells. For quantification of infectious virus titres, serial dilutions of respiratory samples are performed and used for inoculation on the monolayer of cells. In TCID<sub>50</sub>, 3–5 days post-infection, viral-induced cytopathic effect is classically defined using microscopy. In focus-forming assays, cells are fixed 1 day post-infection and immunostaining with virus-specific antibodies is performed to detect groups of infected cells (foci). The foci, indicating the presence of infectious virus, are displayed in blue. In plaque-forming assays, plates are fixed 2–3 days post-infection and stained with crystal violet; wells with individual plaques are used to determine viral titres. The plaques, indicating the presence of infectious virus, are displayed in white.

media used for the sample, storage condition and quality of the sample may further influence the detection of viral RNA and their usefulness and limitations when extrapolating to potential infectiousness.

Although new variants have impacted some gene targets, in most instances, they did not have a major effect on molecular diagnostics, owing to the use of dual-target assays (in which at least two viral genes are detected simultaneously)<sup>31</sup>.

## Antigen-detecting rapid diagnostic tests

Most lateral flow tests are designed to detect SARS-CoV-2 nucleocapsid protein, as a proxy for infectious virus, in nasal or nasopharyngeal swabs<sup>32–36</sup> (Fig. 1). Indeed, most studies on Ag-RDT detection show good concordance with RT-PCR positivity when Ct values are below 25–30, a viral load compatible with the presence of infectious virus, whereas higher Ct values give less reliable results<sup>34,37–41</sup>.

Early time points during infection often give negative results with Ag-RDT in individuals who have tested positive by PCR<sup>29,42</sup>. On average, the first positive Ag-RDT results are obtained about 1–2 days later than positive PCR results<sup>37</sup>, whereas the highest sensitivity in patients was shown during the first 7 dpos in the studies with ancestral SARS-CoV-2 (refs.<sup>42–44</sup>). Antigen tests show highest sensitivity for specimens containing infectious virus and with Ct values below 25 (refs.<sup>45–49</sup>), and their positivity highly correlates with the presence of infectious virus<sup>34,45,47,50</sup>. By contrast, Ag-RDTs are less sensitive to low RNA viral loads (which have higher Ct values)<sup>51</sup>. Several studies have demonstrated a strong correlation between Ag-RDT positivity and the period in which infectious virus can be detected, indicating that Ag-RDTs can add an additional safety layer for deciding when to end isolation<sup>29,39</sup>.

However, some inconsistencies between studies and tests have been noted. For instance, there have been reports (across a range of studies and Ag-RDTs) of failure to detect viral antigens in specimens with a low Ct value and/or containing infectious virus (beyond the early acute phase)<sup>46,50</sup>. Moreover, there are seldom reports of Ag-RDTs remaining positive after more than 10 dpos<sup>42,46,50</sup>. As most studies failed to isolate infectious virus after more than 10 dpos, it remains unclear whether Ag-RDT positivity beyond 10 dpos correlates with infectious virus shedding. One study showed that antigen tests predict infectiousness more accurately at 1–5 dpos, than at 6–11 dpos<sup>52</sup>. Notably, there was a good correlation between Ag-RDT positivity and infectious virus isolation within the first 11 dpos<sup>52</sup>.

Conflicting results were found for sensitivity and specificity of Ag-RDTs for detection of SARS-CoV-2 variants, with large variations between manufacturers, the type of setting in which the Ag-RDTs were used (self-tests versus tests collected by a health-care professional) and the type of sample used for detection (nasal versus oral)<sup>53–57</sup>. With increasing hybrid immunity and the presence of mucosal antibodies, Ag-RDTs may further lose sensitivity<sup>58</sup>.

## Viral load and shedding dynamics

Viral loads are used as a proxy to characterize infectious viral shedding. The exact time for which individuals remain infectious is laborious to estimate and is likely to vary between patients. Viral factors, such as viral variant, and host factors, such as patient age and sex and immune status, influence shedding dynamics.

## Viral load as a key determinant of viral shedding


After the emergence of SARS-CoV-2 in late 2019, the first details on viral load and infectious virus shedding were measured in a cluster of infections that occurred in January 2020 in Germany, assessing

nine immunocompetent individuals with a mild course of disease<sup>4</sup>. Peak RNA viral loads were reached in the early symptomatic period at 5 dpos, a finding that was confirmed by other studies reporting peak viral loads at the time of symptom onset or even shortly before<sup>4,7,28,59</sup>. RNA viral loads gradually declined over the course of the disease in the nasopharyngeal and throat swabs, reaching low or undetectable levels 2 weeks after symptom onset<sup>4,23,59,60</sup> (Fig. 2). Declining RNA viral load is associated with resolution of clinical symptoms and gradual increase in antibody titres, for both binding and neutralizing antibodies<sup>18,23</sup>. However, ongoing detection of viral RNA has been described for prolonged periods up to 28 dpos in otherwise healthy individuals<sup>61</sup>, and some studies have reported low-level detection of RNA by RT-PCR even for months<sup>62</sup>. Participants who continue to shed viral RNA for more than 4 weeks after initial detection by RT-PCR represent a minority of non-severe cases, estimated to be around 3%<sup>63</sup>, 14%<sup>64</sup> or less than 20%<sup>65</sup>.

Infectious virus shedding of the ancestral SARS-CoV-2 strain, as determined by virus isolation in cell culture, was reported to correlate with high RNA viral load in the early acute phase after symptom onset<sup>23</sup>. Importantly, daily longitudinal sampling of respiratory specimens from individuals with mild disease or asymptomatic infection revealed that infectious virus can already be detected before the onset of symptoms<sup>33</sup>. Successful infectious virus isolation was reported within the first 8–10 dpos, but culture probability after this time period rapidly declined<sup>4,7,23,29,66,67</sup>. Studies that assessed infectious virus quantitatively found that infectious virus titres declined over the first 10 dpos<sup>25,29</sup>. In addition, a reduced chance of virus isolation coincided with the time of seroconversion in hospitalized patients and, as a result, infectious virus could no longer be isolated from seroconverted patients with detectable antibody titres<sup>18,68,69</sup>. Although similar seroconversion studies performed on mildly symptomatic patients are missing, the number of immunologically naive individuals is declining and this broadly existing underlying immunity makes such an assessment more complex.

Most studies on infectious virus shedding in the acute symptomatic period were on immunocompetent patients that had mild-to-moderate disease, representing the majority of COVID-19 cases in the community. Therefore, the assessment of the presence of infectious virus in the URT from those studies was used to define the duration of the period of infectiousness and contributed to best public health practices for isolation and quarantine<sup>62,70</sup>. Although the pattern of infection is broadly similar in patients with mild and severe disease, key differences do exist. The first week of illness is comparable in terms of RNA viral load between patients with mild and severe disease. However, patients with severe disease have elevated RNA viral loads in the second week of illness, and RNA was detected for prolonged periods<sup>71</sup>. Moreover, infectious virus was recovered from hospitalized patients for prolonged periods of up to 32 dpos<sup>18,72,73</sup>; however, the median time from symptom onset to viral clearance in culture was similar to that of patients with mild or moderate disease<sup>18,73</sup>. Severe COVID-19 is also characterized by high and persistent RNA viral load in the LRT, whereas non-severe cases have similar viral loads in the URT and LRT<sup>74</sup>.

Prolonged detection of viral RNA was also reported in immunocompromised patients; for example, 224 days after the beginning of the infection, virus was still detected in a man infected with HIV, including the detection of subgenomic RNA (sgRNA) indicating active viral replication<sup>75</sup>. Also, infectious virus was recovered up to 61 dpos in nasopharyngeal swabs collected from immunocompromised patients<sup>76</sup>, and low RNA viral loads were still detected at 60 dpos in another study<sup>77</sup>. Infectious virus was isolated from bronchoalveolar fluids from patients receiving chimeric antigen receptor (CAR) T cell therapy



**Fig. 2 | Kinetics of RNA viral loads and infectious virus for ancestral SARS-CoV-2 in patients with mild-to-moderate disease.** According to different studies, the incubation period for ancestral SARS-CoV-2 was estimated to lie between 4.6 and 6.4 days. On average, symptoms continue to persist for 10 days. RNA can already be detected before the onset of symptoms; RNA levels peak around the onset of symptoms and then gradually decline. Median clearance for RNA viral load is 16 days post-onset of symptoms. Infectious virus titres are highest around symptom onset, and infectious virus can be isolated up to 8 or 10 days post-onset of symptoms. RNA can be detected for prolonged periods

by real-time PCR, when infectious virus is no longer detectable, whereas virus detection by antigen-detecting (rapid) diagnostic tests (Ag-RDTs) was shown to be a better correlate for infectiousness. Gradients reflect variability between individuals (lighter shade towards the end of infection shows that viral loads continue to be detected in some but not all individuals). The grey dashed line marks the initial infection, the blue dashed lines mark the PCR-positive period and the red dashed lines mark Ag-RDT positivity. Details of the underlying studies used to generate Fig. 2 can be found in Supplementary Table 1.

up to 28 days after admission to an intensive care unit<sup>78</sup>. A case report on an immunocompromised patient showed isolation of infectious virus up to 78 dpos<sup>79</sup>. The reports of infectious virus isolation from severely ill or immunocompromised patients are limited (owing to the low number of patients), so it is difficult to define the proportion of cases with prolonged shedding.

The characteristics of viral shedding of other respiratory viruses are outlined in Box 2.

### Viral shedding of SARS-CoV-2 variants

Viral evolution of SARS-CoV-2 over time has led to the emergence of numerous variants. Combined with increasing population immunity due to vaccination or natural infection, this has led to a need to reassess our knowledge of viral shedding patterns.

The WHO designated variants as variants of concern (VOCs) if they were associated with one or more of the following: elevated transmissibility or a detrimental change in COVID-19 epidemiology; increased virulence or a change in clinical disease presentation; or decreased effectiveness of public health measures or available diagnostics, vaccines or therapeutics<sup>80</sup>. To date, five VOCs are recognized: Alpha, Beta, Gamma, Delta and Omicron. In contrast to ancestral SARS-CoV-2, VOCs display some differences in evasion from immunity, viral loads, shedding period or even incubation period, resulting in drastically different levels of transmission<sup>81–85</sup> (Fig. 3).

All VOCs have shown changes in viral load compared with ancestral SARS-CoV-2. One study reported that infection with Alpha leads to approximately tenfold higher RNA viral load and an increased probability of cell culture isolation compared with the ancestral virus<sup>86</sup>.

However, another study did not find a substantial difference in the infectious virus titre between Alpha and ancestral SARS-CoV-2 (ref.<sup>33</sup>). Delta reportedly led to an even higher increase in RNA viral load: one study reported a 1,000× increase relative to the ancestral virus<sup>87</sup>, and other studies reported 1.7× (ref.<sup>88</sup>) or 6.2× higher<sup>89</sup> viral load than Alpha. Furthermore, Delta demonstrated elevated probability of cell culture isolation<sup>90</sup> and higher infectious virus titres than Alpha<sup>91</sup>. Although Omicron was shown to be highly transmissible, lower RNA viral loads<sup>92</sup>, lower cell culture isolation probability<sup>93</sup> and lower infectious virus titres<sup>25</sup> were observed in patients infected with Omicron BA.1 than in those infected with Delta. Even within the Omicron clade, there are differences between sub-lineages, with infection with Omicron BA.2 leading to higher levels of RNA viral loads and longer time to viral clearance than with Omicron BA.1 (refs.<sup>94–96</sup>).

Similarly, VOCs have shown differences in the duration of viral shedding. Analysis of Ct values in respiratory specimens found that Delta showed longer persistence of viral RNA than ancestral SARS-CoV-2 (ref.<sup>97</sup>). Another study demonstrated that there was not significant difference in the mean duration of viral RNA presence in Delta and Omicron BA.1 infections<sup>92</sup>. The duration of infectious virus shedding appears to be similar to that observed with ancestral SARS-CoV-2, with culturable virus obtained at 5 dpos<sup>85</sup> and no replication-competent virus isolated beyond 10 dpos in patients infected with Delta and Omicron BA.1 (refs.<sup>84,98</sup>). It is important to note that pre-existing immunity to SARS-CoV-2, either from infection or vaccination, might influence the duration of infectious virus shedding (alongside immune status and disease severity, as discussed above), which may have driven some of these differences during the course of the pandemic.

## Influence of age and sex on viral shedding

There is some evidence that age-associated and sex-associated differences in innate and adaptive immunity, as well as higher ACE2 expression in adults than in children, result in an increased risk for severe disease in older male patients<sup>99–101</sup>. Moreover, a few studies have found that age and sex influence viral loads and shedding dynamics. In cases of infection with ancestral SARS-CoV-2, resolution of RNA shedding was faster in participants <18 years of age and slower in participants >50 years of age<sup>61</sup>. According to one study, viral RNA can be detected for longer times in male patients infected with ancestral SARS-CoV-2 (ref.<sup>102</sup>), and RNA viral loads were elevated in male patients infected with either Alpha or Delta variants compared with female patients<sup>88</sup>. However, a possible association of viral load dynamics with age or sex is highly debated, as other studies demonstrated that they have no influence on infectious virus<sup>25</sup> or RNA viral loads<sup>59</sup>.

Early studies in ancestral SARS-CoV-2 did not find a difference in virus isolation success<sup>103</sup> or RNA viral loads between children and adults<sup>104–106</sup>, but sample sizes were small. Slightly lower RNA viral loads and a more rapid clearance of viral RNA was observed in children than in adults when analysing much larger cohorts, whereas the patterns of shedding curves over time were similar between children and adults<sup>107</sup>. Furthermore, large-scale analysis of viral loads across different age groups showed no differences of distribution of RNA viral load between children and adults<sup>108</sup> or only slightly lower viral loads (<0.5 log<sub>10</sub> units) in children <5 years of age<sup>86</sup>.

## Symptoms as a correlate for shedding

One of the key epidemiological parameters for SARS-CoV-2 transmission is the incubation period, defined as the time from exposure or infection to the onset of symptoms. Studies on ancestral SARS-CoV-2

## Box 2

### Shedding of respiratory viruses

The dynamics of viral shedding differs between respiratory viruses, which influences their transmission and has an effect on diagnostics and measures applied to contain the outbreaks.

#### SARS-CoV

The epidemic of severe acute respiratory syndrome coronavirus (SARS-CoV) started in November 2002 in the Guangdong province of China and rapidly spread outside China. The virus was airborne and could also be spread via droplets of saliva, but is only moderately transmissible among humans<sup>188</sup>. Only low viral loads were detected in the early symptomatic period, generally peaking in the upper respiratory tract (URT) around 10–14 days post-onset of symptoms (dpos)<sup>189,190</sup>, and then dropping to low levels at 3–4 weeks post-infection<sup>191</sup>. In patients infected with SARS-CoV, viral RNA was detectable for a maximum of 8 weeks in samples collected from the URT<sup>191</sup> and for 52 days in sputum samples<sup>192</sup>, whereas infectious virus was isolated up to 28 dpos from stool and respiratory specimens and up to 36 dpos from urine samples<sup>191,193</sup>. SARS-CoV replicated less efficiently at low temperatures; thus, virus replication was more efficient in the lower respiratory tract (LRT) than in the URT<sup>194</sup>. Notably, asymptomatic or pre-symptomatic viral shedding and transmission were not recorded for SARS-CoV<sup>190,195</sup>; the peaks of transmission occurred around 2 and 10 dpos<sup>195</sup>. As a result, outbreaks were successfully contained through isolation of symptomatic patients infected with SARS-CoV, which reduced onward transmission<sup>196</sup>.

#### MERS-CoV

Middle East respiratory coronavirus (MERS-CoV) was isolated from a patient with pneumonia in Saudi Arabia in 2012 and was shown to be the causative agent of a cluster of severe respiratory tract infections in the Middle East<sup>197</sup>. The disease caused by MERS-CoV is characterized by a wide range of clinical severities and by predominantly respiratory symptoms, such as acute viral pneumonia, with a high case fatality ratio<sup>198</sup>. The virus is capable of

airborne transmission and has low transmissibility among humans, with a maximum estimated reproduction number below 1 (ref.<sup>198</sup>). Higher RNA viral loads were detected in the LRT than in the URT. Estimated mean shedding duration is 15.3 days in the URT and 16.3 days in the LRT<sup>62</sup>. Prolonged PCR positivity and higher RNA viral loads in the URT and LRT were associated with increased disease severity<sup>62,199</sup>. Viral RNA was also detected in the urine, stool and serum<sup>200</sup>. One study reported detection of viral RNA in the blood for 34 days and showed that presence of viral RNA in the blood is associated with higher mortality<sup>201</sup>; however, another study failed to isolate virus from PCR-positive serum samples<sup>200</sup>.


#### Influenza virus

In symptomatic patients, RNA viral loads start to be detectable by real-time PCR 2 days before the onset of symptoms and peak at 1 dpos<sup>202</sup>. Human challenge trials with influenza A viruses show that viral loads already sharply increase at 1 day post-inoculation, reach a peak at 2 days post-inoculation and become undetectable at 8 days post-inoculation. The mean duration of viral shedding for influenza viruses is 4.8 days, and the maximum duration is between 6 and 7 days<sup>203,204</sup>. Kinetics of infectious viral titres were similar to the viral load trends detected by real-time PCR for different strains of influenza<sup>205</sup>. Lower RNA viral loads and shorter infectious viral shedding were noted in asymptomatic patients<sup>202</sup>.

#### Human respiratory syncytial virus

This virus is the most frequent causative agent of LRT infections, leading to morbidity and mortality particularly in young children and older adults<sup>206</sup>. The virus is transmitted by contact with nasal secretions or large aerosols. Viral loads and symptoms increased simultaneously, reaching a peak at 5.4 days<sup>207</sup>. In human challenge trials, respiratory syncytial virus titres were detectable for an average of 4.6 days. Viral RNA could be still detected up to 9 dpos, whereas infectious virus titres could be detected from 1 to 8 dpos in adults<sup>208</sup> and up to 9 dpos in children<sup>209</sup>.





**Fig. 3 | Infectious viral load and symptom onset in SARS-CoV-2 Delta and Omicron BA.1 variants of concern.** Overall patterns of shedding dynamics are conserved between SARS-CoV-2 variants. In comparison to ancestral SARS-CoV-2, Delta and Omicron BA.1 have shorter incubation periods, estimated as approximately 3.7–4 days for Delta and approximately 3–3.4 days for Omicron BA.1. Higher infectious viral loads were detected in patients infected with Delta than in patients infected with Omicron BA.1 or ancestral SARS-CoV-2. Only

a limited number of studies have determined when virus shedding for Delta and Omicron BA.1 ends, so this time point is not well defined. Owing to the low number of studies comparing the end of the infectious period between different SARS-CoV-2 variants of concern, the end point of infectivity is not well defined (shown as a colour gradient). Details of the underlying studies used to generate Fig. 3 can be found in Supplementary Table 2.

have estimated that the incubation period on average is between 4.6 and 6.4 days<sup>59,109–111</sup> (Fig. 2). A human challenge trial with ancestral SARS-CoV-2 demonstrated that symptoms start to appear 2–4 days after inoculation, and RNA viral loads reach their peak 4–5 days after inoculation<sup>29</sup>. Thus, artificial inoculation of the virus confirmed the timing of peak viral loads observed in naturally infected individuals, whereas onset of symptoms was faster in the human challenge cases. In contrast to natural infection, in artificial inoculation, virus-containing drops with high viral load are directly applied in the nose and therefore reach the nasal epithelium more quickly, which might lead to the more rapid appearance of symptoms. For Delta, the estimated incubation period was between 3.7 and 4 days<sup>81–83,97</sup>, whereas infection with Omicron BA.1 was characterized by an even shorter incubation period of 3–3.4 days<sup>83,112,113</sup> (Fig. 3). However, as the time point of infection is rarely known outside of human challenge trials, dpos is most commonly used when analysing viral load and infectious virus.

Considering that high viral loads can be detected in the URT of infected individuals regardless of their clinical manifestations, the presence of symptoms is an unreliable indicator of infectiousness. Notably, individuals infected with SARS-CoV-2 can be infectious before the onset of symptoms<sup>59</sup>, and it was estimated that about half of secondary transmissions take place in the pre-symptomatic phase<sup>59,114</sup>. Moreover, according to population surveys, asymptomatic cases represent around 40% of all SARS-CoV-2 infections with ancestral SARS-CoV-2 (refs. <sup>115–117</sup>), and tracing of close contacts of confirmed cases of SARS-CoV-2 found that up to 23% of infections were asymptomatic<sup>118</sup>.

There are conflicting findings regarding viral shedding differences in symptomatic and asymptomatic patients. Comparison of viral loads between symptomatic and asymptomatic patients remains challenging, as the time of exposure cannot be clearly identified in asymptomatic individuals, and dpos cannot be used when comparing

viral loads with symptomatic individuals. Furthermore, individuals who do not show clinical symptoms at the time of testing can represent either true asymptomatic individuals or pre-symptomatic individuals who will develop symptoms later. Thus, only well-controlled studies with a follow-up of assessed individuals can make a clear distinction between pre-symptomatic and asymptomatic individuals. A study on ancestral SARS-CoV-2, which followed COVID-19 confirmed cases hospitalized for isolation and recorded symptoms daily, found similar initial Ct values between asymptomatic and symptomatic individuals<sup>119</sup>. Similarly, no significant difference in RNA viral loads between symptomatic and asymptomatic patients was found in other studies in which patients were followed longitudinally and the presence of symptoms was either monitored by health-care professionals<sup>120</sup> or was self-reported<sup>115</sup>. By contrast, other studies, in which symptoms were also recorded by clinicians, reported lower RNA viral loads in asymptomatic participants<sup>121,122</sup>. In addition, one study found a faster clearance of viral RNA in asymptomatic than in symptomatic individuals<sup>123</sup>, and another recorded a longer median duration of viral RNA shedding among asymptomatic patients<sup>119</sup>.

There are limited data regarding the presence of infectious virus in asymptomatic patients. One study showed lower virus isolation success from asymptomatic patients<sup>124</sup>, but only a small number of patients were included. Therefore, more studies evaluating infectious virus in asymptomatic patients would help to elucidate the differences in their infectivity compared with symptomatic patients.

### SARS-CoV-2 transmission

Viral loads have a key role in the SARS-CoV-2 transmission. As previously discussed, host (role of vaccination or previous infection) and viral factors (SARS-CoV-2 variants) greatly influence viral load dynamics and therefore further influence viral transmission.

## Influence of viral load on transmission

SARS-CoV-2 can be transmitted via larger droplets and aerosols produced when breathing, speaking, sneezing or coughing and to a lesser extent also by contaminated surfaces. As an infection can only be induced by infectious viral particles and not by remnant RNA or protein alone, the presence of infectious SARS-CoV-2 is required for secondary transmission. Although transmission is a multifactorial process that is also influenced, for example, by environmental and behavioural factors (such as humidity, air quality, exposure time or closeness of contact), the viral load of SARS-CoV-2 in the URT is considered to be a proxy for transmission risk.

An epidemiological study that included viral load analysis found that viral load of an index case strongly correlates with onward transmission, with higher viral loads for ancestral SARS-CoV-2 presenting a greater secondary attack rate risk<sup>125</sup>. In this study, viral load was identified as the main driver of transmission, with a more pronounced effect in household settings than in non-household settings (hospitals and nursing homes, among others). Transmission probability peaks around symptom onset, when infectious virus titres are estimated to be the highest during the course of infection. As viral load decreases with time, the probability of transmission also gradually declines in cases of infection with ancestral SARS-CoV-2 (ref.<sup>126</sup>). On this note, a study of health-care workers infected with ancestral virus documented no transmission from index cases later than 6 dpos, which is in line with findings showing reduced virus isolation success towards the end of week 1 of symptomatic disease<sup>127</sup>.

However, there are limitations when using viral load of an index case as a proxy for transmission. To date, the infectious dose of SARS-CoV-2 required to lead to a secondary transmission is not yet known, and the association between presence of infectious virus in the respiratory tract and infectiousness of the same individuals is poorly understood. In the only available human challenge trial that was conducted with ancestral SARS-CoV-2, an initial infectious dose of 10 TCID<sub>50</sub> did not lead to an infection in 16 of 36 participants<sup>29</sup>. Other factors, such as symptoms, type of contact, protective measures, vaccination status and other host factors may have an additionally strong effect on transmission<sup>128–133</sup>.

Viral load can markedly vary between individuals (as a result of individual susceptibility and of immunity from previous infections or vaccination), which leads to differences in their propensity to transmit the virus. Indeed, differences have been observed in the duration of infectious virus detection and in nasal and oral viral loads for both ancestral SARS-CoV-2 and Alpha<sup>33</sup>. Inter-individual variability was suggested to have a role in the observed heterogeneity of viral load dynamics, as some early immune signatures were significantly associated with higher oropharyngeal RNA viral loads in patients<sup>134</sup>. Therefore, observed heterogeneity between individuals has an important role in ongoing viral transmission<sup>33</sup>.

Such differences can lead to heterogeneity in virus transmission. Modelling with ancestral SARS-CoV-2 and Alpha estimated that individuals who are highly infectious, known as superspreaders, shed 57-fold more virus over the course of infection than those with lowest infectiousness<sup>33</sup>. By contrast, most patients with COVID-19 do not infect other individuals as they expel few to no viral particles from their airways<sup>135</sup>. Indeed, only a minority (about 8%) of patients positive for SARS-CoV-2 infected with ancestral SARS-CoV-2 or Alpha have significantly higher infectious virus titres than the rest of the population (as shown in a study measuring virus isolation probability in a large cohort of patients)<sup>86</sup>. Moreover, only 15%<sup>114</sup> to 19%<sup>136</sup> of individuals that

were infected led to 80% of secondary transmissions of ancestral SARS-CoV-2. Similar trends were confirmed for Omicron BA.1 and BA.2, for which only 9%<sup>137</sup> to 20%<sup>138</sup> of the infectious contacts were responsible for 80% of all transmissions.


Superspreading events are therefore characterized by infectious individuals having close contact with a high number of susceptible individuals and by a higher probability of transmission per contact. Aside from biological factors influencing these events, sociobehavioural and environmental factors contribute to the likelihood of superspreading (for example, large indoor gatherings with poor ventilation and no other infection prevention measures). Moreover, particular locations can represent a higher risk of transmission (for example, many superspreading events take place in crowded indoor settings, such as cruise ships, family gatherings, parties, elderly care centres and hospitals)<sup>139</sup>.

## The role of pre-existing immunity on viral shedding and transmission

All currently licensed SARS-CoV-2 vaccines are administered intramuscularly, leading to a rise in serum antibodies and protection from severe disease and death due to COVID-19, but not to long-term protection from infection<sup>140–142</sup>. The levels of circulating antibodies generated following vaccination decline over time, but can be elevated by a booster dose<sup>143,144</sup>. Furthermore, currently available vaccines were developed against the ancestral SARS-CoV-2 strain using the spike protein of the first sequenced virus, and the degree of protection from severe disease against other genetic variants was shown to vary<sup>145</sup>. Moreover, vaccination leads to limited induction of neutralizing antibodies on mucosal surfaces, which may have a role in mitigating virus replication and prevention of more pronounced disease<sup>146,147</sup>. For instance, secretory component antibodies, which are specific to mucosal surfaces, were detected in the saliva in 58% of participants 2 weeks post-vaccination with mRNA vaccines in one study, but the levels were significantly lower than in convalescent participants, and their neutralizing capacity significantly decayed 6 months post-vaccination<sup>148</sup>. A study on a small group of individuals uninfected or infected with Delta demonstrated that mucosal antibody responses induced by vaccination were low or undetectable, but breakthrough infections led to substantial increases of antibody titres in saliva<sup>149</sup>. However, the role of pre-existing mucosal immunity on infectious virus shedding and the possible correlation between the mucosal antibodies and viral loads in humans has not been elucidated.

As a result of waning antibodies and the emergence of VOCs with immune-evading properties, breakthrough infections have been increasingly reported among vaccinated individuals, mainly since the emergence of the Delta and Omicron VOCs. It has been debated whether vaccination with current SARS-CoV-2 vaccines impacts viral load (and therefore shedding) in breakthrough infections. The effect of vaccination on viral load and shedding is therefore of interest as it would mean that vaccination not only protects the vaccinee but can also help to mitigate virus spread by reducing infectious virus titres or shortening infectious shedding periods, thus having an impact beyond protection of the individual.

Overall, vaccination has been found to lead to reduced viral load (Fig. 4), although this decreases with time. Vaccination with ChAdOx1 vaccine (the Oxford–AstraZeneca vaccine) or BNT162b2 (the Pfizer/BioNTech vaccine) leads to lower RNA viral loads in individuals infected with Alpha, but the effect was weaker for breakthrough infections with Delta<sup>150,151</sup>. Immunization with BNT162b2 led to reduced RNA viral loads in Delta breakthrough infections, although this effect declined



**Fig. 4 | Influence of vaccination on viral load.** Similar RNA viral loads were detected in vaccinated and unvaccinated patients infected with the Delta variant of concern during the first 5 days post-onset of symptoms. However, faster clearance of viral RNA was shown in vaccinated patients. Infectious viral loads (IVLs) were significantly lower in vaccinated individuals and declined

faster than in unvaccinated individuals infected with Delta. Dynamics of viral loads in vaccinated individuals may vary widely in case of infection with another variant. Details of the underlying studies used to generate Fig. 4 can be found in Supplementary Table 3.

2 months after vaccination and ultimately faded 6 months after vaccination<sup>152</sup>. Immunization with ChAdOx1 vaccine also led to a reduction of RNA viral load in breakthrough infections with Alpha VOC<sup>153</sup>. Faster clearance of RNA viral loads was detected in the group of vaccinated patients who mostly received mRNA vaccines<sup>154,155</sup>, and lower probability of isolation of infectious virus from patients vaccinated with mRNA or adenoviral vector vaccines was observed<sup>156,157</sup>. Even though not all studies could demonstrate a reduction of RNA viral loads in Delta breakthrough infections<sup>150,154</sup>, infectious virus titres were reported to be lower in individuals vaccinated with mRNA or adenoviral vector vaccines despite similar levels of viral RNA<sup>25,93,157</sup>. Vaccination was also found to influence infectious virus isolation. Viable virus in cell culture was detected for significantly longer median time periods in unvaccinated patients infected with Delta than in vaccinated patients infected with Delta<sup>155,158</sup>. However, no significant differences in RNA viral loads were found between unvaccinated, fully vaccinated or boosted patients infected with Omicron BA.1 or BA.2 (refs. <sup>93,159</sup>), whereas infectious virus titres, measured quantitatively at 5 dpos, were lower in Omicron BA.1 breakthrough infections only after a booster dose<sup>25</sup>. Other studies showed that vaccination status did not influence infectious virus isolation success<sup>93</sup> or the time from initial positive PCR assay to culture conversion in patients infected with Omicron BA.1 (ref. <sup>85</sup>). These studies indicate that triple vaccination reduces infectious viral load but not the time period during which infectious virus can be isolated from Omicron breakthrough infections.

There are limited data on the effect of previous infection on viral shedding. A study performed on ancestral SARS-CoV-2 demonstrated lower RNA viral loads among seropositive individuals than among seronegative individuals<sup>160</sup>. Although higher levels of reinfection with Omicron BA.1 were demonstrated among unvaccinated patients previously infected with other SARS-CoV-2 variants<sup>161</sup>, there are no relevant data on the effect of previous infections on viral load dynamics.

Together, these findings suggest that vaccinated individuals are less infectious than unvaccinated individuals, although the duration of this effect has not been studied systematically. Nevertheless, there are some conflicting data on the effect of vaccination on onward transmission. An epidemiological study performed in the UK found that, despite RNA viral load declining faster among fully vaccinated than unvaccinated patients infected with Delta, the peak RNA viral loads were similar, and the secondary attack rate among household contacts exposed to fully vaccinated or unvaccinated index cases did not differ<sup>151</sup>. By contrast, data from Israel showed that less Delta transmission took place in households with vaccinated participants than with unvaccinated participants<sup>130</sup>. Another study from the UK showed that both BNT162b2 and ChAdOx1 vaccines led to the reduction of onward transmission from vaccinated index patients, although a stronger reduction was detected for Alpha than for Delta<sup>129</sup>, probably owing to the higher viral loads in the case of infection with Delta, as shown previously<sup>88,89,129</sup>. Finally, another study found that vaccination was associated with reduced onward transmission of Delta breakthrough infection due to shorter duration of viable virus shedding<sup>158</sup>.

Overall, even though the currently used vaccines are still based on the ancestral virus spike protein and elicit mainly a systemic rather than a mucosal immune response, some effect on viral load, infectious virus shedding and transmission has been observed<sup>129,130,162</sup>. Furthermore, with increasing rates of breakthrough infections in the Omicron waves since the end of 2021, many individuals display hybrid immunity consisting of vaccination combined with one or more natural infections before or after vaccination<sup>163,164</sup>. It is thought that such hybrid immunity may provide better control of virus replication in the mucosa<sup>149,163,165</sup>.

With the constant emergence of novel variants that can evade existing immunity, our understanding of the effect of vaccination on viral shedding should be constantly updated<sup>166</sup>. Better understanding of the role of mucosal immunity, and potentially vaccines that elicit

## Glossary

### Chimeric antigen receptor (CAR) T cell therapy

A way to treat cancer by using T cells expressing genetically engineered receptors to target cancer cells.

### Cycle threshold (Ct) value

The number of amplifications required for a target gene to cross the threshold determined by real-time PCR. Arbitrary test-specific Ct values inversely correlate with viral load.

### Focus-forming assays

Assays that count the number of 'foci', defined as a cluster of adjacent cells expressing viral antigen stained by a specific antibody.

### Immunostaining

A method for the detection of specific proteins in individual cells or tissues using antibodies. In the case of SARS-CoV-2, anti-nucleocapsid antibodies are used to detect virus in infected cells.

### Index case

The infected individual who is triggering an outbreak or a cluster by transmitting an infectious agent to others. There might be multiple index cases in an outbreak or epidemiological study.

local rather than systemic immune responses, are needed to aim for viral load reduction as a means to control SARS-CoV-2 circulation<sup>167–169</sup>.

### Influence of SARS-CoV-2 VOCs on transmission

There are several possible underlying causes of increased transmissibility of newly emerging variants, which allow VOCs to quickly outcompete previously circulating strains, including increased viral loads, a lower infectious dose required to establish infection and prolonged period of infectiousness<sup>170</sup>. Furthermore, the immune-evading properties of new variants lead to higher susceptibility of infection for vaccinated and previously infected individuals and result in higher transmissibility, as was observed with Omicron<sup>166,171</sup>.

The rapid emergence of SARS-CoV-2 variants with altered biological properties has shown that knowledge on viral loads, viral kinetics and infectious virus shedding is variant specific, and each emerging variant requires a reassessment. Although understanding of mutational profiles and associated phenotypes of SARS-CoV-2 variants has improved, reasons for enhanced transmissibility are manifold and not all understood yet. To date, shedding characteristics and transmission properties cannot be easily predicted based on sequences. Unlike immune-evasion mechanisms, shedding dynamics, such as kinetics of infectious virus titres or incubation periods of the SARS-CoV-2

variants, cannot be predicted from specific mutation patterns. With a still highly dynamic situation in terms of viral evolution of SARS-CoV-2, understanding viral kinetics and their effect on transmission remains of high public health interest.

### SARS-CoV-2 diagnostics in public health

Our ability to define the presence of infectious virus is key to guiding public health measures, as it will enable the isolation of infectious individuals to limit secondary transmission. Unfortunately, no point-of-care diagnostic test currently exists to determine infectious SARS-CoV-2 in a patient sample<sup>172</sup>, and virus culture as described above is not suited for diagnostic purposes. Thus, a range of approaches have been suggested to find a proxy for infectiousness to guide isolation periods.

One example is the detection of sgRNA transcripts, which are generated during virus replication, and specifically the synthesis of negative-strand RNA. Although sgRNAs are transcribed in infected cells, they are not packaged in the virions and can therefore serve as an indicator of active replication and thus of infectious virus. Specific RT-PCR assays were developed to detect sgRNAs in addition to the diagnostic detection of genomic SARS-CoV-2 RNA, but such assays have not made their way into routine diagnostic use owing to their lower sensitivity than conventional RT-PCR assays. Some studies found that detection of sgRNA correlates with detection of infectious virus<sup>4,173,174</sup>, and that sgRNA was rarely detectable 8 dpos<sup>67</sup>. However, sgRNA was detected in diagnostic samples up to 17 days after initial detection of infection<sup>175</sup> or in culture-negative samples<sup>176</sup>, probably owing to the stability and nuclease resistance of double-membrane vesicles containing sgRNAs. Thus, although the absence of sgRNA would indicate absence of viral replication, the presence of sgRNA does not necessarily indicate infectiousness<sup>19</sup>.

Ct values have also been used as a proxy for infectiousness, as described above. However, as already discussed, low-quality specimens resulting from technical mistakes during the collection process can falsely indicate an absence of infectious virus. Furthermore, owing to the quick increase of RNA viral load at the beginning of the infection, a low viral load, especially in the absence of symptoms or in the early symptomatic period, does not preclude that an individual will not soon enter the infectious period with the highest transmission risk. At such a period, viral loads reach their peak levels, causing the majority of transmission events<sup>59,126</sup>.

Even though Ag-RDTs are less sensitive than RT-PCR, they are less expensive, can be performed outside of laboratory settings and give faster results, and so are useful tools to guide isolation and limit transmission<sup>177</sup>. RT-PCR tests have a limit of detection of  $10^2$ – $10^3$  genome copies per millilitre, whereas Ag-RDTs have a limit of detection corresponding to  $10^4$ – $10^6$  genome copies per millilitre<sup>177–180</sup>. Infectious individuals typically have RNA viral loads of  $>10^6$  genome copies per millilitre, which corresponds largely with a Ct of 25 in most RT-PCR assays<sup>4</sup>, indicating that Ag-RDT is a good proxy for infectiousness<sup>177</sup>. However, the obvious limitations of Ag-RDT, such as lower sensitivity of infectious virus detection towards the end of infection<sup>47,52</sup>, should not be neglected. Ag-RDTs have also shown variation in their sensitivity and specificity for detection of SARS-CoV-2 VOCs<sup>53,54</sup>, which is a challenge as new variants emerge.

Overall, all of the currently available diagnostic methods have certain limitations for detection of infectious virus. However, even if these tests serve only as imperfect tools when used as proxies for infectiousness, their implementation as part of a public health strategy is not intended to prevent every single infection, but rather to reduce the

number of infectious people in the community and thus to decrease the number of secondary transmissions.

## Conclusions

Entering the third year of the pandemic, much knowledge on SARS-CoV-2 viral loads, infectious virus shedding and windows of infectiousness has been gained, although emerging SARS-CoV-2 variants and an increasing population immunity add more complexity to the situation.

Although much progress has been made during the pandemic in the field of diagnostics, to date, no diagnostic tests exist that reliably determine the presence of infectious virus. Continuing evaluation of viral-shedding characteristics under these changing circumstances and understanding the biological properties of novel SARS-CoV-2 variants when it comes to viral shedding remain of importance to guide public health practices.

Published online: 2 December 2022

## References

- Puelles, V. G. et al. Multiorgan and renal tropism of SARS-CoV-2. *N. Engl. J. Med.* **383**, 590–592 (2020).
- Lamers, M. M. & Haagmans, B. L. SARS-CoV-2 pathogenesis. *Nat. Rev. Microbiol.* **20**, 270–284 (2022).
- Peng, L. et al. SARS-CoV-2 can be detected in urine, blood, anal swabs, and oropharyngeal swabs specimens. *J. Med. Virol.* **92**, 1676–1680 (2020).
- Wölfel, R. et al. Virological assessment of hospitalized patients with COVID-2019. *Nature* **581**, 465–469 (2020).
- Zhang, W. et al. Molecular and serological investigation of 2019-nCoV infected patients: implication of multiple shedding routes. *Emerg. Microbes Infect.* **9**, 386–389 (2020).
- Pérez-Bartolomé, F. & Sánchez-Quiros, J. Ocular manifestations of SARS-CoV-2: literature review. *Arch. Soc. Esp. Ophthalmol.* **96**, 32–40 (2021).
- Vetter, P. et al. Daily viral kinetics and innate and adaptive immune response assessment in COVID-19: a case series. *mSphere* <https://doi.org/10.1128/mSphere.00827-20> (2020).
- Jeong, H. W. et al. Viable SARS-CoV-2 in various specimens from COVID-19 patients. *Clin. Microbiol. Infect.* **26**, 1520–1524 (2020).
- Cerrada-Romero, C. et al. Excretion and viability of SARS-CoV-2 in feces and its association with the clinical outcome of COVID-19. *Sci. Rep.* **12**, 7397 (2022).
- Dergham, J. & Delerue, J. Isolation of viable SARS-CoV-2 virus from feces of an immunocompromised patient suggesting a possible fecal mode of transmission. *J. Clin. Med.* <https://doi.org/10.3390/jcm10122696> (2021).
- Xiao, F. et al. Infectious SARS-CoV-2 in feces of patient with severe COVID-19. *Emerg. Infect. Dis.* **26**, 1920–1922 (2020).
- Sun, J. et al. Isolation of infectious SARS-CoV-2 from urine of a COVID-19 patient. *Emerg. Microbes Infect.* **9**, 991–993 (2020).
- Colavita, F. et al. SARS-CoV-2 isolation from ocular secretions of a patient with COVID-19 in Italy with prolonged viral RNA detection. *Ann. Intern. Med.* **173**, 242–243 (2020).
- Matsuyama, S. et al. Enhanced isolation of SARS-CoV-2 by TMPRSS2-expressing cells. *Proc. Natl Acad. Sci. USA* **117**, 7001–7003 (2020).
- Case, J. B., Bailey, A. L., Kim, A. S., Chen, R. E. & Diamond, M. S. Growth, detection, quantification, and inactivation of SARS-CoV-2. *Virology* **548**, 39–48 (2020).
- Baggen, J., Vanstreels, E., Jansen, S. & Daelemans, D. Cellular host factors for SARS-CoV-2 infection. *Nat. Microbiol.* **6**, 1219–1232 (2021).
- Chu, H. et al. Comparative tropism, replication kinetics, and cell damage profiling of SARS-CoV-2 and SARS-CoV with implications for clinical manifestations, transmissibility, and laboratory studies of COVID-19: an observational study. *Lancet Microbe* **1**, e14–e23 (2020).
- van Kampen, J. J. A. et al. Duration and key determinants of infectious virus shedding in hospitalized patients with coronavirus disease-2019 (COVID-19). *Nat. Commun.* **12**, 267 (2021).
- Bruce, E. A. et al. Predicting infectivity: comparing four PCR-based assays to detect culturable SARS-CoV-2 in clinical samples. *EMBO Mol. Med.* **14**, e15290 (2022).
- Essaidi-Laziosi, M. & Perez Rodriguez, F. J. Estimating clinical SARS-CoV-2 infectiousness in Vero E6 and primary airway epithelial cells. *Lancet Microbe* **2**, e571 (2021).
- Liu, R. et al. Positive rate of RT-PCR detection of SARS-CoV-2 infection in 4880 cases from one hospital in Wuhan, China, from Jan to Feb 2020. *Clin. Chim. Acta* **505**, 172–175 (2020).
- Corman, V. M. et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Eur. Surveill.* <https://doi.org/10.2807/1560-7917.ES.2020.25.3.2000045> (2020).
- Bullard, J. et al. Predicting infectious severe acute respiratory syndrome coronavirus 2 from diagnostic samples. *Clin. Infect. Dis.* **71**, 2663–2666 (2020).
- Jefferson, T., Spencer, E. A., Brassey, J. & Heneghan, C. Viral cultures for coronavirus disease 2019 infectivity assessment: a systematic review. *Clin. Infect. Dis.* **73**, e3884–e3899 (2021).
- Puhach, O. et al. Infectious viral load in unvaccinated and vaccinated individuals infected with ancestral, Delta or Omicron SARS-CoV-2. *Nat. Med.* **28**, 1491–1500 (2022).
- van Kasteren, P. B. et al. Comparison of seven commercial RT-PCR diagnostic kits for COVID-19. *J. Clin. Virol.* **128**, 104412 (2020).
- Bentley, E. et al. Collaborative study for the establishment of a WHO international standard for SARS-CoV-2 RNA (WHO, 2020).
- Zou, L. et al. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *N. Engl. J. Med.* **382**, 1177–1179 (2020).
- Killingley, B. et al. Safety, tolerability and viral kinetics during SARS-CoV-2 human challenge in young adults. *Nat. Med.* **28**, 1031–1041 (2022).
- Yu, F. et al. Quantitative detection and viral load analysis of SARS-CoV-2 in infected patients. *Clin. Infect. Dis.* **71**, 793–798 (2020).
- European Centre for Disease Prevention and Control & World Health Organization Regional Office for Europe. Methods for the detection and identification of SARS-CoV-2 variants: second update, August 2022 (WHO, 2022).
- Ke, R. et al. Longitudinal analysis of SARS-CoV-2 vaccine breakthrough infections reveals limited infectious virus shedding and restricted tissue distribution. *Open Forum Infect. Dis.* **9**, ofac192 (2022).
- Ke, R. et al. Daily longitudinal sampling of SARS-CoV-2 infection reveals substantial heterogeneity in infectiousness. *Nat. Microbiol.* **7**, 640–652 (2022).
- Pekosz, A. et al. Antigen-based testing but not real-time polymerase chain reaction correlates with severe acute respiratory syndrome coronavirus 2 viral culture. *Clin. Infect. Dis.* **73**, e2861–e2866 (2021).
- Monel, B. et al. Release of infectious virus and cytokines in nasopharyngeal swabs from individuals infected with non-Alpha or Alpha SARS-CoV-2 variants: an observational retrospective study. *EBioMedicine* **73**, 103637 (2021).
- Kirby, J. E. et al. SARS-CoV-2 antigen tests predict infectivity based on viral culture: comparison of antigen, PCR viral load, and viral culture testing on a large sample cohort. *Clin. Microbiol. Infect.* <https://doi.org/10.1016/j.cmi.2022.07.010> (2022).
- Pickering, S. et al. Comparative performance of SARS-CoV-2 lateral flow antigen tests and association with detection of infectious virus in clinical specimens: a single-centre laboratory evaluation study. *Lancet Microbe* **2**, e461–e471 (2021).
- Tariq, M. et al. Viable severe acute respiratory syndrome coronavirus 2 isolates exhibit higher correlation with rapid antigen assays than subgenomic RNA or genomic RNA. *Front. Microbiol.* **12**, 718497 (2021).
- Chu, V. T. et al. Comparison of home antigen testing with RT-PCR and viral culture during the course of SARS-CoV-2 infection. *JAMA Intern. Med.* **182**, 701–709 (2022).
- Albert, E. et al. Field evaluation of a rapid antigen test (Panbio™ COVID-19 Ag rapid test device) for COVID-19 diagnosis in primary healthcare centres. *Clin. Microbiol. Infect.* **27**, 472.e7–472.e10 (2021).
- Ford, L. et al. Epidemiologic characteristics associated with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) antigen-based test results, real-time reverse transcription polymerase chain reaction (rRT-PCR) cycle threshold values, subgenomic RNA, and viral culture results from university testing. *Clin. Infect. Dis.* **73**, e1348–e1355 (2021).
- Berger, A. et al. Diagnostic accuracy of two commercial SARS-CoV-2 antigen-detecting rapid tests at the point of care in community-based testing centers. *PLoS ONE* **16**, e0248921 (2021).
- Brümmer, L. E. et al. Accuracy of novel antigen rapid diagnostics for SARS-CoV-2: a living systematic review and meta-analysis. *PLoS Med.* **18**, e1003735 (2021).
- Ngo Nsoga, M. T. et al. Diagnostic accuracy of Panbio rapid antigen tests on oropharyngeal swabs for detection of SARS-CoV-2. *PLoS ONE* **16**, e0253321 (2021).
- Korenkov, M. et al. Evaluation of a rapid antigen test to detect SARS-CoV-2 infection and identify potentially infectious individuals. *J. Clin. Microbiol.* **59**, e0089621 (2021).
- Yamayoshi, S. et al. Comparison of rapid antigen tests for COVID-19. *Viruses* **12**, 1420 (2020).
- McKay, S. L. et al. Performance evaluation of serial SARS-CoV-2 rapid antigen testing during a nursing home outbreak. *Ann. Intern. Med.* **174**, 945–951 (2021).
- Nordgren, J. et al. SARS-CoV-2 rapid antigen test: high sensitivity to detect infectious virus. *J. Clin. Virol.* **140**, 104846 (2021).
- Fernandez-Montero, A. & Argemi, J. Validation of a rapid antigen test as a screening tool for SARS-CoV-2 infection in asymptomatic populations. Sensitivity, specificity and predictive values. *EClinicalMedicine* **37**, 100954 (2021).
- Currie, D. W. et al. Relationship of SARS-CoV-2 antigen and reverse transcription PCR positivity for viral cultures. *Emerg. Infect. Dis.* **28**, 717–720 (2022).
- Corman, V. M. et al. Comparison of seven commercial SARS-CoV-2 rapid point-of-care antigen tests: a single-centre laboratory evaluation study. *Lancet Microbe* **2**, e311–e319 (2021).
- Lopera, T. J. & Alzate-Ángel, J. C. The usefulness of antigen testing in predicting contagiousness in COVID-19. *Microbiol. Spectr.* **10**, e0196221 (2022).
- Osterman, A. et al. Impaired detection of Omicron by SARS-CoV-2 rapid antigen tests. *Med. Microbiol. Immunol.* **211**, 105–117 (2022).
- Galliez, R. M. et al. Evaluation of the Panbio COVID-19 antigen rapid diagnostic test in subjects infected with Omicron using different specimens. *Microbiol. Spectr.* **10**, e0125022 (2022).
- Raich-Regué, D. et al. Performance of SARS-CoV-2 antigen-detecting rapid diagnostic tests for Omicron and other variants of concern. *Front. Microbiol.* **13**, 810576 (2022).
- Van Ogtrop, M. L., van de Laar, T. J. W., Eggink, D., Vanhommerig, J. W. & van der Reijden, W. A. Comparison of the performance of the Panbio COVID-19 antigen test in SARS-CoV-2 B.1.1.7 (Alpha) variants versus non-B.1.1.7 variants. *Microbiol. Spectr.* **9**, e0088421 (2021).

57. Lindner, A. K. et al. Head-to-head comparison of SARS-CoV-2 antigen-detecting rapid test with self-collected nasal swab versus professional-collected nasopharyngeal swab. *Eur. Respir. J.* **57**, 2003961 (2021).
58. Meiners, L. & Horn, J. SARS-CoV-2 rapid antigen test sensitivity and viral load in freshly symptomatic hospital employees, December 2020 to February 2022. Preprint at <https://doi.org/10.2139/ssrn.4099425> (2022).
59. He, X. et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat. Med.* **26**, 672–675 (2020).
60. Néant, N. et al. Modeling SARS-CoV-2 viral kinetics and association with mortality in hospitalized patients from the French COVID cohort. *Proc. Natl Acad. Sci. USA* **118**, e2017962118 (2021).
61. Owusu, D. et al. Persistent SARS-CoV-2 RNA shedding without evidence of infectiousness: a cohort study of individuals with COVID-19. *J. Infect. Dis.* **224**, 1362–1371 (2021).
62. Cevik, M. et al. SARS-CoV-2, SARS-CoV, and MERS-CoV viral load dynamics, duration of viral shedding, and infectiousness: a systematic review and meta-analysis. *Lancet Microbe* **2**, e13–e22 (2021).
63. Chen, X. et al. Associations of clinical characteristics and treatment regimens with the duration of viral RNA shedding in patients with COVID-19. *Int. J. Infect. Dis.* **98**, 252–260 (2020).
64. Kim, S. M. & Hwang, Y. J. Prolonged SARS-CoV-2 detection and reversed RT-PCR results in mild or asymptomatic patients. *Infect. Dis.* **53**, 31–37 (2021).
65. Talmy, T. & Tsur, A. Duration of SARS-CoV-2 detection in Israel Defense Forces soldiers with mild COVID-19. *J. Med. Virol.* **93**, 608–610 (2021).
66. Singanayagam, A. et al. Duration of infectiousness and correlation with RT-PCR cycle threshold values in cases of COVID-19, England, January to May 2020. *Euro Surveill.* <https://doi.org/10.2807/1560-7917.ES.2020.25.32.2001483> (2020).
67. Perera, R. et al. SARS-CoV-2 virus culture and subgenomic RNA for respiratory specimens from patients with mild coronavirus disease. *Emerg. Infect. Dis.* **26**, 2701–2704 (2020).
68. Killerby, M. E. et al. Shedding of culturable virus, seroconversion, and 6-month follow-up antibody responses in the first 14 confirmed cases of coronavirus disease 2019 in the United States. *J. Infect. Dis.* **224**, 771–776 (2021).
69. Glans, H. et al. Shedding of infectious SARS-CoV-2 by hospitalized COVID-19 patients in relation to serum antibody responses. *BMC Infect. Dis.* **21**, 494 (2021).
70. Badu, K. et al. SARS-CoV-2 viral shedding and transmission dynamics: implications of WHO COVID-19 discharge guidelines. *Front. Med.* **8**, 648660 (2021).
71. Munker, D. et al. Dynamics of SARS-CoV-2 shedding in the respiratory tract depends on the severity of disease in COVID-19 patients. *Eur. Respir. J.* **58**, 2002724 (2021).
72. Folgueira, M. D. & Luczkowiak, J. Prolonged SARS-CoV-2 cell culture replication in respiratory samples from patients with severe COVID-19. *Clin. Microbiol. Infect.* **27**, 886–891 (2021).
73. Kim, M. C. et al. Duration of culturable SARS-CoV-2 in hospitalized patients with Covid-19. *N. Engl. J. Med.* **384**, 671–673 (2021).
74. Chen, P. Z. et al. SARS-CoV-2 shedding dynamics across the respiratory tract, sex, and disease severity for adult and pediatric COVID-19. *eLife* <https://doi.org/10.7554/eLife> (2021).
75. Cunha, M. D. P. et al. Atypical prolonged viral shedding with intra-host SARS-CoV-2 evolution in a mildly affected symptomatic patient. *Front. Med.* **8**, 760170 (2021).
76. Aydinli, T. et al. Shedding of viable SARS-CoV-2 after immunosuppressive therapy for cancer. *N. Engl. J. Med.* **383**, 2586–2588 (2020).
77. Caillaud, S., Benotmane, I., Gautier Vargas, G., Perrin, P. & Fafi-Kremer, S. SARS-CoV-2 viral dynamics in immunocompromised patients. *Am. J. Transpl.* **21**, 1667–1669 (2021).
78. Roedel, K. et al. Viral dynamics of SARS-CoV-2 in critically ill allogeneic hematopoietic stem cell transplant recipients and immunocompetent patients with COVID-19. *Am. J. Respir. Crit. Care Med.* **203**, 242–245 (2021).
79. Leung, W. F. et al. COVID-19 in an immunocompromised host: persistent shedding of viable SARS-CoV-2 and emergence of multiple mutations: a case report. *Int. J. Infect. Dis.* **114**, 178–182 (2022).
80. European Centre for Disease Prevention and Control. Rapid risk assessment: assessing SARS-CoV-2 circulation, variants of concern, non-pharmaceutical interventions and vaccine rollout in the EU/EEA, 15th update (European Centre for Disease Prevention and Control, 2021).
81. Grant, R. et al. Impact of SARS-CoV-2 Delta variant on incubation, transmission settings and vaccine effectiveness: results from a nationwide case–control study in France. *Lancet Reg. Health Eur.* **13**, 100278 (2022).
82. Ogata, T. & Tanaka, H. Shorter incubation period among unvaccinated Delta variant coronavirus disease 2019 patients in Japan. *Int. J. Environ. Res. Public Health* <https://doi.org/10.3390/ijerph19031127> (2022).
83. Backer, J. A. et al. Shorter serial intervals in SARS-CoV-2 cases with Omicron BA.1 variant compared with Delta variant, the Netherlands, 13 to 26 December 2021. *Eur. Surveill.* <https://doi.org/10.2807/1560-7917.ES.2022.27.6.2200042> (2022).
84. Takahashi, K. et al. Duration of infectious virus shedding by SARS-CoV-2 Omicron variant-infected vaccinees. *Emerg. Infect. Dis.* **28**, 998–1001 (2022).
85. Boucau, J. et al. Duration of shedding of culturable virus in SARS-CoV-2 Omicron (BA.1) infection. *N. Engl. J. Med.* **387**, 275–277 (2022).
86. Jones, T. C. et al. Estimating infectiousness throughout SARS-CoV-2 infection course. *Science* <https://doi.org/10.1126/science.aba5273> (2021).
87. Li, B. et al. Viral infection and transmission in a large, well-traced outbreak caused by the SARS-CoV-2 Delta variant. *Nat. Commun.* **13**, 460 (2022).
88. Bolze, A. et al. SARS-CoV-2 variant Delta rapidly displaced variant Alpha in the United States and led to higher viral loads. *Cell Rep. Med.* **3**, 100564 (2022).
89. Earnest, R. et al. Comparative transmissibility of SARS-CoV-2 variants Delta and Alpha in New England, USA. *Cell Rep. Med.* **3**, 100583 (2022).
90. Luo, C. H. et al. Infection with the SARS-CoV-2 Delta variant is associated with higher recovery of infectious virus compared to the Alpha variant in both unvaccinated and vaccinated individuals. *Clin. Infect. Dis.* **75**, e715–e725 (2021).
91. Despres, H. W. et al. Measuring infectious SARS-CoV-2 in clinical samples reveals a higher viral titer: RNA ratio for Delta and Epsilon vs. Alpha variants. *Proc. Natl Acad. Sci. USA* <https://doi.org/10.1073/pnas.2116518119> (2022).
92. Hay, J. A. et al. Quantifying the impact of immune history and variant on SARS-CoV-2 viral kinetics and infection rebound: a retrospective cohort study. *eLife* **11**, e81849 (2022).
93. Fall, A. et al. The displacement of the SARS-CoV-2 variant Delta with Omicron: an investigation of hospital admissions and upper respiratory viral loads. *EbioMedicine* <https://doi.org/10.1016/j.ebiom.2022.104008> (2022).
94. Lentini, A. & Pereira, A. Monitoring of the SARS-CoV-2 Omicron BA.1/BA.2 lineage transition in the Swedish population reveals increased viral RNA levels in BA.2 cases. *Med* **3**, 636–664 (2022).
95. Qassim, S. H. et al. Effects of BA.1/BA.2 subvariant, vaccination, and prior infection on infectiousness of SARS-CoV-2 Omicron infections. *J. Travel Med.* **29**, taac068 (2022).
96. Marking, U. et al. Correlates of protection, viral load trajectories and symptoms in BA.1, BA.1.1 and BA.2 breakthrough infections in triple vaccinated healthcare workers. Preprint at *medRxiv* <https://doi.org/10.1101/2022.04.02.22273333> (2022).
97. Wang, Y. et al. Transmission, viral kinetics and clinical characteristics of the emergent SARS-CoV-2 Delta VOC in Guangzhou, China. *EClinicalMedicine* **40**, 101129 (2021).
98. Siedner, M. J. et al. Duration of viral shedding and culture positivity with postvaccination SARS-CoV-2 Delta variant infections. *JCI Insight* <https://doi.org/10.1172/jci.insight.155483> (2022).
99. Pierce, C. A. et al. Immune responses to SARS-CoV-2 infection in hospitalized pediatric and adult patients. *Sci. Transl. Med.* **12**, eabd5487 (2020).
100. Takahashi, T. et al. Sex differences in immune responses that underlie COVID-19 disease outcomes. *Nature* **588**, 315–320 (2020).
101. Bunyavanich, S., Do, A. & Vicencio, A. Nasal gene expression of angiotensin-converting enzyme 2 in children and adults. *JAMA* **323**, 2427–2429 (2020).
102. Zheng, S. et al. Viral load dynamics and disease severity in patients infected with SARS-CoV-2 in Zhejiang province, China, January–March 2020: retrospective cohort study. *BMJ* **369**, m1443 (2020).
103. L'Huillier, A. G. & Torriani, G. Culture-competent SARS-CoV-2 in nasopharynx of symptomatic neonates, children, and adolescents. *Emerg. Infect. Dis.* **26**, 2494–2497 (2020).
104. Baggio, S. et al. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) viral load in the upper respiratory tract of children and adults with early acute coronavirus disease 2019 (COVID-19). *Clin. Infect. Dis.* **73**, 148–150 (2021).
105. Chung, E. et al. Comparison of symptoms and RNA levels in children and adults with SARS-CoV-2 infection in the community setting. *JAMA Pediatr.* **175**, e212025 (2021).
106. Han, M. S. et al. Clinical characteristics and viral RNA detection in children with coronavirus disease 2019 in the Republic of Korea. *JAMA Pediatr.* **175**, 73–80 (2021).
107. Bellon, M. et al. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) viral load kinetics in symptomatic children, adolescents, and adults. *Clin. Infect. Dis.* **73**, e1384–e1386 (2021).
108. Madera, S. et al. Nasopharyngeal SARS-CoV-2 viral loads in young children do not differ significantly from those in older children and adults. *Sci. Rep.* **11**, 3044 (2021).
109. Lauer, S. A. et al. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Ann. Intern. Med.* **172**, 577–582 (2020).
110. Elias, C., Sekri, A., Leblanc, P., Cucherat, M. & Vanhems, P. The incubation period of COVID-19: a meta-analysis. *Int. J. Infect. Dis.* **104**, 708–710 (2021).
111. Dhoubi, W. et al. The incubation period during the pandemic of COVID-19: a systematic review and meta-analysis. *Syst. Rev.* **10**, 101 (2021).
112. Brandal, L. T. et al. Outbreak caused by the SARS-CoV-2 Omicron variant in Norway, November to December 2021. *Eur. Surveill.* **26**, 2101147 (2021).
113. Wu, Y. et al. Incubation period of COVID-19 caused by unique SARS-CoV-2 strains: a systematic review and meta-analysis. *JAMA Netw. Open* **5**, e2228008 (2022).
114. Sun, K. et al. Transmission heterogeneities, kinetics, and controllability of SARS-CoV-2. *Science* **371**, eabe2424 (2021).
115. Lavezzo, E. et al. Suppression of a SARS-CoV-2 outbreak in the Italian municipality of Vo'. *Nature* **584**, 425–429 (2020).
116. Gudbjartsson, D. F. et al. Spread of SARS-CoV-2 in the Icelandic population. *N. Engl. J. Med.* **382**, 2302–2315 (2020).
117. Glnet, M. et al. Asymptomatic COVID-19 adult outpatients identified as significant viable SARS-CoV-2 shedders. *Sci. Rep.* **11**, 20615 (2021).
118. Wang, Y. et al. Characterization of an asymptomatic cohort of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infected individuals outside of Wuhan, China. *Clin. Infect. Dis.* **71**, 2132–2138 (2020).
119. Long, Q. X. et al. Clinical and immunological assessment of asymptomatic SARS-CoV-2 infections. *Nat. Med.* **26**, 1200–1204 (2020).
120. Lee, S. et al. Clinical course and molecular viral shedding among asymptomatic and symptomatic patients with SARS-CoV-2 infection in a community treatment center in the Republic of Korea. *JAMA Intern. Med.* **180**, 1447–1452 (2020).

121. Zhou, R. et al. Viral dynamics in asymptomatic patients with COVID-19. *Int. J. Infect. Dis.* **96**, 288–290 (2020).
122. Hall, S. M. et al. Comparison of anterior nares CT values in asymptomatic and symptomatic individuals diagnosed with SARS-CoV-2 in a university screening program. *PLoS ONE* **17**, e0270694 (2022).
123. Kissler, S. M. et al. Viral dynamics of acute SARS-CoV-2 infection and applications to diagnostic and public health strategies. *PLoS Biol.* **19**, e3001333 (2021).
124. Tallmadge, R. L. et al. Viral RNA load and infectivity of SARS-CoV-2 in paired respiratory and oral specimens from symptomatic, asymptomatic, or postsymptomatic individuals. *Microbiol. Spectr.* **10**, e0226421 (2022).
125. Marks, M. et al. Transmission of COVID-19 in 282 clusters in Catalonia, Spain: a cohort study. *Lancet Infect. Dis.* **21**, 629–636 (2021).
126. Marc, A. et al. Quantifying the relationship between SARS-CoV-2 viral load and infectiousness. *eLife* <https://doi.org/10.7554/eLife.69302> (2021).
127. Cheng, H. Y. et al. Contact tracing assessment of COVID-19 transmission dynamics in Taiwan and risk at different exposure periods before and after symptom onset. *JAMA Intern. Med.* **180**, 1156–1163 (2020).
128. Muggleston, M. A. et al. Presymptomatic, asymptomatic and post-symptomatic transmission of SARS-CoV-2: joint British Infection Association (BIA), Healthcare Infection Society (HIS), Infection Prevention Society (IPS) and Royal College of Pathologists (RCPath) guidance. *BMC Infect. Dis.* **22**, 453 (2022).
129. Eyre, D. W. et al. Effect of Covid-19 vaccination on transmission of Alpha and Delta variants. *N. Engl. J. Med.* **386**, 744–756 (2022).
130. Prunas, O. et al. Vaccination with BNT162b2 reduces transmission of SARS-CoV-2 to household contacts in Israel. *Science* **375**, 1151–1154 (2022).
131. Lyngse, F. P. et al. Household transmission of SARS-CoV-2 Omicron variant of concern subvariants BA.1 and BA.2 in Denmark. *Nat. Commun.* **13**, 5760 (2022).
132. Leung, N. H. L. Transmissibility and transmission of respiratory viruses. *Nat. Rev. Microbiol.* **19**, 528–545 (2021).
133. Alihsan, B. et al. The efficacy of facemasks in the prevention of COVID-19: a systematic review. Preprint at *medRxiv* <https://doi.org/10.1101/2022.07.28.22278153> (2022).
134. Hu, Z. et al. Early immune markers of clinical, virological, and immunological outcomes in patients with COVID-19: a multi-omics study. *Elife* **11**, e77943 (2022).
135. Chen, P. Z. et al. Heterogeneity in transmissibility and shedding SARS-CoV-2 via droplets and aerosols. *eLife* <https://doi.org/10.7554/eLife.65774> (2021).
136. Adam, D. C. et al. Clustering and superspreading potential of SARS-CoV-2 infections in Hong Kong. *Nat. Med.* **26**, 1714–1719 (2020).
137. Guo, Z. et al. Superspreading potential of infection seeded by the SARS-CoV-2 Omicron BA.1 variant in South Korea. *J. Infect.* **85**, e77–e79 (2022).
138. Guo, Z. et al. Superspreading potential of COVID-19 outbreak seeded by Omicron variants of SARS-CoV-2 in Hong Kong. *J. Travel Med.* **29**, taac049 (2022).
139. Althouse, B. M. et al. Superspreading events in the transmission dynamics of SARS-CoV-2: opportunities for interventions and control. *PLoS Biol.* **18**, e3000897 (2020).
140. Baden, L. R. et al. Efficacy and safety of the mRNA-1273 SARS-CoV-2 vaccine. *N. Engl. J. Med.* **384**, 403–416 (2021).
141. Lopez Bernal, J. et al. Effectiveness of Covid-19 vaccines against the B.1.617.2 (Delta) variant. *N. Engl. J. Med.* **385**, 585–594 (2021).
142. Feikin, D. R. et al. Duration of effectiveness of vaccines against SARS-CoV-2 infection and COVID-19 disease: results of a systematic review and meta-regression. *Lancet* **399**, 924–944 (2022).
143. Belik, M. et al. Comparative analysis of COVID-19 vaccine responses and third booster dose-induced neutralizing antibodies against Delta and Omicron variants. *Nat. Commun.* **13**, 2476 (2022).
144. Pegu, A. et al. Durability of mRNA-1273 vaccine-induced antibodies against SARS-CoV-2 variants. *Science* **373**, 1372–1377 (2021).
145. Lipsitch, M., Krammer, F., Regev-Yochay, G., Lustig, Y. & Balicer, R. D. SARS-CoV-2 breakthrough infections in vaccinated individuals: measurement, causes and impact. *Nat. Rev. Immunol.* **22**, 57–65 (2022).
146. Mostaghimi, D., Valdez, C. N., Larson, H. T., Kalinich, C. C. & Iwasaki, A. Prevention of host-to-host transmission by SARS-CoV-2 vaccines. *Lancet Infect. Dis.* [https://doi.org/10.1016/S1473-3099\(21\)00472-2](https://doi.org/10.1016/S1473-3099(21)00472-2) (2021).
147. Russell, M. W., Moldoveanu, Z., Ogra, P. L. & Mestecky, J. Mucosal immunity in COVID-19: a neglected but critical aspect of SARS-CoV-2 infection. *Front. Immunol.* **11**, 611337 (2020).
148. Sheikh-Mohamed, S. et al. Systemic and mucosal IgA responses are variably induced in response to SARS-CoV-2 mRNA vaccination and are associated with protection against subsequent infection. *Mucosal Immunol.* **15**, 799–808 (2022).
149. Collier, A. Y. et al. Characterization of immune responses in fully vaccinated individuals after breakthrough infection with the SARS-CoV-2 Delta variant. *Sci. Transl. Med.* **14**, eabn6150 (2022).
150. Pouwels, K. B. et al. Effect of Delta variant on viral burden and vaccine effectiveness against new SARS-CoV-2 infections in the UK. *Nat. Med.* **27**, 2127–2135 (2021).
151. Singanayagam, A. et al. Community transmission and viral load kinetics of the SARS-CoV-2 Delta (B.1.617.2) variant in vaccinated and unvaccinated individuals in the UK: a prospective, longitudinal, cohort study. *Lancet Infect. Dis.* **22**, 183–195 (2021).
152. Levine-Tiefenbrun, M. et al. Viral loads of Delta-variant SARS-CoV-2 breakthrough infections after vaccination and booster with BNT162b2. *Nat. Med.* **27**, 2108–2110 (2021).
153. Emary, K. R. W. et al. Efficacy of ChAdOx1nCoV-19 (AZD1222) vaccine against SARS-CoV-2 variant of concern 202012/01 (B.1.17): an exploratory analysis of a randomised controlled trial. *Lancet* **397**, 1351–1362 (2021).
154. Chia, P. Y. et al. Virological and serological kinetics of SARS-CoV-2 Delta variant vaccine breakthrough infections: a multicentre cohort study. *Clin. Microbiol. Infect.* **28**, 612.e1–612.e7 (2021).
155. Garcia-Knight, M. et al. Infectious viral shedding of SARS-CoV-2 Delta following vaccination: a longitudinal cohort study. *PLoS Pathog.* **18**, e1010802 (2022).
156. Peña-Hernández, M. A. et al. Comparison of infectious SARS-CoV-2 from the nasopharynx of vaccinated and unvaccinated individuals. Preprint at *medRxiv* <https://doi.org/10.1101/2021.12.28.21268460> (2022).
157. Shamier, M. C. et al. Virological characteristics of SARS-CoV-2 vaccine breakthrough infections in health care workers. Preprint at *medRxiv* <https://doi.org/10.1101/2021.08.20.21262158> (2021).
158. Jung, J. et al. Transmission and infectious SARS-CoV-2 shedding kinetics in vaccinated and unvaccinated individuals. *JAMA Netw. Open* **5**, e2213606 (2022).
159. Hirotsu, Y. et al. Similar viral loads in Omicron infections regardless of vaccination status. Preprint at *medRxiv* <https://doi.org/10.1101/2022.04.19.22274005> (2022).
160. Letizia, A. G. et al. SARS-CoV-2 seropositivity and subsequent infection risk in healthy young adults: a prospective cohort study. *Lancet Respir. Med.* **9**, 712–720 (2021).
161. Pulliam, J. R. C. et al. Increased risk of SARS-CoV-2 reinfection associated with emergence of Omicron in South Africa. *Science* **376**, eabn4947 (2022).
162. Harris, R. J. et al. Effect of vaccination on household transmission of SARS-CoV-2 in England. *N. Engl. J. Med.* **385**, 759–760 (2021).
163. Bates, T. A. et al. Vaccination before or after SARS-CoV-2 infection leads to robust humoral response and antibodies that effectively neutralize variants. *Sci. Immunol.* **7**, eabn8014 (2022).
164. Wrátil, P. R. et al. Three exposures to the spike protein of SARS-CoV-2 by either infection or vaccination elicit superior neutralizing immunity to all variants of concern. *Nat. Med.* **28**, 496–503 (2022).
165. Malato, J. et al. Risk of BA.5 infection among persons exposed to previous SARS-CoV-2 variants. *N. Engl. J. Med.* **387**, 953–954 (2022).
166. Cao, Y. et al. BA.2.12.1, BA.4 and BA.5 escape antibodies elicited by Omicron infection. *Nature* **608**, 593–602 (2022).
167. Hassan, A. O. et al. A single-dose intranasal ChAd vaccine protects upper and lower respiratory tracts against SARS-CoV-2. *Cell* **183**, 169–184 (2020).
168. Lapuente, D. et al. Protective mucosal immunity against SARS-CoV-2 after heterologous systemic prime-mucosal boost immunization. *Nat. Commun.* **12**, 6871 (2021).
169. Afkhami, S. et al. Respiratory mucosal delivery of next-generation COVID-19 vaccine provides robust protection against both ancestral and variant strains of SARS-CoV-2. *Cell* **185**, 896–915 (2022).
170. Grubaugh, N. D. & Hodcroft, E. B. Public health actions to control new SARS-CoV-2 variants. *Cell* **184**, 1127–1132 (2021).
171. Carreño, J. M. et al. Activity of convalescent and vaccine serum against SARS-CoV-2 Omicron. *Nature* **602**, 682–688 (2022).
172. Ashcroft, P., Lehtinen, S. & Bonhoeffer, S. Test-trace-isolate-quarantine (TTIQ) intervention strategies after symptomatic COVID-19 case identification. *PLoS ONE* **17**, e0263597 (2022).
173. Santos Bravo, M. et al. Viral culture confirmed SARS-CoV-2 subgenomic RNA value as a good surrogate marker of infectivity. *J. Clin. Microbiol.* **60**, e0160921 (2022).
174. Kim, J. Y. et al. Diagnostic usefulness of subgenomic RNA detection of viable SARS-CoV-2 in patients with COVID-19. *Clin. Microbiol. Infect.* **28**, 101–106 (2022).
175. Alexandersen, S. & Chamings, A. SARS-CoV-2 genomic and subgenomic RNAs in diagnostic samples are not an indicator of active replication. *Nat. Commun.* **11**, 6059 (2020).
176. Bonenfant, G. et al. Surveillance and correlation of SARS-CoV-2 viral RNA, antigen, virus isolation, and self-reported symptoms in a longitudinal study with daily sampling. *Clin. Infect. Dis.* <https://doi.org/10.1093/cid/ciac282> (2022).
177. Peeling, R. W., Heymann, D. L., Teo, Y. Y. & Garcia, P. J. Diagnostics for COVID-19: moving from pandemic response to control. *Lancet* **399**, 757–768 (2022).
178. Pilarowski, G. et al. Performance characteristics of a rapid severe acute respiratory syndrome coronavirus 2 antigen detection assay at a public plaza testing site in San Francisco. *J. Infect. Dis.* **223**, 1139–1144 (2021).
179. Leber, W. et al. Comparing the diagnostic accuracy of point-of-care lateral flow antigen testing for SARS-CoV-2 with RT-PCR in primary care (REAP-2). *EClinicalMedicine* **38**, 101011 (2021).
180. Kohmer, N. et al. The comparative clinical performance of four SARS-CoV-2 rapid antigen tests and their correlation to infectivity in vitro. *J. Clin. Med.* <https://doi.org/10.3390/jcm10020328> (2021).
181. Wang, W. et al. Detection of SARS-CoV-2 in different types of clinical specimens. *JAMA* **323**, 1843–1844 (2020).
182. Lee, R. A., Herigon, J. C., Benedetti, A., Pollock, N. R. & Denkinger, C. M. Performance of saliva, oropharyngeal swabs, and nasal swabs for SARS-CoV-2 molecular detection: a systematic review and meta-analysis. *J. Clin. Microbiol.* **59**, e02881-20 (2021).
183. Tsang, N. N. Y. et al. Diagnostic performance of different sampling approaches for SARS-CoV-2 RT-PCR testing: a systematic review and meta-analysis. *Lancet Infect. Dis.* **21**, 1233–1245 (2021).
184. Peto, T. COVID-19: rapid antigen detection for SARS-CoV-2 by lateral flow assay: a national systematic evaluation of sensitivity and specificity for mass-testing. *EClinicalMedicine* **36**, 100924 (2021).
185. Hay, J. A. et al. Estimating epidemiologic dynamics from cross-sectional viral load distributions. *Science* **373**, eabh0635 (2021).

186. Peccia, J. et al. Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. *Nat. Biotechnol.* **38**, 1164–1167 (2020).
187. Xin, H. et al. The incubation period distribution of coronavirus disease 2019: a systematic review and meta-analysis. *Clin. Infect. Dis.* **73**, 2344–2352 (2021).
188. Yu, I. T. et al. Evidence of airborne transmission of the severe acute respiratory syndrome virus. *N. Engl. J. Med.* **350**, 1731–1739 (2004).
189. Cheng, P. K. et al. Viral shedding patterns of coronavirus in patients with probable severe acute respiratory syndrome. *Lancet* **363**, 1699–1700 (2004).
190. Peiris, J. S. et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. *Lancet* **361**, 1767–1772 (2003).
191. Chan, P. K. S. et al. Laboratory diagnosis of SARS. *Emerg. Infect. Dis.* **10**, 825–831 (2004).
192. Liu, W. et al. Long-term SARS coronavirus excretion from patient cohort, China. *Emerg. Infect. Dis.* **10**, 1841–1843 (2004).
193. Xu, D. et al. Persistent shedding of viable SARS-CoV in urine and stool of SARS patients during the convalescent phase. *Eur. J. Clin. Microbiol. Infect. Dis.* **24**, 165–171 (2005).
194. V'kovski, P. et al. Disparate temperature-dependent virus–host dynamics for SARS-CoV-2 and SARS-CoV in the human respiratory epithelium. *PLoS Biol.* **19**, e3001158 (2021).
195. Pitzer, V. E., Leung, G. M. & Lipsitch, M. Estimating variability in the transmission of severe acute respiratory syndrome to household contacts in Hong Kong, China. *Am. J. Epidemiol.* **166**, 355–363 (2007).
196. Riley, S. et al. Transmission dynamics of the etiological agent of SARS in Hong Kong: impact of public health interventions. *Science* **300**, 1961–1966 (2003).
197. Zaki, A. M., van Boheemen, S., Bestebroer, T. M., Osterhaus, A. D. & Fouchier, R. A. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N. Engl. J. Med.* **367**, 1814–1820 (2012).
198. Breban, R., Riou, J. & Fontanet, A. Interhuman transmissibility of Middle East respiratory syndrome coronavirus: estimation of pandemic risk. *Lancet* **382**, 694–699 (2013).
199. Oh, M. D. et al. Viral load kinetics of MERS coronavirus infection. *N. Engl. J. Med.* **375**, 1303–1305 (2016).
200. Corman, V. M. et al. Viral shedding and antibody response in 37 patients with Middle East respiratory syndrome coronavirus infection. *Clin. Infect. Dis.* **62**, 477–483 (2016).
201. Min, C.-K. et al. Comparative and kinetic analysis of viral shedding and immunological responses in MERS patients representing a broad spectrum of disease severity. *Sci. Rep.* **6**, 25359 (2016).
202. Ip, D. K. et al. Viral shedding and transmission potential of asymptomatic and paucisymptomatic influenza virus infections in the community. *Clin. Infect. Dis.* **64**, 736–742 (2017).
203. Carrat, F. et al. Time lines of infection and disease in human influenza: a review of volunteer challenge studies. *Am. J. Epidemiol.* **167**, 775–785 (2008).
204. Pawelek, K. A. et al. Modeling within-host dynamics of influenza virus infection including immune responses. *PLoS Comput. Biol.* **8**, e1002588 (2012).
205. Ip, D. K. M. et al. The dynamic relationship between clinical symptomatology and viral shedding in naturally acquired seasonal and pandemic influenza virus infections. *Clin. Infect. Dis.* **62**, 431–437 (2016).
206. Nair, H. et al. Global burden of acute lower respiratory infections due to respiratory syncytial virus in young children: a systematic review and meta-analysis. *Lancet* **375**, 1545–1555 (2010).
207. Bagga, B. et al. Comparing influenza and RSV viral and disease dynamics in experimentally infected adults predicts clinical effectiveness of RSV antivirals. *Antivir. Ther.* **18**, 785–791 (2013).
208. DeVincenzo, J. P. et al. Viral load drives disease in humans experimentally infected with respiratory syncytial virus. *Am. J. Respir. Crit. Care Med.* **182**, 1305–1314 (2010).
209. Kutter, J. S. et al. Small quantities of respiratory syncytial virus RNA only in large droplets around infants hospitalized with acute respiratory infections. *Antimicrob. Resist. Infect. Control* **10**, 100 (2021).

## Acknowledgements

The authors thank E. Boehm for help with literature search and proofreading. The work was funded by the COVID-19 National Research Program (grant number 198412) of the Swiss National Science Foundation.

## Author contributions

O.P. and I.E. wrote the manuscript. B.M. created the figures. All authors contributed to the discussion of the content, and reviewed and edited the manuscript before submission.

## Competing interests

The authors declare no competing interests.

## Additional information

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s41579-022-00822-w>.

**Correspondence** should be addressed to Isabella Ecklerle.

**Peer review information** *Nature Reviews Microbiology* thanks Nancy Leung, Quanyi Wang and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© Springer Nature Limited 2022