

Archive ouverte UNIGE

https://archive-ouverte.unige.ch

Article scientifique

Article 2022

Published version

Open Access

This is the published version of the publication, made available in accordance with the publisher's policy.

How is physicians' implicit prejudice against the obese and mentally ill moderated by specialty and experience?

Fitzgerald, Chloé Nathalie Sarah; Mumenthaler, Christian; Berner, Delphine; Schindler, Mélinee; Brosch, Tobias; Hurst, Samia

How to cite

FITZGERALD, Chloé Nathalie Sarah et al. How is physicians" implicit prejudice against the obese and mentally ill moderated by specialty and experience? In: BMC medical ethics, 2022, vol. 23, n° 1, p. 86. doi: 10.1186/s12910-022-00815-7

This publication URL:https://archive-ouverte.unige.ch/unige:166137Publication DOI:10.1186/s12910-022-00815-7

© The author(s). This work is licensed under a Creative Commons Attribution (CC BY) <u>https://creativecommons.org/licenses/by/4.0</u>

RESEARCH

Open Access

How is physicians' implicit prejudice against the obese and mentally ill moderated by specialty and experience?

Chloë FitzGerald^{1*}, Christian Mumenthaler^{2,3}, Delphine Berner¹, Mélinée Schindler¹, Tobias Brosch^{4,5} and Samia Hurst^{1*}

Abstract

Background: Implicit prejudice can lead to disparities in treatment. The effects of specialty and experience on implicit obesity and mental illness prejudice had not been explored. The main objective was to examine how specializing in psychiatry/general medicine and years of experience moderated implicit obesity and mental illness prejudice among Swiss physicians. Secondary outcomes included examining the malleability of implicit bias via two video interventions and a condition of cognitive load, correlations of implicit bias with responses to a clinical vignette, and correlations with explicit prejudice.

Methods: In stage 1, participants completed an online questionnaire including a clinical vignette. In stage 2, implicit prejudice pre- and post- intervention was tested using a 4×4 between-subject design including a control group. In stage 3, explicit prejudice was tested with feeling thermometers and participants were debriefed. Participants were 133 psychiatrists and internists working in Geneva, hospital-based and private practice. Implicit prejudice was assessed using a Weight IAT (Implicit Association Test) and a Mental Illness IAT. Explicit feelings towards the obese and the mentally ill were measured using Feeling Thermometers. A clinical vignette assessed the level of concern felt for a fictional patient under four conditions: control, obese, depression, obese and depression. Linear regression was conducted to test for association of gender, experience, and specialty with responses to vignettes, pre-intervention IATs and explicit attitudes, and to test for association of interventions (or control) with post-intervention IATs and explicit attitudes. Reported effect sizes were computed using Cohen's d. Two-tailed p < 0.05 was selected as the significance threshold.

Results: Compared to internists, psychiatrists showed significantly less implicit bias against mentally vs. physically ill people than internists and warmer explicit feelings towards the mentally ill. More experienced physicians displayed warmer explicit feelings towards the mentally ill and a greater level of concern for the fictional patients in the vignette than the less experienced, except when the patient was described as obese.

Conclusions: Specialty moderates both implicit and explicit mental illness prejudice. Experience moderates explicit mental illness bias and concern for patients. The effect of specialty on implicit prejudice seems to be based principally on self-selection.

*Correspondence: cnsfitzgerald@gmail.com; samia.hurst@unige.ch

¹ iEH2 (Institute for Ethics, History and the Humanities), Faculty of Medicine, University of Geneva, Geneva, Switzerland Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Keywords: Implicit bias, Prejudice, Stereotype, Specialization, Experience, Training, Medical education, Doctor– patient relationships, Vulnerable populations, Mental health, Obesity

Background

Equal treatment of patients regardless of skin colour, weight, or other characteristics is part of the standard of care in medicine and physicians often cite helping others as a reason for practising medicine [1-6]. This standard of care has become even more relevant globally in the recent pandemic, where resources have often been scarce, and where disparities in death rates from Covid-19 correlated with race have been documented in the UK and the US [7, 8]. Fears among patients about discrimination due to factors such as obesity have also increased [9].

Implicit biases against stigmatised groups represent a threat to this standard of care. Implicit biases are associations, which have been characterised as automatic, uncontrollable, unconscious or arational [10], between a category attribute, e.g. being overweight, and a negative evaluation (implicit prejudice), or another category attribute, e.g. being lazy (implicit stereotype). In contrast to explicit biases, evaluations that a person makes consciously, implicit biases are typically manifest in nonverbal behaviour, such as the frequency of eye contact and physical proximity [11]. Their most disturbing aspect is the potential dissociation between what a physician explicitly intends (e.g. treat everyone equally) and the hidden influence of implicit associations on her decisionmaking and action (e.g. perceiving a black child's pain as less severe than a white child's and thus deciding not to prescribe the patient a pain medication) [12].

The effects of medical specialty and experience on implicit obesity and mental illness prejudice have not been explored in the literature. The principal objective of the study was to examine how specializing in psychiatry/general medicine and years of experience moderated implicit obesity and mental illness prejudice among Swiss physicians. The relationship between levels of implicit bias and clinical behaviour remains unclear [14] and we therefore evaluated physicians' intended behaviour through their responses to a clinical vignette. Secondary outcomes included examining the malleability of implicit bias via two video interventions, a condition of cognitive load and correlations with explicit prejudice.

Prejudicial and stereotypical implicit attitudes are widespread among the worldwide population and can influence behaviour outside the laboratory [14, 15]. Many implicit biases are present even among the stigmatized outgroup, e.g. most obese people display an implicit prejudice against the obese [16]. Implicit biases among health care professionals and the potential influence on clinical care is a concern [17–19]. While they are not the whole picture, implicit biases are likely to play a part in health care disparities. In the US, racial health care disparities are widely documented and implicit racial bias is a possible contributing factor [20]. In the UK, a recent study found that a woman is five times more likely to die in childbirth or post-partum if she is black than if she is white [21]. Other health care disparities related to factors such as socio-economic status, sexual orientation and gender are possibly partly due to implicit biases [22].

Implicit bias training is widely on offer in the Englishspeaking world, despite a limited evidence base for effective interventions [23]. But what is the role of training and experience in medicine itself? Do more experienced physicians exhibit more or less implicit bias than trainees? Medical training might reasonably be expected to reduce bias with its emphasis on care and the impartial standard. One study found that mental health training correlated with more positive implicit and explicit attitudes to the mentally ill [24]. Experience in the field could potentially reduce prejudice by making physicians more sympathetic to patients. As they gain experience and knowledge, medical professionals may experience less cognitive strain and thus have more mental resources to devote to ensuring equal treatment of patients. It is possible that exposure to counterstereotipic exemplars that contradict an implicit stereotype can lessen it [25].

On the other hand, some experience will confirm stereotypes rather than contradict them, particularly given confirmation bias [26] and the important role of medical superiors, who may be biased, in training. Current evidence shows that contact with the stigmatised group sometimes decreases prejudice, but can also increase it, depending on the nature of the contact [27-30]. In addition, medical training encourages fast-thinking and shortcuts, which can increase incidence of implicit bias [31]. With experience and training a physician may become more confident about their ability to be impartial and thus more susceptible to bias. There is evidence that if we are confident about our moral capabilities we are more likely to display implicit biases because we register the goal of being impartial as being 'achieved' and monitor ourselves less [32]. There has also been research showing that clinical exposure can reduce empathy [33, 34].

In addition to training and experience, medical specialty may influence levels of implicit bias. One study found that paediatricians displayed less implicit race bias than other physicians [35]. The aforementioned study showing an effect of mental health training on bias towards the mentally ill could be a sign that choice of specialty and the subsequent training and environment influences bias. Specialty, training and experience are likely to interact in complex ways to produce potential effects on implicit bias. No studies have specifically explored the interaction of specialty and training/experience with implicit biases. This study aimed to fill this gap.

Implicit obesity prejudice and implicit mental illness prejudice were chosen partly because they are both tied to characteristics of patients that can be relevant for medical reasons. The former has been found to be present in health care professionals to a similar extent as in the general population [36]; there is less data on implicit mental health bias, but evidence indicates its presence in both physicians and the general population [24, 37-39]. The evidence indicating that mental health training reduced bias made this bias an obvious choice to explore. While black/white race bias has received the most attention due to its relevance in the US, it is not necessarily the most relevant to Swiss physicians, who may discriminate more based on country of origin rather than skin colour [40]. In addition, Geneva city and hospital are highly multicultural, meaning both patients and physicians in the hospital come from a multitude of different countries and typically speak several languages. Unlike race bias, obesity and mental illness bias have been found to exist as explicit biases in addition to implicit among health care professionals [13, 38, 39]. One explanation is that because obesity and mental health are medically relevant characteristics they fail to flag warning signs for prejudice with physicians, as, typically, race would.

Given findings of the presence of implicit obesity bias and implicit mental illness bias among physicians from many other countries [13], some level of implicit bias was expected to be found among Swiss physicians. On the other hand, culturally specific contexts could mean that within Swiss physicians, or specifically physicians in Geneva, we would find higher or lower levels than had been found in other studies [41].

It was hypothesized that training would have an effect on implicit biases, while the direction that this effect would take was uncertain given the mixed evidence to date. It was hypothesized that specializing in psychiatry would correlate with lower mental illness bias based on the one study showing an influence of mental health training on bias [35].

Methods

Study participants

Participants were recruited from psychiatry and general internal medicine. As mentioned in the Introduction,

Page 3 of 11

the evidence so far on the effects of experience and contact on prejudice is mixed. The specialties of psychiatry and general medicine were chosen because in examining bias against illness of the mind and a characteristic of the physical body (obesity) it was deemed interesting to compare specialists in the mind (psychiatrists) and those who did not specialise in the mind and thus could be expected to focus more on physical characteristics (general internists). Female physicians have been found to display weaker implicit bias against the obese than male physicians, hence gender differences in obesity bias were expected [36]. Participants were recruited from the Geneva University Hospitals and from private practices in Geneva by email or physical mail.

Data collection

All participants provided written informed consent. The experiment was described as an investigation into implicit attitudes using a categorisation task with the aim of improving standards of clinical care. The IAT was not named nor were the words 'bias' and 'prejudice' mentioned to avoid influence on responses to the clinical vignette. The order of measures (Fig. 1) was chosen to mask the characteristics under study for as long as possible. The ethics committee approved this procedure for obtaining consent. Participants first completed a demographic questionnaire and a clinical vignette online. Patient characteristics were randomized in the vignette to create four groups each receiving a different version.

A face-to-face meeting was then held with each participant, who completed a Mental Illness IAT and a Weight IAT, followed by either one of two interventions, instructions for the cognitive load condition, or a control condition, then a repeat of both IATs, and finally each responded to two Feeling Thermometers to rate their feelings towards the obese and the mentally ill. The interviewer asked participants to refrain from discussing the contents of the study with colleagues. Interviews were conducted in the local language of the participants, French, and all materials and tests were in French. All methods were carried out in accordance with relevant guidelines and regulations.

Measures

IATs

The most prevalent measure of implicit biases is the Implicit Association Test (IAT), a computerized task where participants rapidly categorize negatively and positively valenced words with images or words. The relative speed of association of, e.g. in a Race IAT, black faces with positively-valenced words (as compared to the other possible associations), indicates the level of bias [42].



Mental illness stimuli were taken from a previously tested IAT, comparing words for physical illnesses and words for mental illnesses paired with negatively and positively valenced words [39]. The Weight IAT used stimuli from the Project Implicit website: silhouette images of thin and obese people paired with negatively and positively valenced words [43]. Words were translated from English to French. D-score interpretation used by Project Implicit indicate a score greater than 0.15 as a slight bias, greater than 0.35 as a moderate bias, and greater than 0.65 as a strong bias. Negative scores represent the inverse association, namely an association between either mental health or obesity (as compared to physical health or thinness) with positive rather than negative words [42, 43].

Vignette

The vignette was taken from a study that found differences in clinical responses to pain correlated with patient gender [44]. It was translated from Portuguese to French and modified to create four versions: a control version with no medical history, a version where the BMI of the patient was 32 and thus indicated clinical obesity, a version were the patient had experienced depressive episodes, and a version were the patient had a BMI of 32 and had experienced depressive episodes. There were six questions that asked participants to evaluate pain intensity, clinical severity, clinical urgency and pain credibility on a scale of 1–7 (Additional file 1: S1 Appendix).

Feeling thermometer

These consisted in a continuous unnumbered line with 'warm feelings' written on the left side of the line and 'cold feelings' on the right. The range was 0-12, but the increments were not indicated on the actual line where participants marked a cross. Instead, the segments were designated as follows:

0–2.75 warm feelings 2.75–5.5 slightly warm feelings 5.5–6.5 neutral feelings 6.5–9.25 slightly cold feelings 9.25–12 cold feelings

Participants marked the point that represented their feelings towards the obese and the mentally ill. Lower scores represent warmer feelings. The feeling thermometer measures explicit feelings, thus is a measure of explicit bias.

Interventions and cognitive load

Few interventions had been tested specifically on implicit weight bias and implicit mental illness bias and even fewer have produced significant results [45–50]. One of the methods that had been found to potentially reduce

anti-age, racial and other implicit biases was perspective-taking, thought to increase empathy with the target group [51, 52]. We decided to test a version of perspective-taking, given that there was as much evidence that it worked as there was for other strategies and it was the one that most appealed to our research team. The participants were required to watch the short (1.25 min) extracts of videos made by the National Health Service in the UK before immediately retaking the IAT. The videos had been designed for training, to help physicians identify and empathise with their mentally ill and obese patients. They showed two genuine female patients in the UK talking about their clinical experiences with obesity and depression. The videos were subtitled in French and are available from the following website: https://www.unige.ch/medecine/ieh2/fr/recherche/ groupe-samia-hurst-manjo/.

Given concerns regarding the test-retest reliability of the IAT [41, 53] and the possibility of a learning effect when participants completed the second IAT, we included a control group. In the control group, participants counted backwards in twos out loud in place of the intervention for the same length of time and then retook the IATs.

A fourth group, in place of the intervention, was instructed on how to proceed with the second set of IATs under a cognitive load condition, consisting in counting backwards in twos out loud during the intervention. They then proceeded to retake the IATs. The effects of implicit bias are thought to increase under stress and time pressure, common working conditions for physicians that can be simulated—albeit approximately—with a cognitive load. While previous research has shown that conditions of cognitive load increase implicit biases [54, 55], we were not aware of other research that looked directly at the effects of cognitive load while performing an IAT.

It was hypothesized that the video interventions would correlate with reduced levels of implicit prejudice and that the cognitive load condition would correlate with increased levels of implicit prejudice when compared with the control group.

Statistical analysis

The sample size was computed using the software GPower and was based on an initial a priori power analysis for ANOVA, targeting a medium-sized interaction effect between two factors (specialty and experience) with two level each with an alpha of 0.05 and a power of 0.80, requiring a total sample size of N=158 (40 per cell). Data was entered into the SPSS statistical software package for analysis. Participants' socio-demographic characteristics were analysed using descriptive statistics. Inter item reliability (the questionnaire's internal

consistency) was tested using Cronbach's alpha. We changed our analysis to enable treatment of IAT scores as continuous variables in response to reviewer comments. Linear regression was conducted to test for association of gender, experience, and specialty with responses to vignettes, pre-intervention IATs and explicit attitudes. Linear regression was also conducted to test for association of interventions (or control) with post-intervention IATs and explicit attitudes. To take possible confounding variables for this second analysis into account we also included gender, experience, and specialty. Reported effect sizes were computed using Cohen's d. Two-tailed p < 0.05 was selected as the significance threshold. The data is publicly available at the following https://doi.org/10.26037/yareta:md2ryexqsrchhb2fafgor6lcmm.

Results

Participants

779 physicians were contacted via email (or physical mail where email addresses were unavailable), and followed up by a second email and a telephone call. The response rate was 24% for this first stage. 133 eligible physicians went on to complete both stages of the study (81.1% from the first stage). Difficulty recruiting less experienced physicians was encountered and so the category was extended by increasing the number of years of practice from < 5 to < 6. This enabled 5 participants who had initially been discarded to complete the second stage when recontacted. Their data are included in the second response rate. The desired number of participants was not achieved in the less-experienced psychiatrist category, with a final total of 13 participants. In the other categories the desired number of 40 participants was reached. Participant characteristics displayed a good range of age, years in practice, and site of practice and an equal gender distribution (Table 1). The 40 participants who only

Table 1 Participant characteristics

Age (years)	Mean (SD) Median Range	39 (11) 36 24–72
Gender	Male Female	49% 51%
Years in practice	Mean (SD) Median Range	14 (11) 10 1–47
	Less than 6 More than 8	40% 60%
Speciality	Internal medicine Psychiatry	60% 40%
Site of practice	Hospital Hospital ambulatory Private practice	59% 23% 18%

completed the first stage of the experiment were statistically compared to the participants who completed both stages and were found to be similar with regard to gender, experience, age, and specialty.

Overall implicit and explicit prejudice and level of concern among physicians

Despite incomplete comparability due to different stimuli, data do suggest greater implicit prejudice towards the obese than towards the mentally ill among physicians overall (D-score 0.53 and 0.09, respectively, Table 2).

The overall mean score for physicians would not count as a mental illness prejudice according to Project Implicit scoring conventions, while obesity prejudice would count as moderate [42, 43]. When a one value t-test was performed to compare the bias to 0, the p value for the Mental Illness IAT was 0.033 and for the Weight IAT was < 0.001.

Explicit feelings were significantly warmer towards the mentally ill than towards the obese (3.94 and 4.89, respectively, p < 0.001, Table 2).

Cronbach's alpha for the responses to the vignette was 0.79 thus we examined responses together rather than individually, as 'level of concern for the fictional patient'. There were no significant effects found.

No significant gender effects were found in the results.

Effect of specialty on prejudice

Compared to internists, psychiatrists appeared to show significantly less implicit bias against mentally ill vs physically ill people (D-score -0.23 and 0.31, respectively, p < 0.001). The overall mean score for psychiatrists showed a slight positive bias and the overall mean score for internists showed a slight negative bias according to Project Implicit scoring conventions [42, 43]. No significant effects were found for specialty on the Weight IAT (Table 2).

Psychiatrists displayed significantly warmer explicit feelings towards the mentally ill on the Feeling Thermometer than did internists (p < 0.001, Table 2).

Effect of experience on prejudice

Data did not show significant differences between more or less experienced physicians in either IAT (Table 2).

More experienced physicians showed significantly warmer feelings towards the mentally ill (p=0.012, Table 2).

The level of concern for the fictional patient was significantly greater among more experienced physicians (p=0.014, Table 2). Experience was significantly correlated with an increased level of concern when the fictional patient was described as depressed and when he was not described as obese (p=0.038, p=0.018, Table 2).

When the fictional patient was obese, experience did not correlate with an increased level of concern.

Effect of interventions on prejudice

Data showed no significant effects of the interventions or of the cognitive load on the IAT or Feeling Thermometer results (Table 3).

Discussion

The findings indicate that specialty correlates with both implicit and explicit mental illness prejudice, but not with obesity prejudice. Although experience did not correlate with implicit bias, it was associated with warmer explicit feelings towards the mentally ill and greater concern for the mentally ill -but not the obese- patient on a clinical vignette.

The physicians overall displayed what Project Implicit qualifies as a moderate implicit bias against the obese [42, 43], slightly weaker than the strong negative bias found among physicians and the general public in a previous study [36]. Previously existing evidence on bias against the mentally ill is less conclusive, partly because different IATs have been used [24, 37, 38]. Our Mental Illness IAT results consisting in a slight positive bias for psychiatrists and a slight negative bias for internists indicate that internists may have similar levels to the general population, whereas psychiatrists display less [39]. However, caution should be taken with the interpretation of these results given that the comparison category used in our IAT was physical illness. Another explanation of the difference between psychiatrists and internists is that, while they had similar levels of implicit prejudice towards the mentally ill, internists had a more positive bias towards physical illness than the psychiatrists.

Scores appear to show a greater implicit obesity prejudice than implicit mental illness prejudice, but image-IATs produce different effects from word-IATs due to the level of stimulus representation so cannot be directly compared [56]. However, this difference was reproduced on other measures: greater obesity prejudice than mental illness prejudice was found in the responses to the vignette and on the Feeling Thermometer. The lack of mental illness prejudice on the vignette could be due to our choice of mental disorder. Depression is not associated with such negative implicit stereotypes as other mental disorders, such as schizophrenia [47]. There may be more awareness among physicians of a risk of bias against the mentally ill than against the obese because of the wider prevalence of campaigns against mental illness stigma.

Compared to internists, psychiatrists showed significantly less implicit bias against mentally ill vs physically ill people. The lack of a significant correlation with

	٩
	Effect size (95% CI)
	Internists Mean (SD)
	Psychiatrists Mean (SD)
Table 2 Specialty, experience, and gender	All physicians

	All physicia	us		Psychiatrists	Internists	Effect size	٩	Less .	More .	Effect size (95% CI)	٩
	Mean (SD)	Effect size (95% CI)	٩	(Uc) Mean	(Uc) Mean	(17 %66)		experienced Mean (SD)	experienced Mean (SD)		
Mental Illness IAT (D score)	0.09 (0.50)			- 0.23 (0.42)	0.31 (0.44)	1.25 (0.87–1.63)	< 0.001	0.14 (0.50)	0.06 (0.51)	— 0.158 (— 0.506 to 0.19)	0.398
Weight IAT (D score)	0.53 (0.36)			0.50 (0.39)	0.54 (0.34)	0.11 (-0.236 to 0.458)	0.674	0.56 (0.33)	0.50 (0.38)	- 0.166 (- 0.514 to 0.181	0.392
Feeling Thermometer Mentally III (Score 0-12)*	3.96 (2.28)	0.409 (0.065–0.752)	< 0.001	2.69 (1.85)	4.80 (2.16)	1.033 (0.813–0.109)	< 0.001	4.58 (2.21)	3.55 (2.25)	— 0.461 (— 0.813 to 0.109)	0.012
Feeling Thermometer Obese (Score 0–12)*	4.89 (2.27)			5.11 (2.69)	4.75 (1.95)	— 0.158 (— 0.506 to 0.189)	0.374	4.85 (1.94)	4.92 (2.48)	0.031 (— 0.316 to 0.378)	0.876
Vignette overall (Total score 6–42)#	29.94 (3.34)			30.15 (3.17)	29.80 (3.47)	— 0.104 (— 0.452 to 0.243)	0.556	29.08 (3.33)	30.51 (3.24)	0.437 (0.085- 0.788)	0.014
Vignette/Depressive patient	29.98 (3.58)	- 0.021 (- 0.361 to 0.319)	0.914	30.86 (2.93)	29.44 (3.86)	— 0.403 (— 0.754 to 0.053)	0.144	28.47 (2.93)	30.61 (3.66)	0.617 (0.041–1.194)	0.038
Vignette/Non- depressive patient	29.91 (3.18)			29.65 (3.28)	30.09 (3.13)	0.138 (0.23 to 0.485)	0.544	29.36 (3.51)	30.41 (2.78)	0.34 (— 0.01 to 0.689)	0.154
Vignette/Obese patient	29.32 (3.34)	0.39 (0.047–0.733)	0.058	29.15 (2.86)	29.43 (3.59)	0.084 (— 0.263 to 0.432)	0.734	28.97 (3.19)	29.62 (3.41)	0.196 (— 0.152 to 0.543)	0.418
Vignette/Non-obese patient	30.61 (3.28)			31.19 (3.18)	30.21 (3.33)	— 0.3 (— 0.649 to 0.049)	0.244	29.24 (3.61)	31.28 (2.93)	0.65 (0.072–1.227)	0.018
P values in bold are stati	stically significa	ant									

Control (N $=$ 34)	Mental illness video (N $=$ 33)	Obesity video (N $=$ 33)
Mean	Mean (SD) Effect size (95% Cl) p	Mean (SD) Effect size (95% Cl) p

Table 3 Effects of interventions

	Control (N=34)	Mental illne	ss video (N=33)	U	Obesity vide	o (N=33)	Cognitive lo	ad (N=33)	
	Mean	Mean (SD)	Effect size (95% CI)	d	Mean (SD)	Effect size (95% Cl) p	Mean (SD)	Effect size (95% CI) p	
Mental illness IAT (D score)	0.043 (0.44)	0.031 (0.37)	- 0.029 (- 0.508 to 0.45)	0.898 (0.012 (0.47)	-0.068 (-0.547 to 0.411) .860	0.005 (0.38)	-0.092 (-0.572 to 0.387) 0.7	778
Weight IAT (D score)	0.382 (0.30)	0.386 (0.38)	0.012 (-0.467 to 0.491)	0.466 (0.372 (0.44)	-0.027 (-0.506 to 0.452) .632	0.240 (0.30)	0.352 (-0.131 to 0.834) 0.0	058
Feeling thermometer mentally III (Score 0–12)*	3.944 (2.10)	3.873 (2.69)	- 0.029 (- 0.508 to 0.45)	0.802	3.967 (2.22)	0.011 (0.468 to 0.49) .986	4.058 (2.18)	0.053 (0.426 to 0.532) 0.7	780
Feeling thermometer obese (Score 0–12)*	4.903 (1.98)	4.958 (2.32)	0.026 (— 0.454 to 0.505)	0.848	4.321 (2.53)	-0.257 (-0.738 to 0.224) .098	5.382 (2.22)	0.228 (— 0.253 to 0.798) 0.1	160

P values in bold are statistically significant

* Lower scores indicate warmer feelings

All interventions were compared to the control

experience suggests that it may be a self-selection effect (which could take place either before or soon after entry into specialization) rather than the result of training or experience. Perhaps those who choose to specialise in psychiatry already have less implicit bias against the mentally ill or have their initial levels of implicit bias lowered very soon after entering the speciality. If this is the case, this finding is of particular practical importance to the specialty of psychiatry. Alternative hypotheses could include that physicians have more positive associations with diseases they feel better equipped to treat. Greater familiarity, or salience, has also been proposed as a confounding factor in IAT testing [57] and could be at play here.

In contrast to other studies suggesting that training affects implicit prejudice, we found no association between experience and implicit bias [24]. However, the more experienced physicians displayed warmer explicit feelings towards the mentally ill and a higher level of concern for the fictional patient in the vignette. This could indicate that experience in the field improves patient care by making physicians more sympathetic to patients, in addition to other ways. This is particularly interesting in light of other studies suggesting that clinical exposure may erase empathy in medical students [33, 34]. When the fictional patient was described as obese, experience did not correlate with an increased level of concern. One explanation could be that experience helps physicians to prevent their implicit attitudes influencing their level of concern for patients, rather than reducing their implicit mental illness prejudice. This may not occur in the case of obese patients because of lack of awareness of the dangers of obesity prejudice.

Unlike other studies, no gender effects were found in our study. The interventions we tested had no significant effects and the cognitive load procedure made no difference to the IAT scores.

A limitation of the study is that it may include a higher percentage of physicians interested in improving patient care than average because those who participated may have been motivated by this, making results vulnerable to a self-selection effect. Another limitation is the small difference in our experienced and less experienced categories: experienced = practicing more than 8 years; less experienced = (after adjustment) practising for less than 6 years, leaving a possible difference of just over 2 years. A separate analysis was run to compare actual number of years of practice, but found the same correlations. Failure to recruit less experienced psychiatrists also limited possible analyses of differences between experienced and less experienced physicians. In one study, it appears that using different scoring algorithms for the IAT may affect the direction in which cognitive load moderates the scores on the IAT [58] and it is a limitation of our study that we did not investigate this. On the other hand, there was a striking lack of an effect in any direction of the cognitive load on the IAT results.

Conclusions

Specializing in psychiatry correlates with lower implicit and explicit mental illness prejudice compared to specializing in general medicine. Experience does not correlate with implicit prejudice, but it does correlate with warmer explicit feelings towards the mentally ill and greater concern towards a fictional patient provided they are not described as obese.

In terms of implications for healthcare practice, further research is needed to tease out potentially interacting factors and thus to help the medical profession tailor implicit and explicit bias interventions to specific groups. Psychiatrists may already be less implicitly biased towards the mentally ill when they enter specialization, in which case they will need a different approach to mental health bias training from physicians in other specialties.

What we can recommend now based on our results is that medical education and training should be specifically targeted towards the levels and varieties of implicit bias known to exist in specialities. For instance, medical trainees and experienced physicians may require different forms of explicit and implicit bias training.

Our study points to the strength and potential lack of awareness of obesity prejudice, both implicit and explicit, among physicians. The profession thus needs to raise awareness of the seriousness of this damaging prejudice and its consequences for patients and to consider how to tackle it. Quite apart from the harm and injustice it can engender, it is likely to be highly counter-productive for patient interaction. While efforts have been made recently to develop tailored interventions to combat specific racial implicit stereotypes [59, 60], there has been less research into interventions targeting obesity prejudice. One study found that explicit obesity prejudice increased in students over the four years of medical school [61], suggesting that urgent intervention is required.

Supplementary Information

The online version contains supplementary material available at https://doi. org/10.1186/s12910-022-00815-7.

Additional file 1. Clinical vignette.

Acknowledgements

Contributors: Many thanks are due to Janice Sabin, Diana Burgess and Alex Mauron for their help and advice with a variety of aspects of the study, particularly in the planning stage. We would also like to thank the anonymous reviewers, as well as all the busy physicians who gave up time to participate in the study. *Funders*: We gratefully acknowledge the support of the Swiss National Science Foundation making this study possible under grant number 32003B_149407.

Author contributions

CF performed background research, conceived of the original idea of the study together with SH, applied for the funding together with SH, held preparatory meetings to decide on methodology, co-ordinated between members of the group, organised the project administration, oversaw development of the necessary software, developed the questionnaire, vignette, and video intervention resources, oversaw recruitment of participants, oversaw data collection, wrote the original draft, designed Fig. 1, drafted all tables together with SH, and reviewed and edited the manuscript in response to comments from the other authors. CM developed the software needed for the IATs, advised on data collection and analysis, and contributed to the manuscript via editing and comments. DB recruited participants, managed project administration, and collected data. MS recruited participants, managed project administration, and collected data. TB contributed to deciding on the methodology of the research, advised on data collection and analysis, and contributed to the manuscript via editing and comments. SH conceived of the original idea of the study together with CF, applied for the funding together with CF, supervised the project overall, oversaw and contributed to deciding on the methodology of the research, supervised the recruitment of participants and data collection, analysed the data formally, and contributed to the manuscript via editing and comments and drafting the display of data in the tables together with CF. All authors read and approved the final manuscript.

Funding

SH was awarded grant number 32003B_149407 from the Swiss National Science Foundation http://www.snf.ch/en/Pages/default.aspx. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Availability of data and materials

The datasets generated and analysed during the current study are available in the Yareta repository: https://doi.org/10.26037/yareta:md2ryexqsrchhb2fafgo r6lcmm

Declarations

Ethics approval and consent to participate

The study was submitted to the chairperson of the Ethics Committee for the Swiss Canton of Geneva (CCER—Commission Cantonale de l'Éthique de la Recherche), who reviewed it and declared that it was exempt from a full review by the committee. Participants' full written informed consent was obtained and they were informed that they could withdraw from the study at any time. After completing the second stage of the study, participants were debriefed by the interviewer who explained the true purpose and why it had been necessary to omit a full explanation from the initial description. Any doubts or concerns were clarified and participants were again offered the possibility of withdrawing from the study. All methods were carried out in accordance with relevant quidelines and regulations.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹iEH2 (Institute for Ethics, History and the Humanities), Faculty of Medicine, University of Geneva, Geneva, Switzerland. ²Department of Information Science, Geneva School of Business Administration, Geneva, Switzerland. ³University of Applied Sciences and Arts, Western Switzerland, Delémont, Switzerland. ⁴CISA (Swiss National Centre for Affective Sciences), University of Geneva, Geneva, Switzerland. ⁵Department of Psychology, University of Geneva, Geneva, Switzerland.

Received: 6 May 2022 Accepted: 20 July 2022 Published online: 24 August 2022

References

- Dall'Alba G. Medical practice as characterised by beginning medical students. Adv Health Sci Educ. 1998;3(2):101–18.
- Draper C, Louw G. What is medicine and what is a doctor? Medical students' perceptions and expectations of their academic and professional career. Med Teach. 2007;29(5):e100–7.
- Coulter ID, Wilkes M, Der-Martirosian C. Altruism revisited: a comparison of medical, law and business students' altruistic attitudes. Med Educ. 2007;41(4):341–5.
- Puljak L, Kraljevic JB, Latas VB, Sapunar D. Demographics and motives of medical school applicants in Croatia. Med Teach. 2007;29(8):e227–34.
- Sousa F, Schwalbach J, Adam Y, Gonçalves L, Ferrinho P. The training and expectations of medical students in Mozambique. Hum Resour Health. 2007;5(1):11.
- Odusanya OO, Alakija W, Akesode FA. Socio demographic profile and career aspirations of medical students in a new medical school. Niger Postgrad Med J. 2000;7(3):112–5.
- Bhala N, Curry G, Martineau AR, Agyemang C, Bhopal R. Sharpening the global focus on ethnicity and race in the time of COVID-19. The Lancet. 2020;395(10238):1673–6.
- 8. Chowkwanyun M, Reed AL Jr. Racial health disparities and Covid-19 caution and context. N Engl J Med. 2020;383:201–3.
- Sutin AR, Robinson E, Daly M, Gerend MA, Stephan Y, Luchetti M, et al. Body mass index, weight discrimination, and psychological, behavioral, and interpersonal responses to the coronavirus pandemic. Obesity. 2020;28:1590–4.
- Brownstein M, Madva A, Gawronski B. What do implicit measures measure? Wiley Interdiscip Rev Cogn Sci. 2019;10(5):e1501.
- Dovidio JF, Kawakami K, Johnson C, Johnson B, Howard A. On the nature of prejudice: Automatic and controlled processes. J Exp Soc Psychol. 1997;33(5):510–40.
- Sabin JA, Greenwald AG. The influence of implicit bias on treatment recommendations for 4 common pediatric conditions: pain, urinary tract infection, attention deficit hyperactivity disorder, and asthma. Am J Public Health. 2012;102(5):988–95.
- FitzGerald C, Hurst S. Implicit bias in healthcare professionals: a systematic review. BMC Med Ethics. 2017;18(1):19.
- 14. Nosek BA, Riskind RG. Policy implications of implicit social cognition. Soc Issues Policy Rev. 2012;6(1):113–47.
- Jost JT, Rudman LA, Blair IV, Carney DR, Dasgupta N, Glaser J, et al. The existence of implicit bias is beyond reasonable doubt: a refutation of ideological and methodological objections and executive summary of ten studies that no manager should ignore. Res Organ Behav. 2009;29:39–69.
- Rudman LA, Feinberg J, Fairchild K. Minority members' implicit attitudes: Automatic ingroup bias as a function of group status. Soc Cogn. 2002;20(4):294–320.
- 17. Burgess D, van Ryn M, Dovidio J, Saha S. Reducing racial bias among health care providers: lessons from social-cognitive psychology. J Gen Intern Med. 2007;22(6):882–7.
- Stone J, Moskowitz GB. Non-conscious bias in medical decision making: what can be done to reduce it? Med Educ. 2011;45(8):768–76.
- Shavers VL, Fagan P, Jones D, Klein WM, Boyington J, Moten C, et al. The state of research on racial/ethnic discrimination in the receipt of health care. Am J Public Health. 2012;102(5):953–66.
- Hall WJ, Chapman MV, Lee KM, Merino YM, Thomas TW, Payne BK, et al. Implicit racial/ethnic bias among health care professionals and its influence on health care outcomes: a systematic review. Am J Public Health. 2015;105(12):e60-76.
- Knight M, Bunch K, Kenyon S, Tuffnell D, Kurinczuk JJ. A national population-based cohort study to investigate inequalities in maternal mortality in the United Kingdom, 2009–17. Paediatr Perinat Epidemiol. 2020;34:392–8.
- 22. Zestcott CA, Blair IV, Stone J. Examining the presence, consequences, and reduction of implicit bias in health care: a narrative review. Group Process Intergroup Relat. 2016;19:528–42.
- FitzGerald C, Martin A, Berner D, Hurst S. Interventions designed to reduce implicit prejudices and implicit stereotypes in real world contexts: a systematic review. BMC Psychol. 2019;7:1–12.
- 24. Peris TS, Teachman BA, Nosek BA. Implicit and explicit stigma of mental illness: Links to clinical care. J Nerv Ment Dis. 2008;196(10):752–60.

- Blair IV, Ma JE, Lenton AP. Imagining stereotypes away: the moderation of implicit stereotypes through mental imagery. J Pers Soc Psychol. 2001;81(5):828.
- 26. Nickerson RS. Confirmation bias: a ubiquitous phenomenon in many guises. Rev Gen Psychol. 1998;2(2):175–220.
- Bermingham S, Kippax S. HIV-related discrimination: a survey of New South Wales general practitioners. Aust N Z J Public Health. 1998;22(1):92–7.
- Pettigrew TF. Intergroup contact theory. Annu Rev Psychol. 1998;49(1):65–85.
- Pettigrew TF, Tropp LR. A meta-analytic test of intergroup contact theory. J Pers Soc Psychol. 2006;90(5):751.
- Brener L, von Hippel W, Kippax S. Prejudice among health care workers toward injecting drug users with hepatitis C: does greater contact lead to less prejudice? Int J Drug Policy. 2007;18(5):381–7.
- Burgess DJ. Are providers more likely to contribute to healthcare disparities under high levels of cognitive load? How features of the healthcare setting may lead to biases in medical decision making. Med Decis Mak. 2010;30(2):246–57.
- Moskowitz GB, Li P. Egalitarian goals trigger stereotype inhibition: a proactive form of stereotype control. J Exp Soc Psychol. 2011;47(1):103–16.
- Neumann M, Edelhäuser F, Tauschel D, Fischer MR, Wirtz M, Woopen C, et al. Empathy decline and its reasons: a systematic review of studies with medical students and residents. Acad Med. 2011;86(8):996–1009.
- Burks DJ, Kobus AM. The legacy of altruism in health care: the promotion of empathy, prosociality and humanism. Med Educ. 2012;46(3):317–25.
- Sabin JA, Rivara FP, Greenwald AG. Physician implicit attitudes and stereotypes about race and quality of medical care. Med Care. 2008;46(7):678–85.
- Sabin JA, Marini M, Nosek BA. Implicit and explicit anti-fat bias among a large sample of medical doctors by BMI, race/ethnicity and gender. PLoS ONE. 2012;7(11):e48448.
- Rüsch N, Todd AR, Bodenhausen GV, Corrigan PW. Do people with mental illness deserve what they get? Links between meritocratic worldviews and implicit versus explicit stigma. Eur Arch Psychiatry Clin Neurosci. 2010;260(8):617–25.
- Sabin JA, Stuber J, Rocha A, Greenwald A. Providers' implicit and explicit stereotypes about mental illnesses and clinical competencies in recovery. Soc Work Ment Health. 2015;13(5):495–513.
- Teachman BA, Wilson JG, Komarovskaya I. Implicit and explicit stigma of mental illness in diagnosed and healthy samples. J Soc Clin Psychol. 2006;25(1):75–95.
- Drewniak D, Krones T, Sauer C, Wild V. The influence of patients' immigration background and residence permit status on treatment decisions in health care. Results of a factorial survey among general practitioners in Switzerland. Soc Sci Med. 2016;161:64–73.
- Cooley E, Payne BK. Using groups to measure intergroup prejudice. Pers Soc Psychol Bull. 2017;43(1):46–59.
- Greenwald AG, McGhee DE, Schwartz JL. Measuring individual differences in implicit cognition: the implicit association test. J Pers Soc Psychol. 1998;74(6):1464.
- 43. https://implicit.harvard.edu/implicit/. Project Implicit.
- Bernardes SF, Lima ML. On the contextual nature of sex-related biases in pain judgments: the effects of pain duration, patient's distress and judge's sex. Eur J Pain. 2011;15(9):950–7.
- Teachman BA, Gapinski KD, Brownell KD, Rawlins M, Jeyaram S. Demonstrations of implicit anti-fat bias: the impact of providing causal information and evoking empathy. Health Psychol. 2003;22(1):68.
- Gapinski KD, Schwartz MB, Brownell KD. Can television change anti-fat attitudes and behavior? 1. J Appl Biobehav Res. 2006;11(1):1–28.
- Lincoln TM, Arens E, Berger C, Rief W. Can antistigma campaigns be improved? A test of the impact of biogenetic vs psychosocial causal explanations on implicit and explicit attitudes to schizophrenia. Schizophr Bull. 2008;34(5):984–94.
- Rukavina PB, Li W, Shen B, Sun H. A service learning based project to change implicit and explicit bias toward obese individuals in kinesiology pre-professionals. Obes Facts. 2010;3(2):117–26.
- O'Brien KS, Puhl RM, Latner JD, Mir AS, Hunter JA. Reducing anti-fat prejudice in preservice health students: a randomized trial. Obesity. 2010;18(11):2138–44.

- Swift JA, Tischler V, Markham S, Gunning I, Glazebrook C, Beer C, et al. Are anti-stigma films a useful strategy for reducing weight bias among trainee healthcare professionals? Results of a pilot randomized control trial. Obes Facts. 2013;6(1):91–102.
- Galinsky AD, Moskowitz GB. Perspective-taking: decreasing stereotype expression, stereotype accessibility, and in-group favoritism. J Pers Soc Psychol. 2000;78(4):708.
- Devine PG, Forscher PS, Austin AJ, Cox WT. Long-term reduction in implicit race bias: a prejudice habit-breaking intervention. J Exp Soc Psychol. 2012;48(6):1267–78.
- Gawronski B, Morrison M, Phills CE, Galdi S. Temporal stability of implicit and explicit measures: A longitudinal analysis. Pers Soc Psychol Bull. 2017;43(3):300–12.
- 54. Bertrand M, Chugh D, Mullainathan S. Implicit discrimination. Am Econ Rev. 2005;95:94–8.
- Alkozei A, Killgore WD, Smith R, Dailey NS, Bajaj S, Haack M. Chronic sleep restriction increases negative implicit attitudes toward Arab Muslims. Sci Rep. 2017;7:4285.
- Foroni F, Bel-Bahar T. Picture-IAT versus Word-IAT: level of stimulus representation influences on the IAT. Eur J Soc Psychol. 2010;40(2):321–37.
- Rothermund K, Wentura D. Underlying processes in the implicit association test: dissociating salience from associations. J Exp Psychol Gen. 2004;133(2):139.
- Schmitz F, Teige-Mocigemba S, Voss A, Klauer KC. When scoring algorithms matter: Effects of working memory load on different IAT scores. Br J Soc Psychol. 2013;52(1):103–21.
- 59. Stone J, Moskowitz GB, Zestcott CA, Wolsiefer KJ. Testing active learning workshops for reducing implicit stereotyping of Hispanics by majority and minority group medical students. Stigma Health. 2020;5(1):94.
- 60. Liu FF, Coifman J, McRee E, Stone J, Law A, Gaias L, et al. A brief online implicit bias intervention for school mental health clinicians. Int J Environ Res Public Health. 2022;19(2):679.
- Phelan SM, Puhl RM, Burke SE, Hardeman R, Dovidio JF, Nelson DB, et al. The mixed impact of medical school on medical students' implicit and explicit weight bias. Med Educ. 2015;49(10):983–92.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

