

Archive ouverte UNIGE

https://archive-ouverte.unige.ch

Article scientifique

Article 2022

Published version

Open Access

This is the published version of the publication, made available in accordance with the publisher's policy.

Kardar-Parisi-Zhang Physics and Phase Transition in a Classical Single Random Walker under Continuous Measurement

Jin, Zizhuo Tony; Martin, David G.

How to cite

JIN, Zizhuo Tony, MARTIN, David G. Kardar-Parisi-Zhang Physics and Phase Transition in a Classical Single Random Walker under Continuous Measurement. In: Physical Review Letters, 2022, vol. 129, p. 260603. doi: 10.1103/PhysRevLett.129.260603

This publication URL:https://archive-ouverte.unige.ch/unige:166536Publication DOI:10.1103/PhysRevLett.129.260603

© This document is protected by copyright. Please refer to copyright holder(s) for terms of use.

Kardar-Parisi-Zhang Physics and Phase Transition in a Classical Single Random Walker under Continuous Measurement

Tony Jin^{1,*} and David G. Martin^{2,†}

¹DQMP, University of Geneva, 24 Quai Ernest-Ansermet, CH-1211 Geneva, Switzerland ²Enrico Fermi Institute, The University of Chicago, 933 East 56th Street, Chicago, Illinois 60637, USA

(Received 12 April 2022; accepted 28 November 2022; published 23 December 2022)

We introduce and study a new model consisting of a single classical random walker undergoing continuous monitoring at rate γ on a discrete lattice. Although such a continuous measurement cannot affect physical observables, it has a nontrivial effect on the probability distribution of the random walker. At small γ , we show analytically that the time evolution of the latter can be mapped to the stochastic heat equation. In this limit, the width of the log-probability thus follows a Family-Vicsek scaling law, $N^{\alpha}f(t/N^{\alpha/\beta})$, with roughness and growth exponents corresponding to the Kardar-Parisi-Zhang (KPZ) universality class, i.e., $\alpha_{\text{KPZ}}^{\text{ID}} = 1/2$ and $\beta_{\text{KPZ}}^{\text{ID}} = 1/3$, respectively. When γ is increased outside this regime, we find numerically in 1D a crossover from the KPZ class to a new universality class characterized by exponents $\alpha_M^{\text{ID}} \approx 1$ and $\beta_M^{\text{ID}} \approx 1.4$. In 3D, varying γ beyond a critical value γ_M^c leads to a phase transition from a smooth phase that we identify as the Edwards-Wilkinson class to a new universality class with $\alpha_M^{\text{3D}} \approx 1$.

DOI: 10.1103/PhysRevLett.129.260603

Universality is a pillar concept of statistical physics, classical and quantum alike. The fact that, under renormalization, different microscopic models can lead to the same scale invariant theory has been the key idea for understanding second-order phase transitions. In particular, the concept of universality classes has found an extremely fertile ground within the study of dynamical interfaces for which a scale invariance property has been reported and documented [1,2]. In this context, one particular fixed point has attracted tremendous interest in the previous decades: the Kardar-Parisi-Zhang (KPZ) universality class and its iconic 1/3 growth exponent [3,4] in 1D. Beyond the eponym KPZ equation, it has been found in a variety of models describing growing interfaces such as the ballistic deposition model [5], the Eden model [6,7], or the restricted solid-on-solid model [8]. Perhaps more surprisingly, in recent years, it has also been discovered in a variety of quantum phenomena such as the growth of entanglement entropy in random unitary circuits [9], stochastic conformal field theory [10], noisy fermions [11], and transport properties of dipolar spin ensembles [12] and integrable spin chains [13-16].

Continuous or weak measurement has enjoyed considerable interest in the previous decades within the quantum community as it provides a nondestructive way to obtain information about a given quantum system [17,18]. Its advent led to many interesting applications such as quantum Zeno effects [19], quantum trajectories [20], quantum Maxwell demons [21], or direct observation of quantum jumps [22]. Recently, a number of studies investigated the

consequences of repeated projections or continuous monitoring on the evolution of quantum many-body systems. For systems undergoing both a random unitary evolution and measurements, a result that has aroused considerable interest lately is the existence of a measurement-induced phase transition (MIPT) in the entanglement entropy [23–35]. Most of these contributions focus on entanglement or Rényi entropies, i.e., information-related quantities which are likely salients in classical systems as well. As such, it is natural to wonder whether the same phenomenology of MIPT also features in classical physics.

In this Letter we unveil a connection between KPZ physics and *classical* information theory by studying a single classical random walker undergoing continuous monitoring and, relying on this connection, we show that this system presents a MIPT in 3D.

We first present the framework that we use to model weak, continuous measurements on a generic Markov process. We then focus on the specific case of a single random walker diffusing on a lattice with the occupancy at each site being continuously monitored.

When the measurement rate γ is small, we find in 1D that the standard deviation of the log-probability follows a Family-Vicsek scaling law with roughness and growth exponents corresponding to the KPZ universality class, i.e., $\alpha_{\text{KPZ}}^{\text{1D}} = 1/2$ and $\beta_{\text{KPZ}}^{\text{1D}} = 1/3$, respectively [3]. By performing a perturbative analysis around $\gamma = 0$, we show analytically that this KPZ-like behavior is due to a direct mapping of the dynamics onto the stochastic heat equation (SHE). As γ is increased further, we see numerically in 1D a size-dependent crossover between the KPZ regime and a new universality class characterized by different exponents $\alpha_M^{\rm ID} \approx 1$ and $\beta_M^{\rm ID} \approx 1.4$. In 3D, instead of a crossover, we see a phase transition between a smooth phase that we identify as the EW class and a rough phase with $\alpha_M^{\rm 3D} \approx 1$. We also show that, both in 1D and 3D, the small γ limit can alternatively be thought of as a short time limit $t \ll \gamma^{-1}$ within which the dynamics is described by the KPZ equation.

Continuous monitoring.—We begin by introducing the formalism of continuous monitoring. It is directly inspired from weak measurement and trajectory frameworks of quantum mechanics [17,36–39] and can be thought of as a simple hidden Markov process [40].

In the absence of monitoring, the system undergoes a stochastic dynamics generated by \mathcal{L} on a classical configuration space $\mathcal{M}(\mathcal{C})$ with total number of configurations Ω . The time evolution of the probability distribution \mathcal{P}_t is given by the master equation

$$\frac{d}{dt}\mathcal{P}_t(\mathcal{C}) = \mathcal{L}[\mathcal{P}_t(\mathcal{C})].$$
(1)

We assume that the stationary state is unique and is further given by the maximally entropic state $\mathcal{P}_{\infty} = \Omega^{-1}$. Weak monitoring takes place via an ancilla that couples to the system for a short amount of time δt such that the generated correlation is of order δt as well. Measuring the ancilla's state provides indirect and noisy information about the system which can be used to write a constrained stochastic evolution for \mathcal{P}_t .

Let Ω be the set $\{X_1, ..., X_N\} := \vec{X}$ where the X_j 's can take values ± 1 : +1 corresponds to an occupied site while -1 to an empty one. We suppose that all sites will be independently monitored. The ancilla monitoring site *j* is also described by a random variable A_j which can take binary values $a_j \in \{-1, 1\}$. We denote by $\mathcal{P}(\vec{X}, A_j)$ the joint probability of the union system + ancillae to be in a given configuration. We fix this probability distribution to positively correlate the state of the system and of the ancilla:

$$\mathcal{P}(\vec{X}, A_j) = \mathcal{P}(\vec{X}) \frac{1 + \frac{\sqrt{\gamma \delta t}}{2} A_j X_j}{2}, \qquad (2)$$

where $\mathcal{P}(\vec{X})$ is the reduced probability of the system only. Once a measurement of the ancilla's state has been made with outcome a_j , the probability distribution is updated with probability $1 + (\sqrt{\gamma \delta t}/2)a_i \langle X_j \rangle$ to

$$\mathcal{P}(\vec{X}) \to \mathcal{P}(\vec{X}|A_j = a_j) = \mathcal{P}(\vec{X}) \frac{1 + \frac{\sqrt{\gamma \delta t}}{2} a_j X_j}{1 + \frac{\sqrt{\gamma \delta t}}{2} a_j \langle X_j \rangle}, \quad (3)$$

where $\langle X_j \rangle := \sum_{\{\vec{X}\}} X_j \mathcal{P}_t(\vec{X})$. In the Supplemental Material (SM) [41], we show that repeating this procedure

M times and taking the limit $M \to \infty$, $\delta t \to 0$ while keeping $M\delta t = t$ fixed leads, in the Itō prescription, to the following evolution for the probability distribution

$$d\mathcal{P}_t(\vec{X}) = \frac{\sqrt{\gamma}}{2} \mathcal{P}_t(\vec{X}) (X_j - \langle X_j \rangle_t) dB_t^j, \qquad (4)$$

where dB_t^j are site-independent Brownian processes with variance dt and Itō rules $dB_t^j dB_t^k = \delta_{j,k} dt$. Note that \mathcal{P}_t is both a probability distribution and a stochastic variable. Consequently, there are two types of averages in the problem: $\langle \cdots \rangle$ denotes the average with respect to \mathcal{P}_t , while $\mathbb{E}[\cdots]$ denotes the average with respect to the Brownian processes $\{B_t^j\}$.

As measurements occur independently on every site, we obtain the stochastic evolution of the monitored system as the sum of (1) and (4):

$$d\mathcal{P}_t = \mathcal{L}(\mathcal{P}_t)dt + \frac{\sqrt{\gamma}}{2} \sum_j \mathcal{P}_t(\vec{X})(X_j - \langle X_j \rangle_t) dB_t^j.$$
(5)

Note that, since \mathcal{L} preserves the total probability and $\sum_{\{\vec{X}\}} \mathcal{P}_t(\vec{X})(X_j - \langle X_j \rangle_t) = 0$, the probability distribution \mathcal{P}_t remains normalized at every time *t* for each realization of the process.

Single-particle problem.—We now consider the specific case of a single random walker. For lightness, the following discussion will be for a 1D system of N sites with periodic boundary conditions but generalization to higher dimensions is straightforward. Let $p_j(t)$ be the probability for the particle to be at site j at time t. We choose \mathcal{L} to be the discrete Laplacian weighted by a diffusion constant D, i.e., $\mathcal{L} = D\Delta$ with $\Delta p_j \coloneqq p_{j-1} - 2p_j + p_{j+1}$. Starting from (5), the evolution of p_j in the presence of monitoring is given by

$$dp_{j} = D\Delta p_{j}dt + \sqrt{\gamma}p_{j}dW_{t}^{j}, \qquad (6)$$

with $dW_t^j := dB_t^j - \sum_m p_m dB_t^m$ (see Ref. [41] for details of the calculation). Note that dW_t^j are site-correlated Gaussian noises such that $\mathbb{E}[dW_t^j] = 0$ and $\mathbb{E}[dW_t^j dW_t^k] = (\delta_{j,k} - (p_j + p_k) + \sum_m p_m^2)dt$.

The diffusive term favors the flat, maximally entropic distribution $p_j = 1/N$ while the measurement term favors the *N* pointer states $p_j = \delta_{j,k}$ for fixed $k \in [\![1,N]\!]$. For finite *D* and γ , the stationary distribution of this model is nontrivial and, to the best of our knowledge, not known with a notable exception for N = 2. In the latter case, it turns out that the dynamics is equivalent to the one of a single qubit undergoing both thermal relaxation and quantum measurements and was treated in [44,45].

Equation (6) is reminiscent of the stochastic heat equation (SHE) with multiplicative noise [46] except that

the noise dW_t^j is the sum of a Brownian process and a nonlocal contribution $\sum_m p_m dB_t^m$. Nonetheless, it turns out that there is a formal correspondence between (6) and the SHE in the regime of small γ .

Small γ regime.—To highlight this correspondence, we now perform a perturbative analysis around $\gamma = 0$ of (6) in the infinite system size limit $N \rightarrow \infty$. Suppose p admits the small γ expansion

$$p = p^{(0)} + \sqrt{\gamma} p^{(1)} + \gamma p^{(2)} + \cdots,$$
 (7)

where $p^{(0)}$ is the stationary flat profile of the maximally entropic state, i.e., $p_j^{(0)}(t) = 1/N$, $\forall (j, t)$. Inserting (7) into (6), we obtain the evolution of $p^{(1)}$ as

$$dp_{j}^{(1)} = D\Delta p_{j}^{(1)}dt + \frac{1}{N} \left(dB_{t}^{j} - \sum_{m} \frac{1}{N} dB_{t}^{m} \right).$$
(8)

The term $\sum_{m} (dB_t^m/N)$ has mean 0 and variance 1/N so it is subleading in the limit $N \to \infty$. In this regime, we get

$$dp_j^{(1)} \approx D\Delta p_j^{(1)} dt + p_j^{(0)} dB_t^j.$$
(9)

The evolution of $p^{(2)}$ is obtained in a similar way:

$$dp_{j}^{(2)} = D\Delta p_{j}^{(2)} dt + p_{j}^{(1)} \left(dB_{t}^{j} - \sum_{m} \frac{1}{N} dB_{t}^{m} \right)$$
$$\underbrace{-\frac{1}{N} \sum_{m} p_{m}^{(1)} dB_{t}^{m}}_{:=II}.$$
(10)

As explained above, the variance of I scales as 1/N. The variance of II is given by $\mathbb{E}[(1/N^2)\sum_j (p_j^{(1)})^2]$. Using translational invariance, we have on the other hand that $\mathbb{E}[(p_j^{(1)})^2] = \mathbb{E}[(1/N)\sum_j (p_j^{(1)})^2]$ so there is a factor of *N* between the variance of the multiplicative noise term $p_j^{(1)}dB_t^j$ and II. Thus, in the limit of large *N*, we can neglect I and II to obtain

$$dp_j^{(2)} \approx D\Delta p_j^{(2)} dt + p_j^{(1)} dB_t^j.$$
 (11)

This equation is structurally equivalent to Eq. (9). Thus, to order γ , the discrete SHE with multiplicative noise

$$dp_j = D\Delta p_j dt + \sqrt{\gamma} p_j dB_t^J \tag{12}$$

is a good approximation of (6). Furthermore, the probability p_j of the SHE is connected to the height h_j of the KPZ equation via the Cole-Hopf transformation [3] $h_j := (1/\sqrt{\gamma}) \log p_j$. Indeed, using standard Itō calculus on (12), we readily obtain the stochastic dynamics of h_j as a discretized version of the celebrated KPZ equation (up to a linear shift in time $h_j \rightarrow h_j + \sqrt{\gamma}t$):

$$dh_j = [D\Delta h_j + D\sqrt{\gamma}(\nabla h_j)^2 - \sqrt{\gamma}]dt + dB_t^J, \quad (13)$$

where ∇ is the discrete derivative $\nabla h_j \coloneqq h_{j+1} - h_j$. Note that since $p_j \in [0, 1]$, $h_j \in] -\infty, 0]$. Through its connection to the SHE, and therefore to the KPZ equation, we expect the dynamics of the monitored random walker to share common features with the physics of interface growth. One of the interesting quantities arising in the study of such interfaces is the so-called width *w* defined as

$$w \coloneqq \left[\frac{1}{N} \sum_{j} (h_j - \bar{h})^2\right]^{1/2},$$
 (14)

where $\bar{h} := (1/N) \sum_{j} h_{j}$. Starting from a flat initial profile, the Family-Vicsek (FV) scaling relation [1,47] conjectures that, for scale-invariant interfaces, the width should behave as

$$w \propto N^{\alpha} f\left(\frac{t}{N^{\alpha/\beta}}\right)$$
 (15)

with $f(u) \propto u^{\beta}$ for $u \ll 1$ and $f(u) \propto \text{const}$ for $u \gg 1$. The parameters α and β are, respectively, called the roughening and growth exponents. For models within the KPZ universality class, it has been shown in 1D [3] that $\alpha_{\text{KPZ}}^{\text{1D}} =$ 1/2 and $\beta_{\text{KPZ}}^{\text{1D}} = 1/3$. We thus expect that the width of the log-probability of the monitored random walker will follow (15) with KPZ exponents when γ is small [see Figs. 2(a) and 2(b)].

Importantly, one can alternatively think of the small γ expansion as a short time limit. Indeed, at short times, $t \ll \gamma^{-1}$, the probability profile will be close to the initial flat distribution. If we assume that the leading term in p_j scales like 1/N, it is easy to check that $\mathbb{E}[(\sum_m p_m dB_t^m)^2] \approx O(N^{-1})dt$ so that the contribution of the nonlocal part of dW_t^j is subleading.

However, in the long-time regime $t \gg \gamma^{-1}$, we expect to be pushed out of the KPZ regime as the roughening of the probability profile makes the nonlocal term of the noise grow.

In addition, the mapping to KPZ physics at short times and/or small γ tells us that a roughening phase transition from a smooth to a rough interface should occur in 3D and above [48–51]. Indeed, at small γ , we can neglect the contribution of the nonlocal part of the noise and thus the perturbative dynamic renormalization flow leads to similar flow equations than those of the KPZ equations [3]. In the smooth phase, the roughening term becomes irrelevant so we can safely neglect the nonlocal part of the noise. There, we expect that our model will flow to the same universality



FIG. 1. (a) Log-Log plot of the rescaled width $w/N^{\alpha_M^{DD}}$ as a function of the rescaled time $t/N^{\alpha_M^{DD}/\beta_{KPZ}^{1D}}$ for a measurement rate $\gamma = 4$ in 1D. Exponents: $\alpha_M^{1D} = 1$ and $\beta_{KPZ}^{1D} = 0.34$. Parameters: D = 1, dt = 0.001. (c) Log-Log plot of the rescaled width $w/N^{\alpha_M^{3D}}$ as a function of the rescaled time $t/N^{\alpha_M^{3D}/\beta_{KPZ}^{3D}}$ for a measurement rate $\gamma = 36$ in 3D. Exponents: $\alpha_M^{3D} = 1$ and $\beta_{KPZ}^{3D} = 0.15$. Parameters: D = 1, dt = 0.0002. (a),(c) In accordance with the perturbative analysis, which is valid at short times, the initial growth exponent is always KPZ-like. At intermediate and large \hat{t} , however, the FV scaling of the width flows to a new universality class characterized by $\beta_M^{1D} \simeq 1.4$ and $\alpha_M^{1D} \approx 1$ in 1D or $\beta_M^{3D} \simeq 1.2$ and $\alpha_M^{3D} \approx 1$ in 3D. (b) Linear-log plot of the width w as a function of time t for a measurement rate $\gamma = 4$ in 3D. As $\gamma < \gamma_M^c$, the system is in the smooth EW phase and the width does not show any dependency on the system size. Parameters: D = 1, dt = 0.002.

class as the KPZ equation, i.e., the Edwards-Wilkinson (EW) class. However, this similarity should break down in the roughening phase where we expect (6) to flow to a different universality class than KPZ.

Although the analytical investigation of the strong γ regime is beyond the scope of this Letter, we performed a series of numerical simulations of (6) in 1D and 3D to confirm the previous qualitative reasoning regarding the renormalization flow.

Numerical results.—We started all our simulations with a flat initial profile $p_j(t=0) = 1/N^d$, i.e., $h_j(t=0) = -(d/\sqrt{\gamma}) \log N$ with d being the dimension. We simulated

(6) using a standard Euler-Maruyama scheme and took the logarithm for every single realization to obtain the evolution of the process h_j . Details about the numerical methods, convergence check, and finite-size scaling are provided in the SM [41].

We plot in Fig. 1(a) the rescaled width $\hat{w} = w/N^{\alpha_M^{D}}$ as a function of the rescaled time $\hat{t} = t/N^{\alpha_M^{ID}/\beta_{KPZ}^{ID}}$ in 1D for different system sizes when $\gamma = 4.0$. In agreement with the connection to KPZ at short times, all curves collapse on the power law $t^{\beta_{KPZ}^{ID}}$ at small \hat{t} . However, beyond this regime, the FV scaling of the width flows to a new universality class characterized by $\beta_M^{ID} \approx 1.4$ and $\alpha_M^{ID} \approx 1$.



FIG. 2. (a) Roughening exponent α_M^{1D} as a function of γ for different system sizes. (b) Second growth exponent β_M^{1D} as a function of γ for different system sizes. (c) Roughening exponent α_M^{3D} as a function of γ for different system sizes. (d) Roughening exponent α_{KPZ}^{3D} as a function of γ for different system sizes. (d) Roughening exponent α_{KPZ}^{3D} as a function of γ for different system sizes. (d) Roughening exponent α_{KPZ}^{3D} as a function of γ for different system sizes. Details about the methods used to extract the α 's and β 's are in the SM [41]. Note that, due to numerical limitations, only the roughening exponent was computed for the 3D case. The 1D case (a),(b) shows a size-dependent crossover from the KPZ exponents $\alpha_{KPZ}^{1D} = 1/2$, $\beta_{KPZ}^{1D} = 1/3$ to a new phase with exponents $\alpha_M^{1D} \approx 1.4$. For the 3D case (c), we observe that α_M^{3D} remains constant close to 0 on a finite interval before jumping to $\alpha_M^{3D} \approx 1$ when γ is greater than a critical value $\gamma_M^c \approx 10$. This steplike behavior indicates a phase transition from the EW class to a new universality class in 3D. For comparison, (d) shows the behavior of α_{KPZ}^{3D} as a function of γ for the standard SHE (12) in 3D where we also find $\gamma_{KPZ}^c \approx 10$.

Figure 1(c) is a similar plot but performed in 3D when $\gamma = 36$ and for which the rescaled width and time are, respectively, given by $\hat{w} = w/N^{\alpha_M^{3D}}$ and $\hat{t} = t/N^{\alpha_M^{3D}/\beta_{\text{KPZ}}^{3D}}$. At small \hat{t} , all curves collapse on the expected power law $t^{\beta_{\text{KPZ}}^{3D}}$ while beyond this regime the FV scaling flows to a new universality class characterized by $\beta_M^{3D} \approx 1.2$ and $\alpha_M^{3D} \approx 1$.

Finally, on Fig. 1(b), we plot w as a function of t in 3D for different system sizes when $\gamma = 4$. For this value of γ , the KPZ equation flows toward the smooth EW class where we expect the nonlocal part of the noise to be irrelevant. In agreement with this intuition, Fig. 1(b) shows indeed that the width does not scale with N.

We report in Fig. 2 the critical exponents as a function of γ for simulations performed on 1D and 3D lattices. For the former case [Figs. 2(a) and 2(b)], we observe a sizedependent crossover between the KPZ phase and a new phase characterized by exponents $\alpha_M^{1D} \approx 1$ and $\beta_M^{1D} \approx 1.4$. For the 3D case, we report in Fig. 2(c) the existence of a finite range over which α_M^{3D} is close to 0, thereby indicating the presence of two distinct phases separated by a critical value $\gamma_M^c \approx 10$. For comparison, Fig. 2(d) shows the behavior of $\alpha_{\rm KPZ}^{\rm 3D}$ with respect to γ when simulating the SHE Eq. (12) in 3D where we find $\gamma_{\text{KPZ}}^c \approx 10$. The fact that the two critical values for the SHE and our model are close corroborates our previous qualitative reasoning concerning the smooth phase in 3D. As we are only interested in the existence of a MIPT, we only reported the behavior of α_M^{3D} as the systematic determination of β_M^{3D} is more involved and left for future works.

Conclusion and perspectives.—In this Letter we introduced and studied a model for a single random walker undergoing continuous measurement. In the regime of weak monitoring, we mapped the time evolution of its probability distribution onto the SHE. We deduced that, in this regime, the width of the log-probability follows the FV scaling relation of the KPZ universality class. In 1D, this corresponds to roughening and growth exponents $\alpha_{\text{KPZ}}^{\text{ID}} =$ 1/2 and $\beta_{\text{KPZ}}^{\text{ID}} = 1/3$. Beyond weak monitoring, we numerically find in 1D that increasing γ leads to a crossover from the KPZ class to a new universality class with exponents $\alpha_M^{\text{ID}} \approx 1$ and $\beta_M^{\text{ID}} \approx 1.4$. In 3D, we showed, again numerically, that this crossover becomes a phase transition between a smooth phase that we identify as the EW class and a new phase with $\alpha_M^{\text{3D}} \approx 1$.

Our study is one of the first characterizations of a MIPT in classically monitored systems and opens the door to several interesting questions. It would be most desirable to have a better analytical characterization of the strong γ regime. Since perturbative methods ought to fail there, nonperturbative RG methods such as the one presented in [51] may be employed there.

While we only considered a flat profile, it is known that different initial distributions lead to different universality classes in KPZ physics [4]. Thus, it would be interesting to investigate various initial states such as wedge or Brownian conditions to assess the effect of continuous monitoring on their corresponding exponents.

Finally, while we only studied a single particle, the continuous measurement process (5) is easily generalized to more intricate, many-body interacting problems. A natural extension would be to consider the symmetric simple exclusion process (SSEP), which describes multiple diffusive particles with hard-core repulsion. Interestingly the SSEP can be promoted to a quantum version called the QSSEP [52,53]. The study of both the SSEP and QSSEP would thus provide a unified framework to disentangle the properties specific to quantum and classical systems under continuous monitoring.

T. J. and D. M. thank D. Bernard, N. Caballero, L. Canet, A. Krajenbrink, V. Lecomte, P. Ledoussal, M. Medenjak, C. Nardini, and L. Piroli for useful discussions. T. J. acknowledges support from the Swiss National Science Foundation under Division II.

Note added.—Recently, two works had a similar objective of studying measurement effects on chaotic, classical systems but with a focus on phase transition [54,55].

^{*}zizhuo.jin@unige.ch

[†]dgmartin@uchicago.edu

- A.-L. Barabási and H. E. Stanley, *Fractal Concepts in Surface Growth* (Cambridge University Press, Cambridge, England, 1995).
- [2] G. Ódor, Rev. Mod. Phys. 76, 663 (2004).
- [3] M. Kardar, G. Parisi, and Y.-C. Zhang, Phys. Rev. Lett. 56, 889 (1986).
- [4] I. Corwin, arXiv:1106.1596.
- [5] P. Meakin, P. Ramanlal, L. M. Sander, and R. C. Ball, Phys. Rev. A 34, 5091 (1986).
- [6] E. Murray, in Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability: Held at the Statistical Laboratory, University of California, 1960 (University of California Press, Berkeley 1960), Vol. 2, p. 223.
- [7] M. Kolb, R. Botet, and R. Jullien, Phys. Rev. Lett. 51, 1123 (1983).
- [8] J. M. Kim and J. M. Kosterlitz, Phys. Rev. Lett. 62, 2289 (1989).
- [9] A. Nahum, J. Ruhman, S. Vijay, and J. Haah, Phys. Rev. X 7, 031016 (2017).
- [10] D. Bernard and P.L. Doussal, Europhys. Lett. 131, 10007 (2020).
- [11] T. Jin, A. Krajenbrink, and D. Bernard, Phys. Rev. Lett. 125, 040603 (2020).
- [12] C. Zu, F. Machado, B. Ye, S. Choi, B. Kobrin, T. Mittiga, S. Hsieh, P. Bhattacharyya, M. Markham, D. Twitchen, A. Jarmola, D. Budker, C. R. Laumann, J. E. Moore, and N. Y. Yao, Nature (London) **597**, 45 (2021).

- [13] J. De Nardis, M. Medenjak, C. Karrasch, and E. Ilievski, Phys. Rev. Lett. **124**, 210605 (2020).
- [14] E. Ilievski, J. De Nardis, S. Gopalakrishnan, R. Vasseur, and B. Ware, Phys. Rev. X 11, 031023 (2021).
- [15] D. Wei, A. Rubio-Abadal, B. Ye, F. Machado, J. Kemp, K. Srakaew, S. Hollerith, J. Rui, S. Gopalakrishnan, N. Y. Yao, I. Bloch, and J. Zeiher, Science **376**, 716 (2022).
- [16] Q. Fontaine, D. Squizzato, F. Baboux, I. Amelio, A. Lemaître, M. Morassi, I. Sagnes, L. Le Gratiet, A. Harouri, M. Wouters, I. Carusotto, A. Amo, M. Richard, A. Minguzzi, L. Canet, S. Ravets, and J. Bloch, Nat. Commun. **608**, 687 (2022).
- [17] Y. Aharonov, D. Z. Albert, and L. Vaidman, Phys. Rev. Lett. 60, 1351 (1988).
- [18] J. M. Raimond, M. Brune, and S. Haroche, Rev. Mod. Phys. 73, 565 (2001).
- [19] W. M. Itano, D. J. Heinzen, J. J. Bollinger, and D. J. Wineland, Phys. Rev. A 41, 2295 (1990).
- [20] S. Kocsis, B. Braverman, S. Ravets, M. J. Stevens, R. P. Mirin, L. K. Shalm, and A. M. Steinberg, Science 332, 1170 (2011).
- [21] N. Cottet, S. Jezouin, L. Bretheau, P. Campagne-Ibarcq, Q. Ficheux, J. Anders, A. Auffèves, R. Azouit, P. Rouchon, and B. Huard, Proc. Natl. Acad. Sci. U.S.A. 114, 7561 (2017).
- [22] Z. K. Minev, S. O. Mundhada, S. Shankar, P. Reinhold, R. Gutiérrez-Jáuregui, R. J. Schoelkopf, M. Mirrahimi, H. J. Carmichael, and M. H. Devoret, Nature (London) 570, 200 (2019).
- [23] X. Cao, A. Tilloy, and A. De Luca, SciPost Phys. 7, 024 (2019).
- [24] B. Skinner, J. Ruhman, and A. Nahum, Phys. Rev. X 9, 031009 (2019).
- [25] Y. Li, X. Chen, and M. P. A. Fisher, Phys. Rev. B 100, 134306 (2019).
- [26] M. Szyniszewski, A. Romito, and H. Schomerus, Phys. Rev. B 100, 064204 (2019).
- [27] Y. Bao, S. Choi, and E. Altman, Phys. Rev. B 101, 104301 (2020).
- [28] M. J. Gullans and D. A. Huse, Phys. Rev. Lett. 125, 070606 (2020).
- [29] M. J. Gullans and D. A. Huse, Phys. Rev. X 10, 041020 (2020).
- [30] A. Zabalo, M. J. Gullans, J. H. Wilson, S. Gopalakrishnan, D. A. Huse, and J. H. Pixley, Phys. Rev. B 101, 060301(R) (2020).
- [31] A. Lavasani, Y. Alavirad, and M. Barkeshli, Nat. Phys. 17, 342 (2021).
- [32] M. Buchhold, Y. Minoguchi, A. Altland, and S. Diehl, Phys. Rev. X 11, 041004 (2021).

- [33] X. Turkeshi, A. Biella, R. Fazio, M. Dalmonte, and M. Schiró, Phys. Rev. B 103, 224210 (2021).
- [34] P. Sierant and X. Turkeshi, Phys. Rev. Lett. 128, 130605 (2022).
- [35] Z. Weinstein, Y. Bao, and E. Altman, Phys. Rev. Lett. 129, 080501 (2022).
- [36] J. Dalibard, Y. Castin, and K. Mølmer, Phys. Rev. Lett. 68, 580 (1992).
- [37] K. Jacobs and D. A. Steck, Contemp. Phys. 47, 279 (2006).
- [38] M. Bauer, D. Bernard, and A. Tilloy, J. Stat. Mech. (2014) P09001.
- [39] D. Bernard, T. Jin, and O. Shpielberg, Europhys. Lett. 121, 60006 (2018).
- [40] L. Rabiner and B. Juang, IEEE ASSP Mag. 3, 4 (1986).
- [41] See Supplemental Material at http://link.aps.org/ supplemental/10.1103/PhysRevLett.129.260603 for a derivation of the weak measurement stochastic evolution and additional details about the numerics, which includes Refs. [42,43].
- [42] R. Mannella, arXiv:cond-mat/9709326.
- [43] Experiments and simulations, edited by F. Moss and P. V. E. McClintock, Noise in nonlinear dynamical systems Vol. 3 (Cambridge University Press, Cambridge [Cambridgeshire]; New York, 1989).
- [44] M. Bauer and D. Bernard, Lett. Math. Phys. 104, 707 (2014).
- [45] A. Tilloy, M. Bauer, and D. Bernard, Phys. Rev. A 92, 052111 (2015).
- [46] L. Bertini and N. Cancrini, J. Stat. Phys. 78, 1377 (1995).
- [47] F. Family and T. Vicsek, J. Phys. A 18, L75 (1985).
- [48] K. Moser, J. Kertész, and D. E. Wolf, Physica (Amsterdam) 178A, 215 (1991).
- [49] H. Yan, D. Kessler, and L. M. Sander, Phys. Rev. Lett. 64, 926 (1990).
- [50] M. F. Torres and R. C. Buceta, J. Stat. Mech. (2018) 033208.
- [51] L. Canet, H. Chaté, B. Delamotte, and N. Wschebor, Phys. Rev. Lett. **104**, 150601 (2010).
- [52] M. Bauer, D. Bernard, and T. Jin, SciPost Phys. 6, 45 (2019).
- [53] D. Bernard and T. Jin, Phys. Rev. Lett. 123, 080601 (2019).
- [54] J. Willsher, S.-W. Liu, R. Moessner, and J. Knolle, Phys. Rev. B 106, 024305 (2022).
- [55] A. Pizzi, D. Malz, A. Nunnenkamp, and J. Knolle, arXiv: 2204.03016.