



Article scientifique

Article

2022

Published version

Open Access

This is the published version of the publication, made available in accordance with the publisher's policy.

---

## Heme cross-feeding can augment *Staphylococcus aureus* and *Enterococcus faecalis* dual species biofilms

---

Ch'ng, Jun-Hong; Muthu, Mugil; Chong, Kelvin K L; Wong, Jun Jie; Tan, Casandra A Z; Koh, Zachary J S; Lopez, Daniel; Matysik, Artur; Nair, Zeus J; Barkham, Timothy; Wang, Yulan; Kline, Kimberly Ann

### How to cite

CH'NG, Jun-Hong et al. Heme cross-feeding can augment *Staphylococcus aureus* and *Enterococcus faecalis* dual species biofilms. In: The ISME journal, 2022, vol. 16, n° 8, p. 2015–2026. doi: 10.1038/s41396-022-01248-1

This publication URL: <https://archive-ouverte.unige.ch/unige:168983>

Publication DOI: [10.1038/s41396-022-01248-1](https://doi.org/10.1038/s41396-022-01248-1)

## ARTICLE OPEN

Heme cross-feeding can augment *Staphylococcus aureus* and *Enterococcus faecalis* dual species biofilms

Jun-Hong Ch'ng <sup>1,2,3,4</sup>, Mugil Muthu<sup>1,5</sup>, Kelvin K. L. Chong<sup>5,6</sup>, Jun Jie Wong <sup>5,7</sup>, Casandra A. Z. Tan <sup>5,7</sup>, Zachary J. S. Koh<sup>1</sup>, Daniel Lopez<sup>5</sup>, Artur Matysik <sup>5</sup>, Zeus J. Nair <sup>5</sup>, Timothy Barkham <sup>1,8</sup>, Yulan Wang <sup>9</sup> and Kimberly A. Kline <sup>5,10</sup>✉

© The Author(s) 2022

The contribution of biofilms to virulence and as a barrier to treatment is well-established for *Staphylococcus aureus* and *Enterococcus faecalis*, both nosocomial pathogens frequently isolated from biofilm-associated infections. Despite frequent co-isolation, their interactions in biofilms have not been well-characterized. We report that in combination, these two species can give rise to augmented biofilms biomass that is dependent on the activation of *E. faecalis* aerobic respiration. In *E. faecalis*, respiration requires both exogenous heme to activate the *cydAB*-encoded heme-dependent cytochrome *bd*, and the availability of O<sub>2</sub>. We determined that the ABC transporter encoded by *cydDC* contributes to heme import. In dual species biofilms, *S. aureus* provides the heme to activate *E. faecalis* respiration. *S. aureus* mutants deficient in heme biosynthesis were unable to augment biofilms whereas heme alone is sufficient to augment *E. faecalis* mono-species biofilms. Our results demonstrate that *S. aureus*-derived heme, likely in the form of released hemoproteins, promotes *E. faecalis* biofilm formation, and that *E. faecalis* gelatinase activity facilitates heme extraction from hemoproteins. This interspecies interaction and metabolic cross-feeding may explain the frequent co-occurrence of these microbes in biofilm-associated infections.

The ISME Journal (2022) 16:2015–2026; <https://doi.org/10.1038/s41396-022-01248-1>

## INTRODUCTION

Biofilms consist of a sessile community of microbes embedded within a matrix of extracellular polymeric substances. Biofilms represent the dominant mode of bacterial life, and biofilm cells exhibit different patterns of behaviour compared to planktonic cells [1]. Characteristically, biofilms confer protection to chemical and physiological stresses [2], sheer forces [3, 4], predation [5], and to antibiotic-mediated clearance [6]—rendering biofilm-associated infections difficult to treat. Behaviourally, biofilm-embedded cells are extremely heterogeneous, exhibiting varying levels of metabolic activity (from very active to quiescent) [7, 8], transcriptional profiles (depending on the microenvironment) [9, 10] and higher frequencies of genetic exchange (via transformation, conjugation and transduction) [11, 12].

Biofilms are frequently polymicrobial, comprising of different species, phyla or kingdoms interacting within the complex community [13]. Typically categorized as commensal, antagonistic or synergistic for simplicity, these multi-species interactions are multifaceted and evolve temporally with fluctuations in the microenvironment such as pH, temperature, oxygen, nutrient and waste levels, and quorum-sensing signals [14]. Although in-depth characterization of molecular interactions in polymicrobial

biofilms is limited, controlled studies have enabled the identification of critical mediating compounds important for interspecies interactions. Notable mediators include c-di-GMP [15, 16], AI-2 [17, 18], alarmone ppGpp [19, 20], bacteriocins [21–23], siderophores [24], L-ornithine [25], lactic acid [26], lipoteichoic acid [27, 28], glycans [29] and indole [30, 31], and these have been reviewed elsewhere [14, 32–42].

*E. faecalis* and *S. aureus* are both opportunistic pathogens and are among the leading causes of nosocomial infections [43, 44]. The biofilm-forming potential of each species is well documented [33, 45] and both have been implicated in biofilm-associated infections such as endocarditis [46, 47], urinary tract infections [48, 49] and chronic wounds [50, 51]. Although, *E. faecalis* and *S. aureus* are commonly co-isolated in chronic wounds such as diabetic foot ulcers [52], venous leg ulcers [53] and pressure wounds [54], studies of their interactions are largely limited to the transfer of vancomycin resistance genes from *E. faecalis* to *S. aureus* in clinical settings [55–57]. Therefore, in this study, we explored the molecular interactions between *E. faecalis* and *S. aureus* in biofilms.

Aerobic respiration in *E. faecalis* requires exogenous heme as a cofactor for cytochrome *bd* [58, 59]. Enterococci are unable to

<sup>1</sup>Department of Microbiology and Immunology, National University of Singapore, Singapore, Singapore. <sup>2</sup>Department of Surgery Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore. <sup>3</sup>Infectious Disease Translational Research Program, National University Health System, Singapore, Singapore. <sup>4</sup>Singapore Centre for Environmental Life Sciences Engineering, National University of Singapore, Singapore, Singapore. <sup>5</sup>Singapore Centre for Environmental Life Sciences Engineering, Nanyang Technological University, Singapore, Singapore. <sup>6</sup>Nanyang Technological University Institute for Health Technologies, Interdisciplinary Graduate School, Nanyang Technological University, Singapore, Singapore. <sup>7</sup>Singapore Centre for Environmental Life Sciences Engineering, Interdisciplinary Graduate Program, Nanyang Technological University, Singapore, Singapore. <sup>8</sup>Department of Laboratory Medicine, Tan Tock Seng Hospital, Singapore, Singapore. <sup>9</sup>Singapore Phenome Center, Lee Kong Chian School of Medicine, Nanyang Technological University, Nanyang, Singapore. <sup>10</sup>School of Biological Sciences, Nanyang Technological University, Singapore, Singapore. ✉email: micchn@nus.edu.sg; kline@ntu.edu.sg

synthesize heme due to an absent TCA cycle that prevents the formation of porphyrin precursors. Through a yet unknown importer, heme is taken into the cell and incorporated into cytochrome *bd* (CydAB), which then converts terminal demethyl-quinol (DMKH<sub>2</sub>) to demethylmenaquinone (DMK), consuming O<sub>2</sub> and releasing H<sub>2</sub>O in the process [60]. DMK is reduced by NADH:quinone oxidoreductase back into DMKH<sub>2</sub> consuming NADH. Importantly, Enterococci do not express other membrane-embedded electron carriers like ubiquinone or menaquinone [60], but only make DMK which is a modified menaquinone lacking a 2-methyl group [61]. Cytochrome *bd* is the key respiratory enzyme for *E. faecalis* and contains two subunits (CydA and CydB) with three cytochromes, *b*<sub>558</sub>, *b*<sub>595</sub> and *d*. The translocation of a proton by cytochrome *bd* establishes a proton motive force that, when coupled with F<sub>0</sub>F<sub>1</sub>-ATP synthase, generates ATP [62, 63]. Additionally, *cydC* and *cydD* are necessary for cytochrome *bd* production and have been suggested to be involved in heme transport and/or cytochrome *bd* assembly [60]. Interestingly, reduction of O<sub>2</sub> by DMK induces extracellular superoxide production, the latter of which was inhibited by exogenous heme [64], while a functional electron transport chain sensitizes *E. faecalis* to oxidative burst and decreased its survival in human blood [65].

In this study, we show that *S. aureus*-derived heme is required to activate *E. faecalis* aerobic respiration, leading to augmented *E. faecalis* growth and augmented dual-species biofilm production. We speculate that this interspecies cross-feeding of heme, where one species consumes metabolic end-products from another, may affect mixed species infection outcomes in heme-restricted host and environmental niches.

## MATERIALS AND METHODS

### Bacterial strains and growth conditions

Strain, isolate and transposon library details are available in Supplementary Information. Overnight cultures of *E. faecalis* were grown in Brain Heart Infusion broth (Becton-Dickinson, United States) whereas *S. aureus* grown in Tryptone Soy Broth (Oxoid, England). Agar Technical Powder No. 3 (Oxoid, England) was used for agar plates. Strains were cultured under static conditions at 37 °C. Overnight cultures were spun down and washed once in phosphate-buffered saline prior to normalization. An OD<sub>600</sub> of 1.3 and 3.0 for *E. faecalis* and *S. aureus* respectively gave about 1 × 10<sup>9</sup> CFU/ml. MRSA Select II agar (Bio-Rad Laboratories, United States) was used to select *S. aureus* USA300LAC from mixed-species cultures whereas rifampicin (25 µg/ml) was added to BHI agar to select for OG1RF. Rifampicin (Sigma-Aldrich, United States) was dissolved in methanol to make a stock of 25 mg/ml and stored at −20 °C. Hemin (Sigma-Aldrich, United States) was dissolved in DMSO to make a stock of 25 mg/ml whereas human hemoglobin (Sigma-Aldrich) was dissolved in dH<sub>2</sub>O and filter sterilized to make stock of 10 mg/ml. For hemin and hemoglobin supplementation, a final concentration of 25 µg/ml and 10 µg/ml respectively were used unless otherwise stated.

### Biofilm assays

Normalized cultures of *E. faecalis* and *S. aureus* were mixed (with equal CFU) in the ratio of 1:1 for dual-species biofilms and 8 µl of single- or dual-species cultures were inoculated in 200 µl of Tryptone Soy Broth in a 96-well flat transparent plates (Thermo Fisher Scientific, United States) and incubated under static conditions at 37 °C for five days unless otherwise stated. For anoxic experiments, plates were instead incubated in an anaerobic jar (Merck, Singapore) with a gas pack (Becton Dickinson, United States) and incubated at 37 °C for five days. Details on Crystal Violet staining and CFU determination is available in Supplementary Information. For biofilm oxygen consumption rate (OCR) assays, 5-day biofilms were grown directly in Seahorse XFe96 FluxPak 96-well plates, with 80 µl of inoculated media added per well. Planktonic cells were removed with three washes of 80 µl PBS using a 96-channel pipettor to prevent cross-contamination and the remaining biofilms were resuspended in 80 µl of PBS. After three baseline measurements were taken, 30 µl of fresh TSB was injected into the wells and OCR measured for 1 h using standard parameters.

### Growth kinetics

Normalized cells were diluted 100× in PBS and 8 µl of diluted cells were added to 200 µl TSB in a 96-well plate. The plate was incubated in Tecan Infinite M200 PRO Spectrophotometer at 37 °C for 20 h and absorbance (600 nm) recorded for every 15 min after shaking the plate for 3 s.

### Transposon library screen and transposon mutants

The *E. faecalis* Transposon Library was provided by Gary M. Dunny [66] and *S. aureus* transposon mutants were from Nebraska Transposon Library [67] provided by the Network on Antimicrobial Resistance in *Staphylococcus aureus* (NARSA) and distributed by BEI resources, NIAID. Details on the screening of the *E. faecalis* transposon library [66, 68] are available in Supplementary Information. The primary screen of dual-species biofilms was performed by adding 5 µl of transposon mutants grown overnight and 3 µl of USA300LAC overnight cultures to 200 µl TSB in the 96-well plate. Each plate had controls of parental strains of OG1RF + USA300LAC. Plates were incubated at 37 °C for five days in a humidified incubator and biofilms quantified by crystal violet. *E. faecalis* transposon mutants that showed reduced staining were selected as hits. Confirmation of primary screen hits was done via a secondary screen (three biological replicates) and secondary screen hits that consistently showed reduced dual-species biofilms were subject to further validation. Transposon mutants without defects in growth kinetics and single-species biofilms, but with reduced dual-species biofilms, were shortlisted as validated hits.

### LC-MS for heme quantification from cell pellet

*E. faecalis* strains (OG1RF, *cydA::Tn*, *cydD::Tn* and *cydC::Tn*) were grown overnight in 20 ml of TSB with or without supplementation of 5 µg/ml hemin. Cell pellets were washed twice with PBS before being normalized by OD600. Cell pellets were mixed with 100 µl of lysozyme (10 mg/ml), incubated in a water bath at 37 °C for 1 h and then stored at −80 °C until LC-MS quantification.

*S. aureus* strains (USA300LAC, *hemA::Tn*, *hemL::Tn*, *hemB::Tn* and *hemE::Tn*) were grown overnight in 20 ml of TSB were resuspended in 950 µl of P1 Buffer. 50 µl of lysostaphin (5 mg/ml) was added and incubated at 37 °C for 1 hr and then stored at −80 °C until LC-MS quantification.

LC-MS details are available in Supplementary Information.

### Gelatinase activity assay

Gelatin tubes were made by combining 3 g of gelatin and 3.7 g of BHI powder with 100 ml water that was autoclaved before 3 ml was added into 15 ml tubes. *E. faecalis* colonies stabbed into the gelatin tubes prior to overnight incubation at 37 °C. Tubes were then refrigerated at 4 °C for an additional 30 min before the tubes were gently tilted to observe for liquefaction of gelatin due to gelatinase activity [69]. Any liquefaction was indicative of positive gelatinase activity. Parental strain OG1RF and OG1RFΔ*gelE* served as positive and negative controls respectively, and assays were repeated four times to confirm gelatinase activity.

### Statistics

Statistical analyses used GraphPad Prism 7 software (version 7.03) (GraphPad, United States). The data was analyzed through unpaired students T-test or one-way analysis of variance (ANOVA) with Tukey's post-hoc testing for multiple inter-group comparison, or with Dunnett's post-hoc testing if comparing to a specific control group. Significance for *p* < 0.05 are reported.


### Ethics

All procedures were performed in accordance with Nanyang Technological University Research Integrity Policy. Anonymized and pure bacterial strains used in this study did not require IRB approval.

## RESULTS

### *S. aureus* augments *E. faecalis* growth within biofilms and overall biomass accumulation

Since *S. aureus* and *E. faecalis* are frequently co-isolated during biofilm-associated infections [55], we first examined the consequence of dual species growth on biofilm biomass accumulation. Using a modified polystyrene microtiter assay followed by crystal violet (CV) staining [70], we observed that *E. faecalis* strain OG1RF



**Fig. 1** Single- and dual-species in vitro biofilms. **A** *E. faecalis* (Ef) and *S. aureus* (Sa) biofilms were grown in 96-well plates alone or in combination (1:1) over five days before biofilm was quantified by crystal violet (CV). Day 0 refers to 1 h post-inoculation. Data shows mean and SD ( $N \geq 3$ ) with \*\*\* $p < 0.001$  and \*\*\*\* $p < 0.0001$  when compared to *E. faecalis*-only biofilms of the same time point using 1-way ANOVA with Tukey's post-hoc test. **B** Day 5 biofilm and **(C)** planktonic cells of Ef or Sa or both were collected and CFU/well determined using selective agar. Data shows mean and SD ( $N \geq 6$ ) with \*\*\* $p < 0.001$  and \*\*\*\* $p < 0.0001$  using 1-way ANOVA and Tukey's post-hoc test.


(a rifampicin and fusidic acid resistant derivative of a human oral isolate OG1, commonly used in molecular manipulation and virulence studies and used throughout this study) biofilm biomass peaked at day 1 and plateaued for the remaining four days, whereas *S. aureus* (strain USA300LAC) formed very poor biofilms under these experimental conditions (Fig. 1A). By contrast, following inoculation of *E. faecalis* together with *S. aureus* at a ratio of 1:1, dual-species biofilms showed significantly greater biomass from day 1 and peaked at day 4. Since the greatest difference in dual-species biomass accumulation occurred at day 4 and 5, we chose day 5 for subsequent assays.

To determine if increased bacterial growth within dual species biofilm contributed to the augmented biomass, biofilms were manually disrupted and CFU determined for both single and dual species biofilms. In day 5 biofilms, we observed a statistically significant 5-fold increase in *E. faecalis* CFU within dual-species biofilms compared to mono-species biofilms (Fig. 1B) which correlated with the increased biomass observed by crystal violet staining. *E. faecalis* out-numbered *S. aureus* by approximately 60-fold in dual-species biofilms. Surprisingly, although *S. aureus* produced very little biofilm biomass compared to *E. faecalis* (Fig. 1A), CFU equivalent to that of *E. faecalis* were recovered, suggesting that *E. faecalis* may produce more abundant biofilm matrix than *S. aureus*. Compared to the increased *E. faecalis* biomass and CFU in the presence of *S. aureus*, there were no significant differences in CFU sampled from non-adherent (planktonic) volume of the same biofilm wells, suggesting that *E. faecalis* growth augmentation by *S. aureus* is specific to biofilms (Fig. 1C).

#### Dual-species biofilm augmentation is strain dependent

To investigate whether *S. aureus* augmentation of *E. faecalis* biofilm is a phenomenon unique to the strains used in the initial studies (*E. faecalis* strain OG1RF and *S. aureus* strain USA300LAC), we assayed for biofilm augmentation using five additional commonly used *S. aureus* laboratory strains, as well as ten clinical wound isolates. We observed that 14 out of the 16 tested *S. aureus* strains/isolates showed a significant biofilm biomass increase when co-cultured with *E. faecalis* OG1RF (Fig. 2A and Supplementary Table 1). Newman strain and isolate C37 did not augment mixed species biofilm biomass, potentially because of anti-enterococcal activity or biofilm-restricting properties of these strains that are discussed later. Together, the data indicates that *S. aureus* strain variation influences the nature of the dual-species relationship with *E. faecalis* vis-à-vis biofilm development.

We next investigated if the *E. faecalis* strain variation also affects dual-species biofilm augmentation by testing three additional *E. faecalis* laboratory strains and 28 clinical isolates derived from bloodstream (VRE isolates), wound (TTSHW-EF05 to EF43), urinary tract infections (UTIEF isolates), or the healthy gastrointestinal tract of children (HCG isolates). In contrast to the relative consistent ability of *S. aureus* strains to augment dual-species biofilms, augmentation observed during co-culture with *S. aureus* USA300LAC was much more heterogeneous with only six out of 32 strains/isolates showing augmented biofilms (Fig. 2B and Supplementary Table 2). Of these six strains, the degree of augmentation observed varied between 2.06–3.10 fold (Supplementary Table 2). Of the 26 that did not produce augmented biofilms, 17 showed single-species biofilm levels that were lower



**Fig. 2 Augmentation of biofilms with multiple *E. faecalis* and *S. aureus* laboratory strains and clinical isolates.** **A** Six *S. aureus* laboratory strains (USA300LAC–ISP479) and ten patient isolates (C01–C50) were grown alone or with *E. faecalis* (OG1RF) for five days before biofilms were quantified by crystal violet (CV). Results show biofilm levels normalized to OG1RF-only control, with mean and SD displayed ( $N \geq 3$ ). Points colored green are significantly different to respective single species biofilms by  $p < 0.05$  using 1-way ANOVA with Bonferroni's post-hoc test (details in Supplementary Table 1). **B** Four laboratory strains of *E. faecalis* (OG1RF–V583), together with 28 *E. faecalis* patient isolates (VRE122–TTSHW-EF43) were grown for five days alone or with *S. aureus* (USA300LAC) before biofilm was quantified by CV. Results show biofilm levels normalized to OG1RF-only controls, with mean and SD ( $N \geq 3$ ). Points colored green are significantly different to respective single species biofilms by  $p < 0.05$  using 1-way ANOVA with Bonferroni's post-hoc test (details in Supplementary Table 2).

than OG1RF under the conditions used (defined arbitrarily as  $<0.5\times$  the biofilm levels of OG1RF), five were defined as having high biofilm levels ( $>2\times$  the biofilm levels of OG1RF), and four had similar levels of biofilm to OG1RF ( $>0.5\times$  but  $<2\times$  of OG1RF) (Supplementary Table 2). Taken together, we conclude that many *E. faecalis* isolates are not susceptible to biofilm augmentation by USA300LAC. This is in keeping with other studies showing strain differences (not just species composition) profoundly affects the nature of microbial interactions [71–73].

#### ***E. faecalis* transposon screen identifies *menA* and *cydA* to be crucial for dual-species biofilm augmentation**


To determine the mechanism by which *E. faecalis* CFU are increased in the presence of *S. aureus*, leading to dual-species biofilm augmentation, we screened a near-saturated *E. faecalis* transposon mutant library [66] for mutants that displayed altered dual-species biofilm accumulation, using the same microtiter CV biofilm assay. After secondary screening to eliminate any mutants that were attenuated in biofilm formation, we validated nine mutants in seven unique genes that reduced dual-species biofilms (Fig. 3A).

Amongst these was *epaOX* that encodes a glycosyltransferase involved in the production of cell wall rhamnopolysaccharide


enterococcal polysaccharide antigen (Epa), is responsible for *E. faecalis* biofilm structure and stability in response to antibiotic stress [74], and is a determinant of biofilm-associated antibiotic resistance [75]. Both the transposon mutant and a clean deletion mutant of *epaOX* showed a reduction in dual-species biofilm levels relative to the single species control which was restored to parental strain levels in the plasmid-complemented strain (Supplementary Fig. 1a) after correcting for variations in single species biofilm levels (Supplementary Fig. 1b). As such, it seems likely that *epaOX* is involved in augmented dual-species biofilms through EPS production.

The roles of four additional validated gene products identified in the transposon screen (an ABC transporter, SufB, RpiR, and a Ser/Thr Phosphatase, Fig. 3A) were not pursued. The essentiality of *sufB* had been suggested in some studies [76] but deemed otherwise by others [58], and its involvement in respiration, via FeS cluster assembly, merits future attention. Most importantly, we identified three unique *menA* (OG1RF\_11661) and one *cydA* (OG1RF\_11666) transposon mutants that failed to undergo dual species augmentation. Both genes encode proteins participating in oxidative respiration (Fig. 3B) [77] prompting closer investigation into this process.

A



B



**Fig. 3 Validation of *E. faecalis* transposon library screen hits. A** The parental strain (OG1RF) and nine transposon mutants were grown alone (circles), or together with *S. aureus* (Sa, USA300LAC, triangles) for five days before biofilm levels were determined by crystal violet (CV). Data shows mean and SD relative to OG1RF-only control ( $N \geq 3$ ). \*\* $p < 0.01$ , \*\*\* $p < 0.001$  and \*\*\*\* $p < 0.0001$  when compared to respective OG1RF-only or OG1RF + Sa control biofilms using 1-way ANOVA with Dunnett's post-hoc test. **B** Oxidative respiration of *E. faecalis* requires MenA, CydA, CydB, heme and  $O_2$ .


### *E. faecalis* OG1RF biofilm augmentation by *S. aureus*, heme and hemoglobin requires $O_2$ and *cydABCD*

We hypothesized that *E. faecalis* respiration is necessary for dual-species biofilm augmentation, and to test this, biofilms were cultured in oxic or anoxic conditions. As predicted, dual-species biofilm augmentation was not observed in anoxic conditions (Fig. 4A) but was increased by over 2-fold in oxic conditions (Fig. 4B), demonstrating that oxygen is required for dual-species biofilm augmentation.

Since the utilization of oxygen by *E. faecalis* is dependent cytochrome *bd* which requires the incorporation of exogenous heme as a cofactor (*E. faecalis* does not synthesize heme [63, 78, 79]), we directly investigated the effect of free and conjugated heme (hemin and haemoglobin respectively) on *E. faecalis* biofilms. Supplementation with either source significantly increased biofilm biomass by over two-fold in oxic conditions (Fig. 4B) and was accompanied by an increase in oxygen consumption rate (Supplementary Fig. 2). Additionally, *E. faecalis* biofilm growth kinetics with hemin supplementation were

measured, with a rapid increase in biofilm staining observed to plateau after day 1 (Supplementary Fig. 3). Under anoxic conditions, hemin supplementation minimally impacted biofilm levels and hemoglobin had no effect (Fig. 4A).

Cytochrome *bd* complex consists of two subunits that are encoded by *cydA* and *cydB*. The same operon (Fig. 5A insert) includes *cydC* and *cydD*, which encode an ATP-binding cassette (ABC)-type transporter required for the expression of a functional cytochrome *bd* complex [63, 80]. As a functional *cydABCD* operon was presumed to be required for aerobic respiration to occur, we tested if disruptions of *cydB*, *cydC* and *cydD* would also attenuate the *E. faecalis* biofilm response to both *S. aureus* and hemin. Biofilm augmentation by both hemin and *S. aureus* was significantly impaired in all the tested transposon mutants of the four genes (Fig. 5A), indicating that that a functional operon was required for augmentation. Deletion mutants of *cydB* and *cydD* likewise failed to augment in the presence of hemin and *S. aureus*, with augmentation completely restored in the complementary strains (Fig. 5A). Additionally, the absence of heme- or



**Fig. 4** *E. faecalis* biofilms augmented under oxic and anoxic conditions. *E. faecalis* (Ef) biofilms were grown for five days under **A** anoxic and **B** oxic conditions, alone or in the presence of *S. aureus* (Sa), hemin or hemoglobin (Hb) before biofilm was quantified by crystal violet (CV). Data shows mean and SD of CV absorbance. \*\*\*\* $p < 0.0001$  when compared to Ef-alone control using Dunnett's post-hoc test ( $N \geq 4$ ).

hemoglobin-induced augmentation in the *menA* transposon mutant shows the requirement for demethylmenaquinone for oxidative respiration to occur (Figs. 5A and 3B).



To ensure that the absence of *E. faecalis* biofilm augmentation by *S. aureus* in the *cydABDC* transposon and deletion mutants was not due to their out-competition by *S. aureus*, the planktonic cells in single- and co-cultured wells were enumerated on selective agar after one day. Of the tested strains, only *epaOX::Tn* had lower CFUs compared to the OG1RF parental strain, both when grown alone and in the presence of *S. aureus*, allowing us to exclude out-competition by *S. aureus* as a confounding factor (Supplementary Fig. 4).

To differentiate the involvement of cytochrome *bd* (CydAB) from the heterodimeric ABC transporter (CydDC) in *E. faecalis* biofilm augmentation, intracellular heme was measured in the cell pellets of the respective transposon and deletion mutants and complementary strains by mass spectrometry. Consistent with the inability of *E. faecalis* to synthesize heme, heme was only detected when *E. faecalis* was cultured in TSB supplemented with hemin, but not in the unsupplemented TSB control (Fig. 5B). Intracellular heme was negligible or not detected in any of the *cyd* mutants when grown in hemin-supplemented TSB (Fig. 5B), suggesting that this operon is essential for heme import. Complementation of *cydD*, but not of *cydB*, partially restored intracellular heme levels (Fig. 5B), potentially due to transcriptional differences arising due to *trans* chromosomal complementation with a non-native (*srtA*) promoter. Notwithstanding, partial restoration of intracellular heme levels in the *cydD* complement indicates that the functional product of the *cydDC* operon is required for heme import. Taken together, these data support that aerobic respiration in *E. faecalis* is activated by heme, hemoglobin or *S. aureus*, leading to augmented biofilm formation, and that import of heme into *E. faecalis* is mediated by CydDC.

#### Heme biosynthesis in *S. aureus* is responsible for dual-species biofilm augmentation


Since *E. faecalis* cannot synthesize heme [63, 79], we hypothesized that *S. aureus*-derived heme might enable the activation of cytochrome *bd*, giving rise to augmented dual-species biofilms. If so, *S. aureus* mutants deficient in heme biosynthesis should be unable to augment *E. faecalis* biofilms. We identified *S. aureus* transposon mutants in four key heme biosynthesis enzymes (*hemA*, *hemB*, *hemE* and *hemL*) in the *S. aureus* Nebraska transposon mutant library [67]. To verify the loss of heme biosynthetic activity, we performed growth assays to query the expected growth defect when exogenous heme was limited [81] and quantified heme in cell pellets by LC-MS. Only *hemB::Tn* displayed a growth defect (Supplementary Fig. 5a) that was restored upon heme supplementation (Supplementary Fig. 5b). The *hemB* mutant was also the only one that showed a 40-fold reduction in heme within cell pellets (Supplementary Fig. 5c). These data indicated that the *hemB* transposon mutant was the only bona fide heme-defective mutant of the four. The *hemA*, *hemE* and *hemL* mutants may have acquired secondary mutations, a documented caveat to this library [82], enabling them to bypass these mutations.

Consistent with our hypothesis, the *S. aureus* *hemB* mutant defective in heme biosynthesis did not give rise to augmented dual-species biofilms (Fig. 6). Notably, the single-species parental *S. aureus* biofilm levels were comparable to the *hemB* mutant, suggesting that growth restriction of this mutant in the absence of heme supplementation is unlikely to account for the drastic reduction in dual-species biofilms. However, this does not rule out the possibility that the *hemB* mutant is outcompeted by *E. faecalis* during biofilm formation. Overall, the data indicate that *S. aureus* and *E. faecalis* dual-species biofilm augmentation is dependent on heme biosynthesis in *S. aureus*.



Recognizing the importance of *S. aureus*-derived heme in driving dual-species biofilm biomass augmentation, we sought to determine if heme and hemoglobin similarly augmented *E. faecalis* biofilm CFU as observed in Fig. 1B. As expected, the un-augmented *cvdA::Tn* biofilms had similar CFU to the *E. faecalis*

Of the *E. faecalis* strains and isolates, it was notable that OG1X, a sister clone of OG1RF lacking in gelatinase activity [84, 85], was not augmented by USA300LAC (Fig. 2B). We therefore investigated whether *gelE* was responsible for use of USA300LAC-derived heme. Significant biofilm augmentation of *E. faecalis* OG1RFΔ*gelE* only occurred during supplementation with hemin, but not *S. aureus* or hemoglobin (Fig. 7). This result demonstrates that free heme (in the form of hemin), but not conjugated heme (in the form of hemoglobin) could be used to augment biofilms in the *gelE* mutant. Moreover, the inability of the *gelE* mutant to augment biofilms in the presence of USA300LAC (Fig. 7) suggests



**Fig. 7 Effects of gelatinase E (*gelE*) deletion on heme-source utilization.** Parental strain *E. faecalis* (OG1RF) and respective *gelE* deletion mutant ( $\Delta$ *gelE*), were grown alone or in combination with hemin (25  $\mu$ g/ml), hemoglobin (Hb, 10  $\mu$ g/ml) or *S. aureus* (Sa, USA300LAC) for five days before biofilm was quantified with crystal violet (CV). Data shows mean and SD ( $N \geq 4$ ). \*\*\*\* $p < 0.0001$  when compared to respective Ef controls using 1-way ANOVA with Bonferroni's post-hoc tests.

that heme produced by USA300LAC is also in a conjugated hemoprotein form. Additional post-hoc testing also revealed that hemin-augmented biofilms of *E. faecalis*  $\Delta$ *gelE* were lower than those of WT, likely due to the established role of GelE in *E. faecalis* biofilm [33] that extend beyond its hemoproteolytic activity.

We then selected a subset of *E. faecalis* strains and isolates (20 out of the original 32, and including OG1RF $\Delta$ *gelE* as control) to assay for biofilm augmentation by hemin and hemoglobin. Hemin supplementation augmented biofilms in 17 of the 21 tested strains/isolates (Supplementary Fig. 8 and Supplementary Table 3). The four that showed little to no augmentation by hemin (HCG5A4, HCG9A2, TTSHW-EF12 and TTSHW-EF30) also did not show augmentation with hemoglobin and *S. aureus*, consistent with the requirement for a functional respiratory chain in *E. faecalis* biofilm augmentation by hemoglobin and *S. aureus*. For TTSHW-EF30, intrinsically high biofilm levels ( $\geq$  four-fold that of OG1RF) may have been less amenable to further augmentation. Of the 17 that were augmented by hemin, 10 were also augmented by hemoglobin (Supplementary Fig. 8 and Supplementary Table 3). Of these ten, eight (80%) were gelatinase-positive whereas two (HCG8E9 and TTSHW-EF16) were gelatinase-negative (Supplementary Table 3). In contrast, out of the seven that were augmented by hemin but not by hemoglobin, only V583 was gelatinase-positive. Together these data suggest that gelatinase activity is important for hemoglobin digestion (heme acquisition) but may not be the exclusive enzyme for this activity and may not be sufficient for heme acquisition.

Of the eight gelatinase-positive strains and isolates that were augmented by both hemin and hemoglobin, five were also augmented by *S. aureus* whereas three were not (Supplementary Fig. 8 and Supplementary Table 3). The three that were not augmented by *S. aureus* (TTSHW-EF9, TTSHW-EF18 and TTSHW-EF28) all produced higher levels of biofilms in the unsupplemented *E. faecalis*-only controls (2.6–2.9 $\times$  more biofilms than OG1RF). This implies that the ability to extract heme from hemoglobin does not necessarily result in dual-species biofilm augmentation. Though there appear to be additional factors that contribute to the response of *E. faecalis* clinical isolates to *S. aureus*, it remains clear that *gelE* facilitates heme acquisition by *E. faecalis* from

hemoglobin and *S. aureus* hemoproteins, highlighting a previously unappreciated role for this established virulence factor in *E. faecalis* polymicrobial biofilm formation.

## DISCUSSION

Despite their co-isolation in biofilm associated infections, the only studies characterizing the molecular interaction between *S. aureus* and *E. faecalis* pertain to vancomycin resistance transfer [55–57]. In this study, we show that *S. aureus* and *E. faecalis* synergize to produce more biofilm and demonstrate that this is dependent on the synthesis of heme by *S. aureus* and cross-feeding by *E. faecalis* to exploit this resource for growth in an oxygen-dependent manner.

To appreciate these findings and provide context to the current literature, it is helpful to understand how *E. faecalis* and *S. aureus* interact with other species. *E. faecalis* produces bacteriocin which reduces growth of *C. perfringens* [21] and also hinders the secretion of botulinum neurotoxin of *C. botulinum* [86]. In oropharyngeal candidiasis, *E. faecalis* bacteriocin, EntV, prevents hyphal morphogenesis of *C. albicans* to suppress hyphal-dependent cytotoxicity [87]. Under iron-limiting conditions, *E. faecalis* secretes L-ornithine which promote siderophore synthesis in *E. coli* [25, 88]. Furthermore, *E. faecalis* suppresses host NF- $\kappa$ B-dependent immune activity to promote the virulence of uropathogenic *E. coli* [89], and in early biofilm stages, Al-2 of *E. faecalis* acts as a chemo-attractant and augments *E. coli* biofilms [90]. *E. faecalis* also promotes biofilm matrix production in *P. aeruginosa* [91] through increasing production of exopolysaccharide Pel. In these studies, the effect of *E. faecalis* on other species is highlighted, yet details on how other species influence *E. faecalis* is conspicuously lacking.

The literature on *S. aureus* interactions with other co-colonizing species is more abundant and has been reviewed [92]. In biofilm-associated infections with *Candida albicans*, *S. aureus* exhibits increased resistance to vancomycin [93], increased mucosal adhesion and pathogenicity [94–96], and increased binding to hyphae via hyphal protein Als3p [97]. By contrast, *S. aureus* interactions with other bacteria appear competitive: *Staphylococcus epidermidis* and *Bacillus subtilis* interferes with *S. aureus* quorum sensing by inhibiting the accessory gene regulatory (*agr*) system, resulting in suppression of virulence genes [98–100]. Additionally, *Pseudomonas aeruginosa*, *Streptococcus pneumoniae* and lactic acid bacteria (LAB) produces toxic products such as pyocyanin, LasA and hydrogen peroxide ( $H_2O_2$ ) to inhibit the growth of *S. aureus* [101–104]. LAB also inhibit the colonization of *S. aureus* by competing for host adhesion sites [105]. As countermeasures, *S. aureus* forms small-colony variants to counter *P. aeruginosa* toxins and produces bacteriocins to inhibit LAB [106, 107]. By contrast, *S. aureus* promotes colonization of *Haemophilus influenzae* within the nasal cavity through hemolysis of erythrocytes and the subsequent release of heme and NAD that drives *H. influenzae* growth [108, 109]. Notably, *S. aureus* responds to hemoglobin in nasal secretions to increase colonization by dampening *agr* expression [110].

One challenge in microbial ecology is in understanding how strain differences contribute to mixed species interactions. Here we show that most *S. aureus* strains augment *E. faecalis* biofilm formation but that the converse is not true. Of the 16 tested *S. aureus* strains grown with *E. faecalis*, only Newman and C37 did not show any biofilm augmentation. In the case of Newman, supernatants inhibit biofilm formation by other *S. aureus* strains via a proteinase-sensitive, heat-tolerant and soluble protein [111]. It is possible that this same biofilm-inhibitory protein could also interfere with *E. faecalis* biofilm formation giving rise to antagonism such that dual-species biofilm levels were lower than *E. faecalis*-only biofilms. Of 31 *E. faecalis* strains tested, only five were amenable to biofilm augmentation in dual-species biofilms.

For several strains, this may be due to their intrinsically lower biofilm producing capacity in the first place. However, the majority appeared to be able to form biofilms yet behaved differently from OG1RF under the conditions tested and restricting the generalizability of dual-species augmentation. This emphasizes the importance of strain differences and raises questions about intrinsic differences in heme uptake and/or aerobic respiration capacity in different *E. faecalis* strains.

The majority of transposon insertion mutants identified as deficient in undergoing biofilm augmentation in the presence of *S. aureus* were in *menA* (1,4-dihydroxy-2-naphthoate isoprenyl-transferase, which converts 1,4-dihydroxy-2-naphthoate to demethylmenaquinone) and *cydA* (cytochrome *d* ubiquinol oxidase subunit I), both of which encode proteins involved in oxidative respiration [60]. For respiration to occur in *E. faecalis*, oxygen and heme are required [60, 112], the latter serving as an essential cofactor for cytochrome *bd* (made up of CydA, CydB and three moieties of cofactor heme) and which acts as the terminal demethylquinol (DMKH<sub>2</sub>) oxidase, generating H<sub>2</sub>O from O<sub>2</sub> and establishing a proton motive force to generate ATP [113]. Since *E. faecalis* is unable to produce porphyrins required to synthesize heme [78], heme must be supplied exogenously. In our experiments, heme was supplied in the form of hemin (free heme) or hemoglobin (protein-conjugated heme). Both hemin and hemoglobin augmented *E. faecalis* biofilms only under oxic conditions, leading to the conclusion that heme augments *E. faecalis* biofilms by enabling aerobic respiration. Clinically, this could translate to dual-species biofilm being more prominent in sites where oxygen is available, such as superficial wounds, and less of a concern in anoxic regions like the bladder [114, 115].

Through a very different mechanism, the shift from anaerobic to aerobic respiration because of interspecies interactions has been described for *Aggregatibacter actinomycetemcomitans*. Though primarily fermentative, the opportunistic oral pathogen *A. actinomycetemcomitans* switched to respiratory metabolism when grown in the presence of *Streptococcus gordonii* in response to enhanced O<sub>2</sub> bioavailability during coinfection [116]. Although it was not determined how O<sub>2</sub> levels were augmented, the assumed provision of electron-acceptors (O<sub>2</sub>) from *S. gordonii* (which requires O<sub>2</sub> to produce H<sub>2</sub>O<sub>2</sub>) resulted in increased *A. actinomycetemcomitans* growth and persistence, giving rise to the term “cross-respiration”. Opposite to this, *S. aureus* has been shown to increase biofilms when grown in anoxic conditions via a SrrAB-dependent programmed cell lysis, whereas the inactivation of the heme production (*hemB*) permitted greater biofilm growth under oxic conditions [83]. Differences between the previous *S. aureus* study and ours could be due to differences in biofilm assay conditions, in particular the duration of incubation (22 hr used previously compared to 5 d used in our study).

Although exogenous heme utilization has been established for both oxidative respiration and catalase (KatA) stability and activity in *E. faecalis* [117, 118], no heme import machinery has been defined. Incidental to our main findings is the discovery that *cydDC* is essential for heme import. Previously shown to be essential for cytochrome *bd* assembly in *E. coli*, the CydDC proteins are ABC transporters that export redox-active thiol compounds such as cysteine and glutathione [119–122]. In *E. coli*, it has also been suggested that CydDC may bind heme to enhance ATP hydrolysis and thiol export [78, 123–125]. Given that no intracellular heme could be detected in *E. faecalis* *cydD* or *cydC* mutants cultured in heme-supplemented media, we propose that the CydDC heterodimer may also function as a key, if not the sole, heme importer in this species. However, others have suggested that CydDC is not involved in heme import or cytochrome assembly based on everted *E. coli* membrane vesicles involving a *cydD1* point mutant that showed similar patterns of hemin uptake as those in the parental strain and may instead be important for maintenance of a suitable redox environment in the periplasm for

conversion to heme *d* [126, 127]. Further work is needed to rule out the possibility that cytochrome *bd* assembly require CydDC, which could also impact heme levels in the membrane, or that other import mechanisms may exist since heme-dependent KatA activity has been described in *cydABDC* deletion mutants [78]. Moreover, we only performed heme quantification on whole-cell lysates, and fractionation experiments could ascertain if heme is internalized into the cytosol or directly incorporated into the cell membrane.

Loss-of-function mutations to the *cydABDC* operon are enriched in *E. faecalis* isolates resistant to killing in the hemoglobin-rich environment of whole blood [128] and this has been attributed to the induction of respiration in parental strain of *E. faecalis* in the presence of heme which increased vulnerability to neutrophil killing [65]. It is therefore possible, in the context of host infection, that only within the immune-shielded environment of biofilms can heme utilization by *E. faecalis* safely take place without concomitant increasing their susceptibility to immune clearance – hence the importance of *E. faecalis* augmenting biofilms in response to heme.

In vitro, it is unclear why *E. faecalis* augmentation by *S. aureus* occurs in biofilms but not planktonic cells. It has been reported by others (and our own studies agree, data not shown) that *E. faecalis* planktonic growth is unaffected by heme supplementation [118]. We speculate that the spatially confined microenvironment of biofilms, or the unique transcriptional profile of biofilm cells, results in sensitivity to heme-based growth augmentation.

Biofilm augmentation of many *E. faecalis* strains and isolates by hemin but not by hemoglobin or heme-producing USA300LAC suggests that an important secondary *E. faecalis* determinant is involved in this mixed species interaction. Comparative genomics led us to identify *gelE* as required for *E. faecalis* OG1RF biofilm augmentation in the presence of hemoglobin and USA300LAC. Free heme could augment biofilm formation in the *gelE* deletion mutant, but this strain was unable to extract heme from hemoglobin or *S. aureus* hemoproteins. Together with its established role in biofilms, *gelE* has previously been shown to hydrolyse hemoglobin [129] which would result in heme release and subsequent import, potentially by CydDC, and incorporation into cytochrome *bd* of *E. faecalis* to allow aerobic respiration to augment biofilms. We speculate that *S. aureus* hemoproteins, whether secreted or released during cell death, may likewise need to be hydrolysed for heme release and subsequent uptake by *E. faecalis*.

In summary, we show that augmentation of *S. aureus* and *E. faecalis* dual-species biofilms requires the biosynthesis of heme by *S. aureus* which facilitates oxidative respiration in *E. faecalis*. In most cases, successful cross-feeding of heme (likely in the form of secreted hemoproteins) will require gelatinase-mediated heme acquisition by *E. faecalis*. It is conceivable that the increased energy derived from aerobic respiration enables production of complex and “expensive” extracellular biofilm-associated proteins like EpaOX or surface-anchored pili (EbpABC) [130]. Since *epaOX* (along with *fsrA*, *gelE* and *epal*) correlate with biofilm-associated antibiotic resistance [75], the potential impact of this inter-species interaction on antibiotic efficacy is worth considering, especially since this has been demonstrated for other mixed species interactions [131, 132]. The findings of this study highlight the importance of understanding inter-species interactions in biofilms and underscore the usefulness of identifying potentiating determinants, like heme, to develop interventions relevant in a complex host setting.

## REFERENCES

1. Watnick P, Kolter R. Biofilm, city of microbes. *J Bacteriol*. 2000;182:2675–9.
2. Yin W, Wang Y, Liu L, He J. Biofilms: the microbial “protective clothing” in extreme environments. *Int J Mol Sci*. 2019;20:3423.

3. Islam N, Kim Y, Ross JM, Marten MR. Proteomic analysis of *Staphylococcus aureus* biofilm cells grown under physiologically relevant fluid shear stress conditions. *Proteome Sci.* 2014;12:21.
4. Thomen P, Robert J, Monmeyran A, Bitbol AF, Douarche C, Henry N. Bacterial biofilm under flow: first a physical struggle to stay, then a matter of breathing. *PLoS One.* 2017;12:1–24.
5. Derlon N, Peter-Varbanets M, Scheidegger A, Pronk W, Morgenroth E. Predation influences the structure of biofilm developed on ultrafiltration membranes. *Water Res.* 2012;46:3323–33.
6. Sharma D, Misba L, Khan AU. Antibiotics versus biofilm: an emerging battleground in microbial communities. *Antimicrob Resist Infect Control.* 2019;8:76.
7. Solokhina A, Bruckner D, Bonkat G, Braissant O. Metabolic activity of mature biofilms of *Mycobacterium tuberculosis* and other non-tuberculous mycobacteria. *Sci Rep.* 2017;7:9225.
8. Wan N, Wang H, Ng CK, Mukherjee M, Ren D, Cao B, et al. Bacterial metabolism during biofilm growth investigated by  $^{13}\text{C}$  tracing. *Front Microbiol.* 2018;9:2657.
9. Qi Z, Chen L, Zhang W. Comparison of transcriptional heterogeneity of eight genes between batch *Desulfovibrio vulgaris* biofilm and planktonic culture at a single-cell level. *Front Microbiol.* 2016;7:597.
10. Nett JE, Lepak AJ, Marchillo K, Andes DR. Time course global gene expression analysis of an in vivo *Candida* biofilm. *J Infect Dis.* 2009;200:307–13.
11. Ghigo JM. Natural conjugative plasmids induce bacterial biofilm development. *Nature.* 2001;412:442–5.
12. Molin S, Tolker-Nielsen T. Gene transfer occurs with enhanced efficiency in biofilms and induces enhanced stabilisation of the biofilm structure. *Curr Opin Biotechnol.* 2003;14:255–61.
13. Gabriliska RA, Rumbaugh KP. Biofilm models of polymicrobial infection. *Future Microbiol.* 2015;10:1997–2015.
14. Toyofuku M, Inaba T, Kiyokawa T, Obana N, Yawata Y, Nomura N. Environmental factors that shape biofilm formation. *Biosci Biotechnol Biochem.* 2016;80:7–12.
15. Lori C, Ozaki S, Steiner S, Bohm R, Abel S, Dubey BN, et al. Cyclic di-GMP acts as a cell cycle oscillator to drive chromosome replication. *Nature.* 2015;523:236–9.
16. Mills E, Petersen E, Kulasekara BR, Miller SI. A direct screen for c-di-GMP modulators reveals a *Salmonella Typhimurium* periplasmic L-arginine-sensing pathway. *Sci Signal.* 2015;8:ra57.
17. Ahmed NA, Petersen FC, Scheie AA. Al-2/LuxS is involved in increased biofilm formation by *Streptococcus intermedius* in the presence of antibiotics. *Antimicrob Agents Chemother.* 2009;53:4258–63.
18. Antonova ES, Hammer BK. Quorum-sensing autoinducer molecules produced by members of a multispecies biofilm promote horizontal gene transfer to *Vibrio cholerae*. *FEMS Microbiol Lett.* 2011;322:68–76.
19. Chavez de Paz LE, Lemos JA, Wickstrom C, Sedgley CM. Role of (p)ppGpp in biofilm formation by *Enterococcus faecalis*. *Appl Environ Microbiol.* 2012;78:1627–30.
20. Abranches J, Martinez AR, Kajfasz JK, Chavez V, Garsin DA, Lemos JA. The molecular alarmone (p)ppGpp mediates stress responses, vancomycin tolerance, and virulence in *Enterococcus faecalis*. *J Bacteriol.* 2009;191:2248–56.
21. Han SK, Shin MS, Park HE, Kim SY, Lee WK. Screening of bacteriocin-producing *Enterococcus faecalis* strains for antagonistic activities against *Clostridium perfringens*. *Korean J Food Sci Anim Resour.* 2014;34:614–21.
22. Lemme A, Grobe L, Reck M, Tomasch J, Wagner-Dobler I. Subpopulation-specific transcriptome analysis of competence-stimulating-peptide-induced *Streptococcus mutans*. *J Bacteriol.* 2011;193:1863–77.
23. van der Ploeg JR. Regulation of bacteriocin production in *Streptococcus mutans* by the quorum-sensing system required for development of genetic competence. *J Bacteriol.* 2005;187:3980–9.
24. Kang D, Kirienko NV. Interdependence between iron acquisition and biofilm formation in *Pseudomonas aeruginosa*. *J Microbiol.* 2018;56:449–57.
25. Keogh D, Tay WH, Ho YY, Dale JL, Chen S, Umashankar S, et al. Enterococcal metabolite cues facilitate interspecies niche modulation and polymicrobial infection. *Cell Host Microbe.* 2016;20:493–503.
26. Kubota H, Senda S, Nomura N, Tokuda H, Uchiyama H. Biofilm formation by lactic acid bacteria and resistance to environmental stress. *J Biosci Bioeng.* 2008;106:381–6.
27. Ahn KB, Baik JE, Yun CH, Han SH. Lipoteichoic acid inhibits *Staphylococcus aureus* biofilm formation. *Front Microbiol.* 2018;9:327.
28. Fabretti F, Theilacker C, Baldassarri L, Kaczynski Z, Kropec A, Holst O, et al. Alanine esters of enterococcal lipoteichoic acid play a role in biofilm formation and resistance to antimicrobial peptides. *Infect Immun.* 2006;74:4164–71.
29. Wu H, Zeng M, Fives-Taylor P. The glycan moieties and the N-terminal polypeptide backbone of a fimbria-associated adhesin, Fap1, play distinct roles in the biofilm development of *Streptococcus parasanguinis*. *Infect Immun.* 2007;75:2181–8.
30. Lee J, Jayaraman A, Wood TK. Indole is an inter-species biofilm signal mediated by SdiA. *BMC Microbiol.* 2007;7:42.
31. Chu W, Zere TR, Weber MM, Wood TK, Whiteley M, Hidalgo-Romano B, et al. Indole production promotes *Escherichia coli* mixed-culture growth with *Pseudomonas aeruginosa* by inhibiting quorum signaling. *Appl Environ Microbiol.* 2012;78:411–9.
32. Pereira CS, Thompson JA, Xavier KB. Al-2-mediated signalling in bacteria. *FEMS Microbiol Rev.* 2013;37:156–81.
33. Ch'ng JH, Chong KKL, Lam LN, Wong JJ, Kline KA. Biofilm-associated infection by enterococci. *Nat Rev Microbiol.* 2019;17:82–94.
34. Jenal U, Reinders A, Lori C. Cyclic di-GMP: second messenger extraordinaire. *Nat Rev Microbiol.* 2017;15:271–84.
35. Mathur H, Field D, Rea MC, Cottrell PD, Hill C, Ross RP. Fighting biofilms with lantibiotics and other groups of bacteriocins. *NPJ Biofilms Microbiomes.* 2018;4:9.
36. Hu M, Zhang C, Mu Y, Shen Q, Feng Y. Indole affects biofilm formation in bacteria. *Indian J Microbiol.* 2010;50:362–8.
37. Kuramitsu HK, He X, Lux R, Anderson MH, Shi W. Interspecies interactions within oral microbial communities. *Microbiol Mol Biol Rev.* 2007;71:653–70.
38. Percy MG, Grundling A. Lipoteichoic acid synthesis and function in gram-positive bacteria. *Annu Rev Microbiol.* 2014;68:81–100.
39. Colomer-Winter C, Flores-Mireles AL, Kundra S, Hultgren SJ, Lemos JA. (p)ppGpp and CodY promote *Enterococcus faecalis* virulence in a murine model of catheter-associated urinary tract infection. *mSphere.* 2019;4:e00392–19.
40. Gomes Von Borowski R, Gnoatto SCB, Macedo AJ, Gillet R. Promising antibiofilm activity of peptidomimetics. *Front Microbiol.* 2018;9:2157.
41. Harrison F, Buckling A. Siderophore production and biofilm formation as linked social traits. *ISME J.* 2009;3:632–4.
42. Dworkin J. The medium is the message: interspecies and interkingdom signaling by peptidoglycan and related bacterial glycans. *Annu Rev Microbiol.* 2014;68:137–54.
43. Ruoff KL, de la Maza L, Murtagh MJ, Spargo JD, Ferraro MJ. Species identities of enterococci isolated from clinical specimens. *J Clin Microbiol.* 1990;28:435–7.
44. Valaperta R, Tejada MR, Frigerio M, Moroni A, Ciulla E, Cioffi S, et al. *Staphylococcus aureus* nosocomial infections: the role of a rapid and low-cost characterization for the establishment of a surveillance system. *N Microbiol.* 2010;33:223–32.
45. Archer NK, Mazaitis MJ, Costerton JW, Leid JG, Powers ME, Shirtliff ME. *Staphylococcus aureus* biofilms: properties, regulation, and roles in human disease. *Virulence.* 2011;2:445–59.
46. Dahl A, Bruun NE. *Enterococcus faecalis* infective endocarditis: focus on clinical aspects. *Expert Rev Cardiovasc Ther.* 2013;11:1247–57.
47. Fernandez Guerrero ML, Gonzalez Lopez JJ, Goyenechea A, Fraile J, de Gorgolas M. Endocarditis caused by *Staphylococcus aureus*: a reappraisal of the epidemiologic, clinical, and pathologic manifestations with analysis of factors determining outcome. *Med (Baltim).* 2009;88:1–22.
48. Zheng JX, Bai B, Lin ZW, Pu ZY, Yao WM, Chen Z, et al. Characterization of biofilm formation by *Enterococcus faecalis* isolates derived from urinary tract infections in China. *J Med Microbiol.* 2018;67:60–7.
49. Muder RR, Brennen C, Rihs JD, Wagener MM, Obman A, Stout JE, et al. Isolation of *Staphylococcus aureus* from the urinary tract association of isolation with symptomatic urinary tract infection and subsequent staphylococcal bacteremia. *Clin Infect Dis.* 2006;42:46–50.
50. Fazli M, Bjarnsholt T, Kirketerp-Moller K, Jorgensen B, Andersen AS, Krogfelt KA, et al. Nonrandom distribution of *Pseudomonas aeruginosa* and *Staphylococcus aureus* in chronic wounds. *J Clin Microbiol.* 2009;47:4084–9.
51. Rajkumari N, Mathur P, Misra MC. Soft tissue and wound infections due to *Enterococcus* spp. among hospitalized trauma patients in a developing country. *J Glob Infect Dis.* 2014;6:189–93.
52. Bowler PG, Duerden BI, Armstrong DG. Wound microbiology and associated approaches to wound management. *Clin Microbiol Rev.* 2001;14:244–69.
53. Gjodsbol K, Christensen JJ, Karlsmark T, Jorgensen B, Klein BM, Krogfelt KA. Multiple bacterial species reside in chronic wounds: a longitudinal study. *Int Wound J.* 2006;3:225–31.
54. Cataldo MC, Bonura C, Caputo G, Aleo A, Rizzo G, Geraci DM, et al. Colonization of pressure ulcers by multidrug-resistant microorganisms in patients receiving home care. *Scand J Infect Dis.* 2011;43:947–52.
55. Weigel LM, Donlan RM, Shin DH, Jensen B, Clark NC, McDougal LK, et al. High-level vancomycin-resistant *Staphylococcus aureus* isolates associated with a polymicrobial biofilm. *Antimicrob Agents Chemother.* 2007;51:231–8.
56. Zhu W, Murray PR, Huskins WC, Jernigan JA, McDonald LC, Clark NC, et al. Dissemination of an *Enterococcus* Inc18-Like *vanA* plasmid associated with vancomycin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother.* 2010;54:4314–20.
57. Flannagan SE, Chow JW, Donabedian SM, Brown WJ, Perri MB, Zervos MJ, et al. Plasmid content of a vancomycin-resistant *Enterococcus faecalis* isolate from a

- patient also colonized by *Staphylococcus aureus* with a VanA phenotype. *Antimicrob Agents Chemother*. 2003;47:3954–9.
58. Gilmore MS, Salamzade R, Selleck E, Bryan N, Mello SS, Manson AL, et al. Genes contributing to the unique biology and intrinsic antibiotic resistance of *Enterococcus faecalis*. *mBio*. 2020;11:e02962–20.
  59. Pritchard GG, Wimpenny JW. Cytochrome formation, oxygen-induced proton extrusion and respiratory activity in *Streptococcus faecalis* var. *zymogenes* grown in the presence of haematin. *J Gen Microbiol*. 1978;104:15–22.
  60. Ramsey M, Hartke A, Huycke M. The physiology and metabolism of enterococci. In: Gilmore MS, Clewell DB, Ike Y, Shankar N, editors. *Enterococci: from commensals to leading causes of drug resistant infection*. Boston. 2014.
  61. Baum RH, Dolin ML. Isolation of 2-solaneyl-1,4-naphthoquinone from *Streptococcus faecalis*, 10Cl. *J Biol Chem*. 1965;240:3425–33.
  62. Ritchey TW, Seely HW Jr. Distribution of cytochrome-like respiration in streptococci. *J Gen Microbiol*. 1976;93:195–203.
  63. Winstedt L, Frankenberg L, Hederstedt L, von Wachenfeldt C. *Enterococcus faecalis* V583 contains a cytochrome *bd*-type respiratory oxidase. *J Bacteriol*. 2000;182:3863–6.
  64. Huycke MM, Moore D, Joyce W, Wise P, Shepard L, Kotake Y, et al. Extracellular superoxide production by *Enterococcus faecalis* requires demethylmenaquinone and is attenuated by functional terminal quinol oxidases. *Mol Microbiol*. 2001;42:729–40.
  65. Painter KL, Hall A, Ha KP, Edwards AM. The electron transport chain sensitizes *Staphylococcus aureus* and *Enterococcus faecalis* to the oxidative burst. *Infect Immun*. 2017;85:e00659–17.
  66. Kristich CJ, Nguyen VT, Le T, Barnes AM, Grindle S, Dunny GM. Development and use of an efficient system for random mariner transposon mutagenesis to identify novel genetic determinants of biofilm formation in the core *Enterococcus faecalis* genome. *Appl Environ Microbiol*. 2008;74:3377–86.
  67. Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, et al. A genetic resource for rapid and comprehensive phenotype screening of nonessential *Staphylococcus aureus* genes. *mBio*. 2013;4:e00537–12.
  68. Dale JL, Beckman KB, Willett JLE, Nilson JL, Palani NP, Baller JA, et al. Comprehensive functional analysis of the *Enterococcus faecalis* core genome using an ordered, sequence-defined collection of insertional mutations in strain OG1RF. *mSystems*. 2018;3:e00062–18.
  69. Rahimi N, Poursina F, Ghaziasgar FS, Sepehrpor S, Hassanzadeh A. Presence of virulence factor genes (*gelE* and *esp*) and biofilm formation in clinical *Enterococcus faecalis* and *Enterococcus faecium* isolated from urinary tract infection in Isfahan, Iran. *Gene Rep*. 2018;13:72–5.
  70. O'Toole GA, Pratt LA, Watnick PI, Newman DK, Weaver VB, Kolter R. Genetic approaches to study of biofilms. *Methods Enzymol*. 1999;310:91–109.
  71. Biyikoglu B, Ricker A, Diaz PI. Strain-specific colonization patterns and serum modulation of multi-species oral biofilm development. *Anaerobe*. 2012;18:459–70.
  72. MacKenzie KD, Palmer MB, Koster WL, White AP. Examining the link between biofilm formation and the ability of pathogenic *Salmonella* strains to colonize multiple host species. *Front Vet Sci*. 2017;4:138.
  73. Vanhommerig E, Moons P, Pirici D, Lammens C, Hernalsteens JP, De Greve H, et al. Comparison of biofilm formation between major clonal lineages of methicillin resistant *Staphylococcus aureus*. *PLoS One*. 2014;9:e104561.
  74. Dale JL, Nilson JL, Barnes AMT, Dunny GM. Restructuring of *Enterococcus faecalis* biofilm architecture in response to antibiotic-induced stress. *NPJ Biofilms Microbiomes*. 2017;3:15.
  75. Dale JL, Cagnazzo J, Phan CQ, Barnes AM, Dunny GM. Multiple roles for *Enterococcus faecalis* glycosyltransferases in biofilm-associated antibiotic resistance, cell envelope integrity, and conjugative transfer. *Antimicrob Agents Chemother*. 2015;59:4094–105.
  76. Garsin DA, Urbach J, Hugueta-Tapia JC, Peters JE, Ausubel FM. Construction of an *Enterococcus faecalis* Tn917-mediated-gene-disruption library offers insight into Tn917 insertion patterns. *J Bacteriol*. 2004;186:7280–9.
  77. Huycke MM, Moore D, Joyce W, Wise P, Shepard L, Kotake Y, et al. Extracellular superoxide production by *Enterococcus faecalis* requires demethylmenaquinone and is attenuated by functional terminal quinol oxidases. *Mol Microbiol*. 2001;42:729–40.
  78. Baureder M, Hederstedt L. Genes important for catalase activity in *Enterococcus faecalis*. *PLoS One*. 2012;7:e36725.
  79. Bryan-Jones DG, Whittenbury R. Haematin-dependent oxidative phosphorylation in *Streptococcus faecalis*. *J Gen Microbiol*. 1969;58:247–60.
  80. Winstedt L, Yoshida K, Fujita Y, von Wachenfeldt C. Cytochrome *bd* biosynthesis in *Bacillus subtilis*: characterization of the *cydABCD* operon. *J Bacteriol*. 1998;180:6571–80.
  81. von Eiff C, Heilmann C, Proctor RA, Woltz C, Peters G, Gotz F. A site-directed *Staphylococcus aureus* *hmbB* mutant is a small-colony variant which persists intracellularly. *J Bacteriol*. 1997;179:4706–12.
  82. Hammer ND, Skaar EP. Powerful genetic resource for the study of methicillin-resistant *Staphylococcus aureus*. *mBio*. 2013;4:e00166–13.
  83. Mashruwala AA, Guchte AV, Boyd JM. Impaired respiration elicits SrrAB-dependent programmed cell lysis and biofilm formation in *Staphylococcus aureus*. *Elife*. 2017;6:e23845.
  84. Ike Y, Craig RA, White BA, Yagi Y, Clewell DB. Modification of *Streptococcus faecalis* sex pheromones after acquisition of plasmid DNA. *Proc Natl Acad Sci USA*. 1983;80:5369–73.
  85. Su YA, Sulavik MC, He P, Makinen KK, Makinen PL, Fiedler S, et al. Nucleotide sequence of the gelatinase gene (*gelE*) from *Enterococcus faecalis* subsp. *liquefaciens*. *Infect Immun*. 1991;59:415–20.
  86. Kruger M, Shehata AA, Schrod W, Rodloff A. Glyphosate suppresses the antagonistic effect of *Enterococcus* spp. on *Clostridium botulinum*. *Anaerobe*. 2013;20:74–8.
  87. Graham CE, Cruz MR, Garsin DA, Lorenz MC. *Enterococcus faecalis* bacteriocin EntV inhibits hyphal morphogenesis, biofilm formation, and virulence of *Candida albicans*. *Proc Natl Acad Sci USA*. 2017;114:4507–12.
  88. Hughes ER, Winter SE. *Enterococcus faecalis*: *E. coli*'s siderophore-inducing sidekick. *Cell Host Microbe*. 2016;20:411–2.
  89. Tien BYQ, Goh HMS, Chong KKL, Bhaduri-Tagore S, Holec S, Dress R, et al. *Enterococcus faecalis* promotes innate immune suppression and polymicrobial catheter-associated urinary tract infection. *Infect Immun*. 2017;85:e00378–17.
  90. Laganenka L, Sourjik V. Autoinducer 2-dependent *Escherichia coli* biofilm formation is enhanced in a dual-species coculture. *Appl Environ Microbiol*. 2018;84:e02638–17.
  91. Lee K, Lee KM, Kim D, Yoon SS. Molecular determinants of the thickened matrix in a dual-species *Pseudomonas aeruginosa* and *Enterococcus faecalis* biofilm. *Appl Environ Microbiol*. 2017;83:e01182–17.
  92. Nair N, Biswas R, Gotz F, Biswas L. Impact of *Staphylococcus aureus* on pathogenesis in polymicrobial infections. *Infect Immun*. 2014;82:2162–9.
  93. Harriott MM, Noverr MC. *Candida albicans* and *Staphylococcus aureus* form polymicrobial biofilms: effects on antimicrobial resistance. *Antimicrob Agents Chemother*. 2009;53:3914–22.
  94. El-Azizi MA, Starks SE, Khardori N. Interactions of *Candida albicans* with other *Candida* spp. and bacteria in the biofilms. *J Appl Microbiol*. 2004;96:1067–73.
  95. Peters BM, Noverr MC. *Candida albicans*-*Staphylococcus aureus* polymicrobial peritonitis modulates host innate immunity. *Infect Immun*. 2013;81:2178–89.
  96. Morales DK, Hogan DA. *Candida albicans* interactions with bacteria in the context of human health and disease. *PLoS Pathog*. 2010;6:e1000886.
  97. Peters BM, Jabra-Rizk MA, Scheper MA, Leid JG, Costerton JW, Shirtliff ME. Microbial interactions and differential protein expression in *Staphylococcus aureus*-*Candida albicans* dual-species biofilms. *FEMS Immunol Med Microbiol*. 2010;59:493–503.
  98. Otto M, Echner H, Voelter W, Gotz F. Pheromone cross-inhibition between *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Infect Immun*. 2001;69:1957–60.
  99. Lina G, Boutite F, Tristan A, Bes M, Etienne J, Vandenesch F. Bacterial competition for human nasal cavity colonization: role of *Staphylococcus aureus* *agr* alleles. *Appl Environ Microbiol*. 2003;69:18–23.
  100. Piewngam P, Zheng Y, Nguyen TH, Dickey SW, Joo HS, Villaruz AE, et al. Pathogen elimination by probiotic *Bacillus* via signalling interference. *Nature*. 2018;562:532–7.
  101. Voggu L, Schlag S, Biswas R, Rosenstein R, Rausch C, Gotz F. Microevolution of cytochrome *bd* oxidase in *Staphylococcus* and its implication in resistance to respiratory toxins released by *Pseudomonas*. *J Bacteriol*. 2006;188:8079–86.
  102. Kessler E, Safrin M, Olson JC, Ohman DE. Secreted LasA of *Pseudomonas aeruginosa* is a staphylolytic protease. *J Biol Chem*. 1993;268:7503–8.
  103. Regav-Yochay G, Trzcinski K, Thompson CM, Malley R, Lipsitch M. Interference between *Streptococcus pneumoniae* and *Staphylococcus aureus*: in vitro hydrogen peroxide-mediated killing by *Streptococcus pneumoniae*. *J Bacteriol*. 2006;188:4996–5001.
  104. Ocana VS, de Ruiz Holgado AA, Nader-Macias ME. Growth inhibition of *Staphylococcus aureus* by H<sub>2</sub>O<sub>2</sub>-producing *Lactobacillus paracasei* subsp. *paracasei* isolated from the human vagina. *FEMS Immunol Med Microbiol*. 1999;23:87–92.
  105. Zarate G, Nader-Macias ME. Influence of probiotic vaginal lactobacilli on in vitro adhesion of urogenital pathogens to vaginal epithelial cells. *Lett Appl Microbiol*. 2006;43:174–80.
  106. Biswas L, Biswas R, Schlag M, Bertram R, Gotz F. Small-colony variant selection as a survival strategy for *Staphylococcus aureus* in the presence of *Pseudomonas aeruginosa*. *Appl Environ Microbiol*. 2009;75:6910–2.
  107. Charlier C, Cretenet M, Even S, Le, Loir Y. Interactions between *Staphylococcus aureus* and lactic acid bacteria: an old story with new perspectives. *Int J Food Microbiol*. 2009;131:30–9.
  108. Margolis E, Yates A, Levin BR. The ecology of nasal colonization of *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Staphylococcus aureus*: the role of

- competition and interactions with host's immune response. *BMC Microbiol.* 2010;10:59.
109. Artman M, Domenech E, Weiner M. Growth of *Haemophilus influenzae* in simulated blood cultures supplemented with heme and NAD. *J Clin Microbiol.* 1983;18:376–9.
  110. Pynnonen M, Stephenson RE, Schwartz K, Hernandez M, Boles BR. Hemoglobin promotes *Staphylococcus aureus* nasal colonization. *PLoS Pathog.* 2011;7:e1002104.
  111. Cue D, Junecko JM, Lei MG, Blevins JS, Smeltzer MS, Lee CY. SaeRS-dependent inhibition of biofilm formation in *Staphylococcus aureus* Newman. *PLoS One.* 2015;10:e0123027.
  112. Ritchey TW, Seeley HW. Cytochromes in *Streptococcus faecalis* var. *zymogenes* grown in a haematin-containing medium. *J Gen Microbiol.* 1974;85:220–8.
  113. Hederstedt L, Gorton L, Pankratova G. Two routes for extracellular electron transfer in *Enterococcus faecalis*. *J Bacteriol.* 2020;202:e00725–19.
  114. Martin-Gutierrez G, Rodriguez-Beltran J, Rodriguez-Martinez JM, Costas C, Aznar J, Pascual A, et al. Urinary tract physiological conditions promote ciprofloxacin resistance in low-level-quinolone-resistant *Escherichia coli*. *Antimicrob Agents Chemother.* 2016;60:4252–8.
  115. Giannakopoulos X, Evangelou A, Kalfakakou V, Grammeniatas E, Papandropoulos I, Charalambopoulos K. Human bladder urine oxygen content: implications for urinary tract diseases. *Int Urol Nephrol.* 1997;29:393–401.
  116. Stacy A, Fleming D, Lamont RJ, Rumbaugh KP, Whiteley M. A commensal bacterium promotes virulence of an opportunistic pathogen via cross-respiration. *mBio.* 2016;7:e00782–16.
  117. Frankenberg L, Brugna M, Hederstedt L. *Enterococcus faecalis* heme-dependent catalase. *J Bacteriol.* 2002;184:6351–6.
  118. Baureder M, Barane E, Hederstedt L. In vitro assembly of catalase. *J Biol Chem.* 2014;289:28411–20.
  119. Yamashita M, Shepherd M, Booth WI, Xie H, Postis V, Nyathi Y, et al. Structure and function of the bacterial heterodimeric ABC transporter CydDC: stimulation of ATPase activity by thiol and heme compounds. *J Biol Chem.* 2014;289:23177–88.
  120. Holyoake LV, Poole RK, Shepherd M. The CydDC family of transporters and their roles in oxidase assembly and homeostasis. *Adv Micro Physiol.* 2015;66:1–53.
  121. Pittman MS, Corker H, Wu G, Binet MB, Moir AJ, Poole RK. Cysteine is exported from the *Escherichia coli* cytoplasm by CydDC, an ATP-binding cassette-type transporter required for cytochrome assembly. *J Biol Chem.* 2002;277:49841–9.
  122. Shepherd M. The CydDC ABC transporter of *Escherichia coli*: new roles for a reductant efflux pump. *Biochem Soc Trans.* 2015;43:908–12.
  123. Borisov VB, Gennis RB, Hemp J, Verkhovsky MI. The cytochrome *bd* respiratory oxygen reductases. *Biochim Biophys Acta.* 2011;1807:1398–413.
  124. Baureder M, Hederstedt L. Heme proteins in lactic acid bacteria. *Adv Micro Physiol.* 2013;62:1–43.
  125. Poole RK, Cozens AG, Shepherd M. The CydDC family of transporters. *Res Microbiol.* 2019;170:407–16.
  126. Goldman BS, Kranz RG. ABC transporters associated with cytochrome *c* biogenesis. *Res Microbiol.* 2001;152:323–9.
  127. Cook GM, Poole RK. Oxidase and periplasmic cytochrome assembly in *Escherichia coli* K-12: CydDC and CcmAB are not required for haem-membrane association. *Microbiol (Read).* 2000;146:527–36.
  128. Van Tyne D, Manson AL, Huycke MM, Karanickolas J, Earl AM, Gilmore MS. Impact of antibiotic treatment and host innate immune pressure on enterococcal adaptation in the human bloodstream. *Sci Transl Med.* 2019;11:eaat8418.
  129. Makinen PL, Clewell DB, An F, Makinen KK. Purification and substrate specificity of a strongly hydrophobic extracellular metalloendopeptidase ("gelatinase") from *Streptococcus faecalis* (strain OG1-10). *J Biol Chem.* 1989;264:3325–34.
  130. Smith DR, Chapman MR. Economical evolution: microbes reduce the synthetic cost of extracellular proteins. *mBio.* 2010;1:e00131–10.
  131. Radlinski L, Rowe SE, Kartchner LB, Maile R, Cairns BA, Vitko NP, et al. *Pseudomonas aeruginosa* exoproducts determine antibiotic efficacy against *Staphylococcus aureus*. *PLoS Biol.* 2017;15:e2003981.
  132. Little W, Black C, Smith AC. Clinical implications of polymicrobial synergism effects on antimicrobial susceptibility. *Pathogens.* 2021;10:144.

## ACKNOWLEDGEMENTS

We are grateful to the following people for input and resources provided: Eric Skaar for input on the study, Yap Shao Jun Nicole for contributing unpublished data to the initial characterization of the interaction, and Singapore Phenome Center (Tham Wai Kin, Cheah Yeong Cheng, Liang Xu) for measurement of heme. Bacterial strains, isolates, mutants, and transposon mutant libraries were generously shared with us by Gary Dunny, Jennifer Dale, Hailyn V. Nielsen, Scott J. Hultgren, Phillip I. Tarr, Stéphane Mesnage, Alexander R. Horswill, Yang Liang, Eric P. Skaar, Lynn Hancock, Barnes-Jewish Hospital microbiology laboratory in St. Louis, Tan Tock Seng Hospital in Singapore, American Type Culture Collection, Network on Antimicrobial Resistance in *Staphylococcus aureus* (NARSA), BEI resources and NIAID.

## AUTHOR CONTRIBUTIONS

KKLC and KAK conceived the study. JHC, KKLC, and KAK designed the experiments. JHC and MM conducted and analysed all experiments, with contributions from JJW, CAZT, ZJSK, AM, ZJN and DL. KAK supervised the research. TMSB provided clinical isolates. YW guided the mass spectrometry analyses. JHC wrote the manuscript, with contributions from KKLC, MM, JJW, CAZT, AM, and KK. All co-authors critically reviewed and approved the submitted manuscript.

## COMPETING INTERESTS

This work was carried out at the Singapore Centre for Environmental and Life Science Engineering (SCELSE), whose research is supported by the National Research Foundation Singapore, Ministry of Education, to Nanyang Technological University and the National University of Singapore under its Research Centre of Excellence Programme. This work was also supported by supported by The Lee Foundation Grant, Department of Surgery, Yong Loo Lin School of Medicine, National University of Singapore, Singapore.

## ADDITIONAL INFORMATION

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s41396-022-01248-1>.

**Correspondence** and requests for materials should be addressed to Jun-Hong Ch'ng or Kimberly A. Kline.

**Reprints and permission information** is available at <http://www.nature.com/reprints>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022