



Article
scientifique

Rapport de
cas

2020

Published
version

Open
Access

This is the published version of the publication, made available in accordance with the publisher's policy.

Typical takotsubo syndrome triggered by SARS-CoV-2 infection

Meyer, Philippe; Degrauwe, Sophie; Van Delden, Christian; Ghadri, Jelena-Rima; Templin, Christian

How to cite

MEYER, Philippe et al. Typical takotsubo syndrome triggered by SARS-CoV-2 infection. In: European heart journal, 2020, vol. 41, n° 19, p. 1860. doi: 10.1093/eurheartj/ehaa306

This publication URL: <https://archive-ouverte.unige.ch/unige:163121>

Publication DOI: [10.1093/eurheartj/ehaa306](https://doi.org/10.1093/eurheartj/ehaa306)

Typical takotsubo syndrome triggered by SARS-CoV-2 infection

Philippe Meyer ^{1*}, Sophie Degrauwe ¹, Christian Van Delden², Jelena-Rima Ghadri³, and Christian Templin ³

¹Cardiology Service, Geneva University Hospitals, Geneva, Switzerland;; ²Infectious diseases Service, Geneva University Hospitals, Geneva, Switzerland; and ³Department of Cardiology, University Hospital of Zurich, Zurich, Switzerland

*Corresponding author. Cardiology Service, Geneva University Hospitals, Rue Gabrielle-Perret-Gentil 4, 1205 Geneva, Switzerland. Tel: +41 22 372 95 97, Fax: +41 22 372 37 45, Email: philippe.meyer@hcuge.ch

Takotsubo syndrome (TTS) is an important differential diagnosis of myocardial injury, which may play a significant role in the COVID-19 pandemic. This report describes a typical case of TTS triggered by SARS-CoV-2 infection.

An 83-year-old lady treated for chronic hypertension was hospitalized for acute chest pain on 18 March 2020, 21 days after the COVID-19 outbreak started in Geneva, Switzerland. She described a 5/10 oppressive mediasternal non-radiating chest discomfort associated with mild breathlessness and dry cough since 15 March. Physical examination was unremarkable. ECG (*Panel A*) showed <1 mm ST-segment elevation in all precordial leads with deep T-wave inversions. High-sensitive cardiac troponin T was elevated at 1142 ng/L (<14 ng/L). Chest X-ray did not detect any pulmonary opacity. Echocardiography revealed typical left ventricular apical ballooning with hyperkinetic basal segments. Coronary angiography showed non-significant lesions (*Panel B*) with a typical takotsubo syndrome (TTS) image on ventriculography (*Panel C*). At day 3, the patient started developing fever, showing increasing biological signs of inflammation, and clear bilateral lung X-ray opacities (*Panel D*). Nasopharyngeal

swab was negative for SARS-CoV-2, but the initial positive immunoglobulin A and negative immunoglobulin G serology pattern proved acute infection. The patient recovered progressively on conventional heart failure medication without the need for oxygen/ventilation, and echocardiography showed only mild residual apical hypokinesis on the day of discharge (day 10).

This is to our knowledge the first case of TTS described in the COVID-19 pandemic. The huge emotional stress at the population level and respiratory infections caused by COVID-19 may represent potential triggers in this context. Myocardial injury, frequently reported in patients with COVID-19, is usually attributed to sepsis and/or hypoxaemia and/or underlying coronary artery disease, as well as myocarditis. We believe that TTS may also play a significant role in the COVID-19 pandemic.

We would like to thank Nicolas Johnner, MD who helped in designing the figure.

Authors' contributions: all authors participated in writing and approved the final manuscript. Written consent for publication was obtained from the patient.

