



Article scientifique

Article

2017

Published version

Open Access

This is the published version of the publication, made available in accordance with the publisher's policy.

Modification of the Surgical Strategy for the Dissection of the Recurrent Laryngeal Nerve Using Continuous Intraoperative Nerve Monitoring

Marin Arteaga, Andrés; Pelsoni, Giuseppe; Leuchter, Igor; Bedat, Benoît; Karenovics, Wolfram; Triponez, Frédéric; Sadowski, Samira Mercedes

How to cite

MARIN ARTEAGA, Andrés et al. Modification of the Surgical Strategy for the Dissection of the Recurrent Laryngeal Nerve Using Continuous Intraoperative Nerve Monitoring. In: World Journal of Surgery, 2017. doi: 10.1007/s00268-017-4277-z

This publication URL: <https://archive-ouverte.unige.ch/unige:99635>

Publication DOI: [10.1007/s00268-017-4277-z](https://doi.org/10.1007/s00268-017-4277-z)

Modification of the Surgical Strategy for the Dissection of the Recurrent Laryngeal Nerve Using Continuous Intraoperative Nerve Monitoring

Andres Marin Arteaga^{1,2} · Giuseppe Peloni^{2,3} · Igor Leuchter⁴ · Benoit Bedat² · Wolfram Karenovics² · Frederic Triponez² · Samira Mercedes Sadowski²

© Société Internationale de Chirurgie 2017

Abstract

Background The aim of this study was to describe first experiences and changes in management using continuous intraoperative neuromonitoring (C-IONM) in thyroid and parathyroid surgery.

Method Retrospective analysis of patients who underwent surgery with C-IONM since 2012. Surgical maneuvers were modified when electrophysiologic events occurred. Patients with persistent loss of signal (LOS) underwent postoperative laryngoscopy.

Results One hundred and one patients (of 1586 neck surgeries) were included and 19 had events: In 13 these were temporary (resolved before end of surgery) and led to intraoperative modifications in surgical approach; in all cases traction was released, and in 8, recurrent laryngeal nerve (RLN) approach was changed [superior approach (2), inferior approach (2), both (4)]. Six patients had persistent LOS (5.9%, present at end of procedure), with RLN palsy (RLNP) on postoperative day 1: In three, LOS occurred at electrode placement on the vagus nerve, leading to distal placement of the electrode allowing ipsilateral dissection under continuous monitoring; all three had complete recovery at 6 months. In the three other patients, LOS occurred on the RLN: one probable thermal, one traction lesion and one accidental section of the anterior RLN branch. The RLN recovered within 6 months in two patients, and in the third, RLNP persisted after 6 months (1/101 = 1%).

Conclusion C-IONM provides real-time evaluation of the RLN function, allowing for adaptation of surgical maneuvers to prevent RLNP. It seems particularly useful in difficult cases like redo neck surgery, invasive thyroid cancer and intrathoracic or large goiter. Care should be given at electrode placement on the vagus nerve.

Introduction

Recurrent laryngeal nerve palsy (RLNP) is a feared complication of neck surgery and has decreased secondary to routine visual identification of the RLN [1, 2]. Impairment in voice function after unilateral RLNP can greatly affect quality of life and is fortunately transitory in most cases. This is not the case in the dreaded bilateral RLNP [3, 4]. There continues to be a debate regarding intermittent intraoperative neuromonitoring of the RLN (I-IONM), as in recent literature on the systematic use of I-IONM there was no reduction in RLNP rates [5] and the number of patients needed to be treated to avoid a RLN is relatively high [2]. A recent meta-analysis and a randomized study

Paper presented at the IAES/WCS congress in Basel, August 2017.

✉ Samira Mercedes Sadowski
samira.sadowski@hcuge.ch

- 1 General Surgery, University Hospitals of Geneva, Geneva, Switzerland
- 2 Thoracic and Endocrine Surgery, University Hospitals of Geneva, Rue Gabrielle Perret-Gentil 4, 1211 Geneva 14, Switzerland
- 3 Ospedale Regionale di Mendrisio, Via Turconi 23, 6850 Mendrisio, Switzerland
- 4 Ear-Nose-Throat Division, University Hospitals of Geneva, Geneva, Switzerland

show benefit in reducing transient RLNP in complex cases [6, 7]. Another advantage of I-IONM is the prevention of bilateral RLNP, with implementation of strict operative procedures for two-staged thyroidectomy [8] and standardized procedures for I-IONM [9].

Continuous intraoperative neuromonitoring (C-IONM) is a more recent technique and represents the potential for incremental improvement, potentially enabling the surgeon to react before irreversible damage to the RLN occurs [10, 11]. An electrode is placed on the vagus nerve during the entire operation in order to measure the electromyographic activity of the muscles innervated by the RLN. The lesions to the RLN are most frequently caused by traction, followed by clamping, transection or thermal injury, as shown in a recent multicenter study with 80% of 115 cases with loss of signal (LOS) caused by traction to the RLN [12]. The difference between each type of injury is critical when it comes to C-IONM because not all of them can be prevented by IONM, and because traction injuries, if tension is released when detected early by IONM, are reversible. This has been shown in porcine models [13, 14] and in humans, with Phelan [11] and Schneider et al. [10] showing that more than 73 and 82% of events, respectively, were reversible during surgery when prompt corrective measures were applied. When comparing RLNP between I-IONM and C-IONM, they conclude that C-IONM offers a higher sensitivity for early postoperative RLNP [10]. Furthermore, a safety analysis of 400 vagal nerve dissections and electrode positioning was reported to be safe [15]. On the other hand, Terris et al. [16] report one case of bradycardia and hypotension using the vagal electrode and one case of trauma to the vagal nerve by electrode application with LOS resulting in temporary RLNP, concluding this to be an invasive and harmful technique. Another group looking at high-risk neck surgeries reports one LOS caused by the vagal electrode and multiple situations of electromyographic (EMG) artifacts [17].

C-IONM was introduced in Geneva in 2012 and used selectively in difficult surgical cases such as re-interventions, invasive cancer, retrosternal goiter and Graves' disease. In the present study, we aim to present our experience with C-IONM.

Methods

This is a retrospective single-center study of consecutive neck surgery cases with C-IONM conducted between August 2012 and October 2016 at the University Hospitals of Geneva. This study was approved by the Institutional Review Board.

The University Hospitals of Geneva are a tertiary center with approximately 400 thyroid and parathyroid surgeries

per year, and the use of I-IONM is standard of care since 2008 [9], as well as two-staged thyroidectomy in cases of LOS on the first side, as published by our group [8].

The choice to use C-IONM was decided intraoperatively by the surgeon, based on expected difficulties, and used selectively in difficult cases. Combined use of handheld probe for RLN identification and continuous electrode on the vagal nerve was done in all cases. The vagal nerve was dissected for the placement of a 2- or 3-mm-diameter-sized automatic periodic stimulation (APS[®]) electrode (Medtronic), with baseline calibration, and the size of the electrode was decided by the surgeon after vagal nerve visualization. Patients were intubated using commercial EMG endotracheal tubes, and NIM-Response 3.0 system was used for monitoring (Medtronic Inc., Jacksonville, USA).

All procedures were performed by two surgeons (each with annual volume >100/year), using a standard approach: lateral approach to the RLN after medializing the thyroid and using energy devices (harmonic focus[®]). If significant events associated with a surgical maneuver occurred, that maneuver was immediately aborted and another approach was performed. A LOS was defined as a response in EMG below the threshold of 100 μ V. An event was defined as a decrease >50% of EMG amplitude from baseline or an increase in latency of more than 10%, with an acoustic signal to the surgeon. LOS was classified into type 1 (segmental) and type 2 (diffuse) injuries. All events and changes in surgical maneuvers were recorded. In cases of LOS, direct laryngoscopy (DL) was performed by our ear-nose-throat (ENT) specialist at postoperative day (POD) 1, and at follow-up. Current practice is follow-up at POD10 and patients with LOS are enrolled in a protocol with ENT follow-up of RLNP and therapy, or until resolution of paresis. Transient and permanent RLNP was defined as resolved within or present at 6 months after surgery, respectively.

Exclusion criteria were patients with RLNP on preoperative laryngoscopy and thyroid cancer invading the RLN (intentional section).

Descriptive analyses were performed to determine the incidence of RLNP and changes in surgical approach. Fischer's exact test was used for comparisons. Two-tailed $P < .05$ was considered statistically significant. Continuous data are presented as mean \pm standard deviation (SD) or median [range], as appropriate. Statistical analyses were performed using SPSS, version 21 (IBM[®]).

Results

Between August 2012 and October 2016, 101 patients with 159 nerves at risk were included using C-IONM, from a total of 1586 neck surgeries performed over the same time

period. Mean age of patients was 55 ± 16.8 years, with 65.4% being female, as shown in Table 1. The average time of surgery was 156.6 ± 63.5 min and the most frequent indication for surgery was benign disease (63.4%, comprising 24.8% of retrosternal or large compressive goiters), and suspected malignancy in 36.6%. Overall, 41.6% total thyroidectomies were performed and 14.9% with neck dissections, 25.7% unilateral lobectomies, 8.9% completion surgeries, 4.0% neck dissections and 4.9% parathyroidectomies. On pathology, 36.6% had malignant disease, with 14 patients presenting stage III and IV disease.

Loss of signal with C-IONM and modification of surgical maneuver

In all cases, the electrode placement was successfully achieved after dissection of the vagal nerve. C-IONM did not result in any intraoperative anesthesiological events. Overall, in 19 cases an event was observed ($19/101 = 18.8\%$). Thirteen patients had temporary events (resolved before end of surgery) and six had a persistent LOS (no signal at the end of the procedure). In all, the current maneuver provoking the events was aborted and traction released to wait for an increase in signal.

The 13 temporary events led to the following modifications in surgical approach: In five cases, traction was released, and in eight cases, in addition to traction release, the surgical strategy for RLN approach was changed, in two patients, a superior approach, in two an inferior approach and in four both approaches were used.

The six patients with persistent LOS ($6/101 = 5.9\%$) are listed in Table 2. All six had a unilateral RLNP confirmed at DL on POD1. Histopathology showed one case with recurrent central lymph nodes of papillary thyroid cancer (PTC), one PTC with lateral and central lymph nodes, one PTC, two large goiters and one case of Grave's disease. In three cases LOS occurred at electrode placement, with the vagus nerve anatomically intact and no visible lesions (cases 4–6). The IONM signal was lost on the proximal side of the electrode, which led to distal placement of the electrode on the vagus nerve. Ipsilateral dissection was then performed with C-IONM. In two cases, the planned bilateral surgery was interrupted after the first side due to LOS, with RLNP confirmed at POD1. They both had a complete recovery of RLN within 2 months, followed by surgery on the second side. In the third case, with previous lateral and central neck dissection, DL showed RLNP on POD1, but complete recovery at 6 months.

In the other three cases (cases 1–3), only lobectomy was planned or LOS happened on the second side: One case had LOS probably due to traction with an anatomically intact RLN, and RLNP at POD1 but full recovery within

Table 1 Clinical characteristics of cohort with C-IONM ($n = 101$)

Variables	Values
Mean age in years	55 \pm 16.8
Female gender	66 (65.4)
Mean body mass index (kg/m ²)	26.3 \pm 5.4
ASA score	
ASA1	10 (9.9)
ASA2	80 (79.2)
ASA3	11 (10.9)
Mean duration of surgery (min)	156.6 \pm 63.5
Indication for surgery	
Benign ^a	52 (51.5)
Suspect nodule ^b	15 (14.9)
Malignant (PTC, FTC, MTC, ATC)	22 (21.8)
Grave's disease, Hashimoto thyroiditis	12 (11.9)
Type of surgery	
Total thyroidectomy	42 (41.6)
Thyroidectomy and neck dissection	15 (14.9)
Unilateral lobectomy	26 (25.7)
Completion thyroidectomy	9 (8.9)
Neck dissections	4 (4.0)
Parathyroidectomy	5 (4.9)
Pathology	
Benign ^c	54 (53.5)
Grave's disease, Hashimoto thyroiditis	10 (9.9)
Malignant	37 (36.6)
PTC (including micro-papillary)/FVPTC	15/14
FTC	2
MTC	2
PDTC/ATC	3/1
Stage ^d I/II/III/IVa + IVb	21/2/6/7 + 1
Number of nerves at risk, total	159
Unilateral	43
Bilateral	58
EMG events, total	19
Events, temporary	13 (12.9)
Persistent LOS	6 (5.9)
RLNP	
Temporary	6 (5.9)
Permanent	1 (1.0)

Values are presented as mean \pm standard deviation or n (%)

^aMultinodular goiter, toxic nodules, chronic thyroiditis, cystic lesions, hyperparathyroidism

^bSuspect nodules (Bethesda classification III–V) and non-specific tumors

^cThyroid hyperplasia and follicular adenoma, multinodular goiter, parathyroid hyperplasia and adenoma, cysts

^dAccording to the AJCC Cancer staging Manual, 7th edition [25]

ASA American Society of Anesthesiologists, ATC anaplastic thyroid carcinoma, EMG electromyographic, FTC follicular thyroid carcinoma, FVPTC follicular variant PTC, MTC medullary thyroid carcinoma, PDTC poorly differentiated thyroid carcinoma, PTC papillary thyroid carcinoma, RLNP recurrent laryngeal nerve palsy

1 month; one case had possible thermal lesion, with RLNP at DL and recovery within 6 months; in the last case, there was an accidental section of the anterior branch of RLN, with suture of nerve extremities during surgery, and with persistent RLNP at DL. Overall, there was one persistent RLNP in this cohort ($1/101 = 1\%$).

Analyzing risk factors for RLNP and LOS, such as redo neck surgery, neck dissections, invasive thyroid cancer and intrathoracic or large goiter, none were found to be significant for RLNP.

Twelve patients presented transient postoperative hypoparathyroidism and two had persistent hypoparathyroidism at 6 months. Three patients presented non-surgical postoperative hematoma.

Comparison with loss of signal using I-IONM

Of all 1586 cases performed using I-IONM during the same time period, 64 patients had persistent LOS, with 42 cases presenting with RLNP at POD1 ($42/1485 = 2.8\%$, excluding those who underwent C-IONM), and in 22 of the cases, the DL on POD1 was normal (I-IONM with false positive LOS or early recovery of RLNP).

Analyzing the causes of LOS, there were 26 due to traction mechanism ($26/64 = 40.6\%$), of those 20 had RLNP on POD1. Eleven presented with difficult dissection around the RLN due to adhesions, three had thermic injury to the nerve, one accidental section, two after compression (for bleeding) and the rest was unknown.

RLNP persisted in 13 patients at 3 months, and in one patient at 6 months. Three patients were lost to follow-up.

Discussion

This study presents the first 101 cases performed with C-IONM in neck surgery at the University Hospitals of Geneva. The results show that in 13 patients the events were temporary after a change in surgical approach was performed and prevented a LOS from occurring. In three cases, LOS occurred at electrode placement on the vagal nerve and in one case with an accidental section of the anterior branch of RLN, RLNP persisted after 6 months ($1/101 = 1\%$).

Our findings reveal a direct risk to the vagal nerve by electrode placement in three cases, resulting in a rate of RLNP at POD1 due to the electrode of 2.97% ($=3/101$ cases, or $1.9\% = 3/159$ nerves at risk). The vagal nerve was intact in all three cases, and placement was carefully performed under direct vision, without hematoma. The exact reason for this trauma is unclear, but might be due to perineural bleeding or traction by surgical instruments when exposing the nerve. In our first case (Grave's disease,

#4), the dissection was not particularly difficult but the vagal nerve was positioned posterior and deep to the vessels. In the second case (recurrent PTC, #5), vagal nerve dissection was difficult, because the patient had undergone previous central and lateral lymph node dissection. And in the third case (PTC N1b metastasis, #6), the vagal nerve was dissected full length for the lateral neck dissection. Also, the signal was lost proximally to the electrode and RLNP was transient in all three cases. Only a few such cases have been described in the literature, with Brauckhoff et al. [17] citing two cases of LOS by the electrode in high-risk surgery, one at placement due to perineural bleeding, and one caused by an undetected torsion of the electrode. Mangano et al. [15], in a safety study of 400 vagal nerve dissections and electrode placements, report a mean increase in vagal nerve size after electrode placement of .82 mm and an electrode dislocation rate of 11% per procedure, thus highlighting the importance of selecting the suitable electrode size and using careful, standardized technique of vagal nerve dissection and electrode placement. Every new surgical technique has a learning curve [18]. The three LOS at electrode placement occurred over the 5-year study period, the last case happening after standard dissection of more than 100 vagal nerves. This led us to believe that LOS might be caused by the upward traction of the right-angle dissector used for the 360° nerve dissection necessary for electrode placement. We believe it is mandatory to verify the signal on the vagus nerve above the site of electrode placement (proximal) at the end of surgery, in addition to the C-IONM signal from the electrode, as this is not sufficient to guarantee postoperative RLN function.

With regard to the 13 temporary events ($13/101 = 12.9\%$), the decrease in amplitude of the initial IONM value was seen when medializing the thyroid. The most logical explanation for this phenomenon is the mobilization of the thyroid, which causes the nerve to stretch and leads to a decreased conduction capacity. The perioperative solution in response to C-IONM was to release the traction, which in addition to a change in approach, led to full recovery of all RLN signals. Going further, we could assume that some of the 26 traction injuries in the I-IONM cases could have been avoided if C-IONM had been used, as the traction would have been immediately released before definitive injury occurs. This concurs with findings from other studies that report events to be reversible when prompt corrective measures are applied during surgery [10, 11, 19]. Thus, our study shows that the RLN recovers when appropriate measures are taken.

We had one case of accidental section of the anterior RLN branch while using C-IONM, resulting in an overall permanent RLNP rate of 1% ($1/101$ cases). This RLNP was

Table 2 Cases with intraoperative loss of signal at C-IONM

Case	Age sex	Planned surgery	LOS type	LOS cause	Surgery change	Pathology, TNM	Size/ weight	DL D1	Second DL at follow-up	Follow-up
1	78 m	Total thyroidectomy	2	Traction, on second side	Release of traction No change for approach RLN	FV PTC, T1b (m)	1.2 cm	uRLNP	M1: no uRLNP	
2	44 m	Total thyroidectomy	1	Section of ant. branch RLN, on second side	No change for approach RLN	MNG	96 g	uRLNP	M6: uRLNP	
3	50 f	Lobectomy	1	Thermal lesion	Superior approach to RLN	MNG	92 g	uRLNP	M6: no uRLNP	
4	51 f	Total thyroidectomy	1	LOS at electrode placement on vagus nerve	Distal placement of electrode Stop after 1st side	Grave's disease	62 g	uRLNP	M2: no uRLNP	M2: Surgery 2nd side
5	49 f	Recurrent CND	1	LOS at electrode placement on vagus nerve	Distal placement of electrode	Metastatic LN PTC (3/19)	3.8 cm	uRLNP	M6: no uRLNP	
6	48 f	Total thyroidectomy, LND and CND	1	LOS at electrode placement on vagus nerve	Distal placement of electrode Superior approach to RLN Stop after 1st side	PTC T3 N1b (7/16)	2 cm	uRLNP	D15: no uRLNP	D15: Surgery 2nd side

CND central lymph node dissection, D day, DL direct laryngoscopy, f female, FV PTC follicular variant of papillary thyroid cancer, g gram, LN lymph node, LND lateral lymph node dissection, LOS type loss of signal: type 1 = segmental; type 2 = global, M month, m male, MNG multinodular goiter, RLN recurrent laryngeal nerve, uRLNP unilateral recurrent laryngeal nerve palsy

not prevented by C-IONM and reflects a segmental LOS, which is caused by abrupt trauma, such as thermic, transection or clamping. These segmental lesions usually happen without delay and do not permit the surgeon to take protective action, contrary to the gradual, global LOS due to traction. They are therefore a limitation of C-IONM.

Overall, both permanent and temporary RLNP rates were similar to another study using C-IONM in high-risk neck surgeries [17]; however, they were higher than in other studies, which had no permanent RLNP using C-IONM [10] when compared to I-IONM [11]. Furthermore, no risk factors for RLNP have been found, looking at extent of disease and difficulty of surgery.

Regarding the risk of dislocation of the electrode during surgery, cited by other studies to be between 11 and 30% [11, 15], we feel that the most important factor causing dislocations is the surgical maneuvers used when placing the electrode and during thyroid surgery. Our group

dissects the vagal nerve staying underneath the sternothyroid and sterno-hyoid muscles. Certain groups have used an entirely lateral approach (to the muscles) or even going in between the pre-thyroidal muscles [20] for access. Retracting carefully and laterally the pre-thyroidal muscles seemed to distance the electrode from the surgical field and protect it from dislocations. Usually, the electric wire and its electrode were not touched by the retractors and thus not under traction.

Also, as mentioned before, a two-staged thyroidectomy is performed at our center in case of LOS on the first side [8]. This approach is controversial and not always used by other teams [21, 22], but we prefer this approach in order to avoid the risk of bilateral RLNP, and patients are informed prior to surgery and understand the reasoning. A recent study also suggests that this approach is cost-effective [23].

The present study has limitations. First, the number of patients with C-IONM was limited, as they were only high-

risk cases, depending strictly on the perioperative decision of the surgeon. This introduces a selection bias in the comparison with I-IONM. Second, due to the non-randomized study design, the final conclusion regarding the potential benefit of C-IONM is weakened, and the small number of events did not allow for robust analysis. Last, the use of pre- and postoperative laryngoscopies was not systematic, which limits the interpretation of the study's results regarding RLNP and outcome of C-IONM. To obtain more reliable outcome data regarding C-IONM, future randomized studies are needed.

In conclusion, C-IONM provides a real-time evaluation of the RLN function, allowing for adaptation of surgical maneuvers to prevent RLNP. Because it implies a risk at electrode placement, the risk–benefit ratio of C-IONM should be further evaluated. In our opinion, C-IONM is particularly useful in situations where the RLN is difficult to identify by the standard lateral approach, like in redo surgery, in patients with posterior nodules or with large and/or retrosternal goiters. In these latter situations (large and retrosternal goiters), we favor a superior approach to the RLN (also called the toboggan technique [24]) and find C-IONM very helpful for this technique. Furthermore, when using C-IONM, we believe that a vagal stimulation above the site of electrode placement should be performed at the end of dissection to ensure integrity of the vagal nerve and RLN and to avoid false negative IONM result (a good IONM signal with true RLNP).

Acknowledgements There was no funding for this retrospective study.

Compliance with ethical standards

Conflict of interest All authors declare that they have no conflict of interest.

References

- Rosenthal LH, Benninger MS, Deeb RH (2007) Vocal fold immobility: a longitudinal analysis of etiology over 20 years. *Laryngoscope* 117:1864–1870
- Dralle H, Sekulla C, Haerting J et al (2004) Risk factors of paralysis and functional outcome after recurrent laryngeal nerve monitoring in thyroid surgery. *Surgery* 136:1310–1322
- Rosato L, Avenia N, Bernante P et al (2004) Complications of thyroid surgery: analysis of a multicentric study on 14,934 patients operated on in Italy over 5 years. *World J Surg* 28:271–276
- Bergenfels A, Jansson S, Kristofferson A et al (2008) Complications to thyroid surgery: results as reported in a database from a multicenter audit comprising 3,660 patients. *Langenbecks Arch Surg* 393:667–673
- Higgins TS, Gupta R, Ketcham AS et al (2011) Recurrent laryngeal nerve monitoring versus identification alone on post-thyroidectomy true vocal fold palsy: a meta-analysis. *Laryngoscope* 121:1009–1017
- Yang S, Zhou L, Lu Z et al (2017) Systematic review with meta-analysis of intraoperative neuromonitoring during thyroidectomy. *Int J Surg* 39:104–113
- Barczynski M, Konturek A, Cichon S (2009) Randomized clinical trial of visualization versus neuromonitoring of recurrent laryngeal nerves during thyroidectomy. *Br J Surg* 96:240–246
- Sadowski SM, Soardo P, Leuchter I et al (2013) Systematic use of recurrent laryngeal nerve neuromonitoring changes the operative strategy in planned bilateral thyroidectomy. *Thyroid* 23:329–333
- Randolph GW, Dralle H, International Intraoperative Monitoring Study G et al (2011) Electrophysiologic recurrent laryngeal nerve monitoring during thyroid and parathyroid surgery: international standards guideline statement. *Laryngoscope* 121(Suppl 1): S1–S16
- Schneider R, Sekulla C, Machens A et al (2015) Postoperative vocal fold palsy in patients undergoing thyroid surgery with continuous or intermittent nerve monitoring. *Br J Surg* 102:1380–1387
- Phelan E, Schneider R, Lorenz K et al (2014) Continuous vagal IONM prevents recurrent laryngeal nerve paralysis by revealing initial EMG changes of impending neuropraxic injury: a prospective, multicenter study. *Laryngoscope* 124:1498–1505
- Schneider R, Randolph G, Dionigi G et al (2016) Prospective study of vocal fold function after loss of the neuromonitoring signal in thyroid surgery: The International Neural Monitoring Study Group's POLT study. *Laryngoscope* 126:1260–1266
- Wu CW, Dionigi G, Sun H et al (2014) Intraoperative neuromonitoring for the early detection and prevention of RLN traction injury in thyroid surgery: a porcine model. *Surgery* 155:329–339
- Lamade W, Bechu M, Lauzana E et al (2016) The weepy nerve-different sensitivity of left and right recurrent laryngeal nerves under tensile stress in a porcine model. *Langenbecks Arch Surg* 401:983–990
- Mangano A, Kim HY, Wu CW et al (2016) Continuous intraoperative neuromonitoring in thyroid surgery: Safety analysis of 400 consecutive electrode probe placements with standardized procedures. *Head Neck* 38(Suppl 1):E1568–E1574
- Terris DJ, Chaung K, Duke WS (2015) Continuous Vagal Nerve Monitoring is Dangerous and Should not Routinely be Done During Thyroid Surgery. *World J Surg* 39:2471–2476
- Brauckhoff K, Vik R, Sandvik L et al (2016) Impact of EMG changes in continuous vagal nerve monitoring in high-risk endocrine neck surgery. *World J Surg* 40:672–680
- Schneider R, Przybyl J, Pliquet U et al (2010) A new vagal anchor electrode for real-time monitoring of the recurrent laryngeal nerve. *Am J Surg* 199:507–514
- Schneider R, Sekulla C, Machens A et al (2016) Dynamics of loss and recovery of the nerve monitoring signal during thyroidectomy predict early postoperative vocal fold function. *Head Neck* 38(Suppl 1):E1144–E1151
- Dionigi G, Kim HY, Wu CW et al (2013) Vagus nerve stimulation for standardized monitoring: technical notes for conventional and endoscopic thyroidectomy. *Surg Technol Int* 23:95–103
- Sitges-Serra A, Fontane J, Duenas JP et al (2013) Prospective study on loss of signal on the first side during neuromonitoring of the recurrent laryngeal nerve in total thyroidectomy. *Br J Surg* 100:662–666
- Khamsy L, Constanthin PE, Sadowski SM et al (2015) Loss of neuromonitoring signal during bilateral thyroidectomy: no systematic change in operative strategy according to a survey of the French Association of Endocrine Surgeons (AFCE). *BMC Surg* 15:95

23. Al-Qurayshi Z, Kandil E, Randolph GW (2017) Cost-effectiveness of intraoperative nerve monitoring in avoidance of bilateral recurrent laryngeal nerve injury in patients undergoing total thyroidectomy. *Br J Surg*
24. Scarpa CR, Triponez F (2014) Retrosternal Thyroid Goiter: Recurrent Laryngeal Nerve Dissection According to the Charles Proye's Toboggan Technique. *VideoEndocrinology* 1
25. Edge S, Labow D, Carduci M (2010) *AJCC Cancer Staging Manual*, 7th edn. Springer, New York, NY