



Article scientifique

Article

2003

Published version

Open Access

This is the published version of the publication, made available in accordance with the publisher's policy.

Revised Small Subunit rRNA Analysis Provides Further Evidence that Foraminifera Are Related to Cercozoa

Berney, Cédric; Pawlowski, Jan Wojciech

How to cite

BERNEY, Cédric, PAWLOWSKI, Jan Wojciech. Revised Small Subunit rRNA Analysis Provides Further Evidence that Foraminifera Are Related to Cercozoa. In: Journal of Molecular Evolution, 2003, vol. 57, p. S120–S127. doi: 10.1007/s00239-003-0015-2

This publication URL: <https://archive-ouverte.unige.ch/unige:113627>

Publication DOI: [10.1007/s00239-003-0015-2](https://doi.org/10.1007/s00239-003-0015-2)

Revised Small Subunit rRNA Analysis Provides Further Evidence that Foraminifera Are Related to Cercozoa

Cédric Berney, Jan Pawlowski

Department of Zoology and Animal Biology, University of Geneva, Geneva, Switzerland

Received: 31 July 2002 / Accepted: 11 December 2002

Abstract. There is accumulating evidence that the general shape of the ribosomal DNA-based phylogeny of Eukaryotes is strongly biased by the long-branch attraction phenomenon, leading to an artifactual basal clustering of groups that are probably highly derived. Among these groups, Foraminifera are of particular interest, because their deep phylogenetic position in ribosomal trees contrasts with their Cambrian appearance in the fossil record. A recent actin-based phylogeny of Eukaryotes has proposed that Foraminifera might be closely related to Cercozoa and, thus, branch among the so-called crown of Eukaryotes. Here, we reanalyze the small-subunit ribosomal RNA gene (SSU rDNA) phylogeny by removing all long-branching lineages that could artifactually attract foraminiferan sequences to the base of the tree. Our analyses reveal that Foraminifera branch together with the marine testate filosean *Gromia oviformis* as a sister group to Cercozoa, in agreement with actin phylogeny. Our study confirms the utility of SSU rDNA as a phylogenetic marker of megaevolutionary history, provided that the artifacts due to the heterogeneity of substitution rates in ribosomal genes are circumvented.

Key words: Foraminifera — Cercozoa — Eukaryotes — *Gromia oviformis* — Molecular phy-

logeny — Long-branch attraction — Small-subunit ribosomal DNA

Introduction

Phylogenetic studies based on the small-subunit ribosomal RNA gene (SSU rDNA) have yielded a “classical” view of eukaryotic evolution, in which several assumed primitive, isolated lineages diverged very early, while other groups, supposedly more recent and apparently simultaneously diverging, cluster in the so-called eukaryotic crown (Sogin 1991; Knoll 1992). However, the primitive status of “early” Eukaryotes, as defined by SSU rDNA phylogenies, has been a subject of long debate (see, e.g., Sogin 1997). Protein-based phylogenies are now providing accumulating evidences that the distinction between “early” and “crown” Eukaryotes is no longer accurate, because many—if not all—“early” Eukaryotes are probably highly derived groups, which are artifactually attracted to the base of SSU rDNA trees by the long-branch attraction (LBA) phenomenon (Embley and Hirt 1998; Philippe and Adoutte 1998; Philippe et al. 2000). Most striking is the example of the Microsporidia, which have been clearly shown to be highly derived, parasitic members of the Fungi (Keeling and Doolittle 1996; Hirt et al. 1999; Keeling et al. 2000; see Van de Peer et al. [2000b] for a review).

Because protein data are still lacking for a large part of the known eukaryotic diversity, the phylogenetic position of many groups of protists remains

Correspondence to: Cédric Berney, Université de Genève, Station de Zoologie, 154 route de Malagnou, 1224 Chêne-Bougeries, Genève, Switzerland; email: cedric.berney@zoo.unige.ch

unclear. Among these groups, the Foraminifera are of particular interest, because their evolutionary history can be inferred from their exceptionally well preserved fossil record. Foraminifera are a large and morphologically diverse group of mainly marine, testate protists with typical filamentous, granuloreticulate pseudopodia (Lee et al. 2000). Molecular data from both large- and small-subunit rDNAs have suggested a very ancient origin for the group (Pawlowski et al. 1994, 1996, 1999). But as discussed before (Pawlowski et al. 1996; Sogin 1997), rRNA genes from the Foraminifera are highly divergent. Therefore, a very early branching of the group must be taken suspiciously.

A recent actin-based phylogeny of Eukaryotes showed a close relationship among the Foraminifera, *Cercomonas*, and *Chlorarachnion* (Keeling 2001), suggesting a possible inclusion of Foraminifera within Cercozoa. This group of “crown” Eukaryotes was recently described on the basis of molecular data (Bhattacharya et al. 1995). The Cercozoa consist of a diverse assemblage of organisms, such as *Cercomonas*-like amoeboid flagellates, the euglyphid testate filose amoebae, the plasmodiophorid plant pathogens, and the chlorarachniophyte green amoebae (Cavalier-Smith 1998; see also Cavalier-Smith and Chao 1997; Atkins et al. 2000; Kühn et al. 2000; Bulman et al. 2001; Vickerman et al. 2002). Clear morphological evidence is yet lacking to support the Cercozoa, but a close relationship between *Cercomonas* and *Chlorarachnion* was also recovered on the basis of both alpha- and beta-tubulin phylogenies (Keeling et al. 1998, 1999).

A possibility to avoid LBA artifacts, and to find the true phylogenetic position of an assumed fast-evolving, derived group of Eukaryotes, is to suppress all other possibly fast-evolving taxa from the analyses and, as suggested by Swofford et al. (1996), to reconstruct a tree including only ingroup taxa. These two conditions are necessary in order to avoid the attraction of the tested group by both the sequences from the outgroup and any other fast-evolving taxa. If the group examined is really a primitive, independent lineage of Eukaryotes, then its position in such phylogenetic trees should be left unresolved, with no clear affinity for any group of crown Eukaryotes. But if it is actually a derived member of one of these crown clades, then one can hope that there is still enough phylogenetic signal in its SSU rDNA sequences so that its true phylogenetic position might be retrieved. This approach was used by Van de Peer et al. (2000b), which showed that when considering among-site rate variation, reanalysis of large-subunit rDNA sequences from “crown” Eukaryotes, including two sequences of Microsporidia, supports the placement of the latter within the Fungi. In this study, we use this approach in order to test if there is

any support in SSU rDNA sequences for a cercozoan affiliation of Foraminifera.

Materials and Methods

The complete SSU rDNA sequences of 5 Foraminifera and 49 other Eukaryotes were manually aligned using the Genetic Data Environment software (Larsen et al. 1993), following the secondary structure models proposed by Neefs et al. (1993) and Wuyts et al. (2000). Sequences were chosen so that representatives of all well-defined domains of the “crown” were included in the data set, using the studies by Van de Peer and De Wachter (1997) and Van de Peer et al. (2000a) as references. Eukaryotic lineages that were excluded from our data set comprise all groups (Diplomonadida, Parabasalids, Microsporidia, Euglenozoa, Heterolobosea, Radiolaria, Mycetozoa, Entamoebidae, and Pelobiontida) that are thought to have undergone a process of accelerated rate of evolution and, thus, may act as long branches in phylogenetic analyses (see, e.g., Stiller and Hall 1999; Morin 2000; Philippe and Germot 2000; Bolivar et al. 2001; López-García et al. 2002). Relative rate tests performed with RRTree (Robinson-Rechavi and Huchon 2000) confirmed at both 5 and 1% levels that the representatives of these groups display significantly higher rates of substitution compared to crown species. The names, phylogenetic position, and GenBank accession numbers of the sequences used in this study are indicated in Table 1.

Evolutionary trees were inferred using the neighbor-joining (NJ) method (Saitou and Nei 1987), the maximum-parsimony (MP) method, and the maximum-likelihood (ML) method (Felsenstein 1981). The reliability of internal branches was assessed using the bootstrap method (Felsenstein 1985), with 1000 replicates for NJ analyses, 500 replicates for MP analyses, and 100 replicates for ML analyses. The Phylo_Win program (Galtier et al. 1996) was employed for distance computations and NJ building and bootstrapping, using the HKY85 model of substitution (Hasegawa et al. 1985). MP and ML analyses were performed with PAUP* (Swofford 1998). The most parsimonious trees for each MP bootstrap replicate were determined using a heuristic search procedure with 10 random-addition-sequence replicates and tree bisection-reconnection branch-swapping. The transitions cost was set to twice the transitions cost. For ML analyses, the TrN model of substitution was used (Tamura and Nei 1993), taking into account a proportion of invariable sites, and a gamma-shaped distribution of rates of substitution among sites, with eight categories. All necessary parameters were estimated from the data set using Modeltest (Posada and Crandall 1998). Starting trees of ML searches were obtained via NJ and swapped with the tree bisection-reconnection algorithm.

Results and Discussion

All methods of tree reconstruction yielded the same result: when no other sequences with particularly high rates of substitutions are included in the analyses, the Foraminifera branch together with the marine testate filosean *Gromia oviformis* as a sister group to the Cercozoa (Fig. 1). Bootstrap support values for the clade consisting of Foraminifera + *Gromia* + Cercozoa are good: 88, 94, and 88%, for ML, NJ, and MP analyses, respectively. As the ML tree in Fig. 1 contains only ingroup sequences, it is presented in an unrooted format, with a basal trichotomy separating

Table 1. List of the taxa used in our analyses

Taxonomic position	Species name	GenBank accession no.
Amoebozoa	<i>Acanthamoeba castellanii</i>	M13435
	<i>Echinamoeba exundans</i>	AF293895
	<i>Hartmannella vermiformis</i>	M95168
	<i>Leptomyxa reticulata</i>	AF293898
Opisthokonta	<i>Spizellomyces acuminatus</i>	M59759
	<i>Schizosaccharomyces pombe</i>	X58056
	<i>Amanita muscaria</i>	AF026631
	<i>Nuclearia simplex</i>	AF349566
	<i>Diaphanoeca grandis</i>	L10824
	<i>Leucosolenia</i> sp.	AF100945
	<i>Trichoplax adhaerens</i>	L10828
	<i>Mytilus edulis</i>	L33448
Cryptophyta	<i>Goniomonas truncata</i>	U03072
	<i>Guillardia theta</i>	X57162
Glaucophyta	<i>Cyanophora paradoxa</i>	X68483
Rhodophyta	<i>Flintiella sanguinaria</i>	AF168621
	<i>Composopogon coeruleus</i>	AF087128
	<i>Glaucosphaera vacuolata</i>	AB045583
Viridiplantae	<i>Dunaliella salina</i>	M84320
	<i>Mesostigma viride</i>	AJ250108
	<i>Huperzia phlegmaria</i>	X81964
	<i>Helianthus annuus</i>	AF107577
Alveolata	<i>Blepharisma americanum</i>	M97909
	<i>Oxytricha nova</i>	X03948
	<i>Colpoda inflata</i>	M97908
	<i>Perkinsus marinus</i>	AF126013
Stramenopiles	<i>Crypthecodinium cohnii</i>	M64245
	<i>Prorocentrum concavum</i>	Y16237
	<i>Cafeteria roenbergensis</i>	L27633
	<i>Thraustochytrium multirudimentale</i>	AB022111
	<i>Blastocystis hominis</i>	U51151
	<i>Apodachlya brachynema</i>	AJ238663
	<i>Chrysosolepidomonas dendrolepidota</i>	AF123297
	<i>Ectocarpus siliculosus</i>	L43062
	<i>Coccolithus pelagicum</i>	AJ246261
	<i>Phaeocystis cordata</i>	AF163147
Haptophyta	<i>Pavlova salina</i>	L34669
	<i>Gromia oviformis</i> 1	AJ457811
Cercozoa	<i>Gromia oviformis</i> 2	AJ457812
	<i>Cercomonas longicauda</i>	AF101052
	<i>Euglypha rotunda</i>	X77692
	<i>Pseudodiffugia</i> cf. <i>gracilis</i>	AJ418794
	<i>Nuclearia</i> -like filose amoeba NPor	AF174374
	<i>Massisteria marina</i>	AF174374
	<i>Chlorarachnion reptans</i>	U03477
	<i>Lotharella vacuolata</i>	AF054890
	<i>Plasmodiophora brassicae</i>	U18981
	<i>Spongospora subterranea</i>	AF310899
Foraminifera	<i>Phagomyxa odontellae</i>	AF310904
	<i>Allogromia</i> sp.	X86093
	<i>Astrammia rara</i>	AJ318223
	<i>Reticulomyxa filosa</i>	AJ132367
	<i>Trochammina</i> sp.	X86095
	<i>Bolivina spathulata</i>	AJ318227

Opisthokonta, Amoebozoa, and Bikonta, following a recent suggestion of Stechmann and Cavalier-Smith (2002). The stem branch leading to the Foraminifera

was reduced to half of its actual size to make the rest of the tree clearer, which emphasizes the high divergence of foraminiferan SSU rDNA sequences. However, their inclusion in the analyses does not seem to disrupt the global phylogenetic informativeness of the molecule, because all groups of “crown” Eukaryotes that are now firmly established on the basis of molecular data (see, e.g., Van de Peer et al. 2000a) are recovered with confident bootstrap support.

Given the fact that highly diverging outgroup sequences generally join the longest branch in the ingroup (Wheeler 1990), the position of the Foraminifera in the tree shown in Fig. 1 might be regarded as the result of the artificial clustering of the foraminiferan sequences to one of the most rapidly evolving group of “crown” Eukaryotes, i.e., the Cercozoa. In order to test this hypothesis, we performed additional analyses on two enhanced data sets, incorporating sequences that were primarily rejected by relative rate tests. The first data set included some more diverging members of the Amoebozoa, like Mastigamoebidae, and the second one included some rapidly evolving members of the polycystine Radiolaria. In both cases, the phylogenetic position of the Foraminifera remained unchanged in ML analyses (data not shown). These results show that Foraminifera branch with Cercozoa even in the presence of some faster-evolving lineages, excluding the hypothesis that their grouping together in Fig. 1 was an artifact. Besides, analyses of the first data set showed that the ML topology is even recovered with NJ analyses, provided that enough slowly evolving members of the Amoebozoa are included (data not shown). Furthermore, ML analyses of the second data set interestingly revealed that the Radiolaria might represent a sister group to the clade comprising Foraminifera, *Gromia*, and Cercozoa, as already hypothesized by Burki et al. (2002). In this case, however, NJ analyses failed to recover the sister-group relationship between *Gromia* and the Foraminifera, and the latter were attracted to the fast-evolving polycystine Radiolaria in the NJ tree (data not shown).

The fact that Foraminifera and Cercozoa branch together even in the presence of some faster-evolving lineages suggests that a true phylogenetic signal for this relationship exists in SSU rDNA sequences. Indeed, a screening of the primary structure of the sequences used in our analyses revealed that seven of the eight nucleotide states that specifically define the Cercozoa (i.e., that are absent in all other crown Eukaryotes) are also present in the foraminiferan sequences. Besides, Foraminifera and Cercozoa share the same character states in most other informative but potentially homoplasious positions, i.e., where other groups or subgroups of crown Eukaryotes also display the same nucleotide. Interestingly, these




Fig. 1. Phylogenetic position of the Foraminifera based on the analysis of 54 SSU rDNA sequences from diverse Eukaryotes, using 1117 unambiguously aligned positions. The tree is the ML tree, which is presented in an unrooted format (see text); its log likelihood value is -14,413.669. When no other sequences with high rates of substitution are included in the analyses, the Foraminifera cluster with the Cercozoa, as a sister group to the marine testate

filosean *Gromia oviformis* (shaded area). Numbers at internal nodes are the bootstrap support values for ML, NJ, and MP analyses after 100, 1000, and 500 replicates, respectively; dashes indicate values under 50%. When all three bootstrap support values are inferior to 50%, they are not indicated. All branches are drawn to scale, except the stem-branch leading to the Foraminifera, which was reduced to half its actual size.




Fig. 2. Alignment of helices 24 to 26 of the SSU rDNA sequences of 4 Foraminifera and 16 “crown” Eukaryotes, highlighting some of the specific positions that indicate a close relationship among Cercozoa, Foraminifera, and the testate marine forams *Gromia oviformis* (light gray box). The dark gray boxes show two complementary sequences signatures (GCYG and TRGC) uniting Foraminifera and *G. oviformis*. Dots indicate identity with the consensus sequence, which was defined as the most frequent nucleotide in each position. The limits of each helix, as defined by Wuyts et al. (2000), are indicated below the alignment. Numbers at the top refer to the sequence of *Cercomonas longicauda* (GenBank accession number AF101052).

cercozoan-specific signatures are not completely randomly scattered through the SSU rDNA, but are concentrated mainly in the region comprising helices 24 to 26 (Fig. 2), suggesting local rearrangements of the secondary structure of the molecule. In our opinion, this feature is not likely to be due to random convergence, because in that case one might expect the foraminiferan sequences to show similar convergences toward the specific signatures of any other group of crown Eukaryotes, which is not the case.

The preservation of cercozoan-specific nucleotides in Foraminifera is particularly significant, given the high divergence of the foraminiferan stem lineage, as illustrated by about 130 specific nucleotide sites which define this group in the conserved regions of our alignment. Analyses of partial SSU rDNA sequences from a large sampling of extant foraminiferan lineages, using calibration points obtained from the rich fossil record of the group, revealed that an episodic change of substitution rates occurred in the stem lineage leading to the Foraminifera. According to our calibration, the rate of substitution in the most conserved regions of the SSU rDNA averaged at least 1.0 to 1.65 substitutions/1000 sites/ 10^6 years during the stem lineage evolution of Foraminifera. This rapid burst of the evolutionary rate was followed by a return to a "normal" value of about 0.03 substitutions/1000 sites/ 10^6 years during the subsequent radiation of the different foraminiferan lineages (Pawlowski and Berney, unpublished), comparable to evolutionary rates calculated in other groups of Eukaryotes (see, e.g., Sorhannus 1996).

Our data confirm a recent analysis of actin sequences (Keeling 2001) and are also congruent with unpublished analyses of RNA polymerase II coding sequences (Longet, personal communication). All these studies clearly show that Foraminifera are related to Cercozoa, a relationship that is also strongly supported by polyubiquitin structure (Archibald, personal communication). However, whether Foraminifera are a derived lineage of Cercozoa or their sister group remains disputable. Although the Foraminifera and *Gromia oviformis* appear as sister groups to Cercozoa in the ML tree (Fig. 1), this is not the case in other analyses. In both the NJ tree and one of the two most parsimonious MP trees, the clade comprising Foraminifera and *G. oviformis* formed a sister group to the plasmodiophorid plant pathogens (data not shown), proning the inclusion of both *Gromia* and the Foraminifera within the Cercozoa.

The close relationship between Foraminifera and *G. oviformis* suggested by our analyses (Fig. 1) could argue in favor of a cercozoan origin of Foraminifera. In fact, for a long time *G. oviformis* was considered a member of the Testacea filosa (class Filosea), whose representatives (e.g., *Euglypha*, *Paulinella*) branch within Cercozoa in molecular analyses (see, e.g.,

Cavalier-Smith and Chao 1997; Cavalier-Smith 1998). However, recent analyses of SSU rDNA sequences of *Gromia* (Burki et al. 2002) showed that this genus is not closely related to the Euglyphida (*Euglypha*, *Paulinella*) but, rather, branches near the base of the cercozoan clade. This may suggest that the common ancestor of Foraminifera and *Gromia* evolved from the cercozoan stem lineage before the radiation of the living representatives of the Cercozoa. Further studies of protein-coding genes should allow resolution of this problem, as well as attainment of a clearer idea of the true evolutionary relationships among the different cercozoan lineages.

Our findings bring further support to the idea that the so-called eukaryotic crown is actually an artifactual clustering of slowly evolving sequences, to the exclusion of a few rapidly evolving lineages (Stiller and Hall 1999; Philippe et al. 2000). The relationship between Foraminifera and Cercozoa also suggests that although diversity within the protists remains largely underestimated (see, e.g., Moreira and López-García 2002), most eukaryotic lineages might have evolved from a reduced number of large "super-groups." Finally, our results show that even if recent protein-based analyses have cast strong doubts on the phylogenetic potential of SSU rDNA sequences to recover deep evolutionary relationships (see, e.g., Baldauf et al. 2000; Moreira et al. 2000), SSU rDNA data might still remain a source of valuable phylogenetic information. Although protein data are rapidly accumulating, they are available for a very limited taxonomic sampling of unicellular Eukaryotes. Because of technical constraints, the evolutionary studies of many noncultivable protozoans, such as Foraminifera and Radiolaria, remain largely dependent on ribosomal genes phylogenies, hence their crucial importance in megaevolutionary studies.

Acknowledgments. We would like to thank Colomán de Vargas, Juan Montoya, and three anonymous referees for helpful comments on an early version of our manuscript. This study was supported by Swiss NSF Grant 31-59145.99.

References

- Atkins MS, Teske AP, Anderson OR (2000) A survey of flagellate diversity at four deep-sea hydrothermal vents in the eastern Pacific ocean using structural and molecular approaches. *J Eukaryot Microbiol* 47:400–411
- Baldauf SL, Roger AJ, Wenk-Siefert I, Doolittle WF (2000) A kingdom level phylogeny of Eukaryotes based on combined protein data. *Science* 290:972–977
- Bhattacharya D, Helmchen T, Melkonian M (1995) Molecular evolutionary analyses of nuclear-encoded small subunit ribosomal RNA identify an independent rhizopod lineage containing the Euglyphina and the Chlorarachniophyta. *J Eukaryot Microbiol* 42:65–69
- Bolivar I, Fahrni JF, Smirnov A, Pawlowski J (2001) SSU rRNA-based phylogenetic position of the genera *Amoeba* and *Chaos*

- (Lobosea, Gymnamoebia): The origin of Gymnamoebae revisited. *Mol Biol Evol* 18:2306–2314
- Bulman SR, Kühn SF, Marshall JW, Schnepf E (2001) A phylogenetic analysis of the SSU rRNA from members of the Plasmodiophorida and Phagomyxida. *Protist* 152:43–51
- Burki F, Berney C, Pawlowski J (2002) Phylogenetic position of *Gromia oviformis* Dujardin inferred from nuclear-encoded small subunit ribosomal DNA. *Protist* 153:251–260
- Cavalier-Smith T (1998) A revised six-kingdom system of life. *Biol Rev Camb Philos Soc* 73:203–266
- Cavalier-Smith T, Chao EE (1997) Sarcomonad ribosomal RNA sequences, rhizopod phylogeny, and the origin of euglyphid amoebae. *Arch Protistenkd* 147:227–236
- Embley TM, Hirt RP (1998) Early branching Eukaryotes? *Curr Opin Genet Dev* 8:624–629
- Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* 17:368–376
- Felsenstein J (1985) Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* 39:783–791
- Galtier N, Gouy M, Gautier C (1996) SEAVIEW and PHYLO_WIN: Two graphic tools for sequence alignment and molecular phylogeny. *Comput Appl Biosci* 12:543–548
- Hasegawa M, Kishino H, Yano T (1985) Dating the human-ape split by a molecular clock of mitochondrial DNA. *J Mol Evol* 22:160–174
- Hirt RP, Logsdon Jr JM, Healy B, Dorey MW, Doolittle WF, Embley TM (1999) Microsporidia are related to Fungi: Evidence from the largest subunit of RNA polymerase II and other proteins. *Proc Natl Acad Sci USA* 96:580–585
- Keeling PJ (2001) Foraminifera and Cercozoa are related in actin phylogeny: two orphans find a home? *Mol Biol Evol* 18:1551–1557
- Keeling PJ, Doolittle WF (1996) Alpha-tubulin from early-diverging eukaryotic lineages and the evolution of the tubulin family. *Mol Biol Evol* 13:1297–1305
- Keeling PJ, Deane JA, McFadden GI (1998) The phylogenetic position of alpha- and beta-tubulins from the *Chlorarachnion* host and *Cercomonas* (Cercozoa). *J Eukaryot Microbiol* 45:561–570
- Keeling PJ, Deane JA, Hink-Schauer C, Douglas SE, Maier UG, McFadden GI (1999) The secondary endosymbiont of the Cryptomonad *Guillardia theta* contains alpha-, beta-, and gamma-tubulin genes. *Mol Biol Evol* 16:1308–1313
- Keeling PJ, Luker MA, Palmer JD (2000) Evidence from beta-tubulin phylogeny that Microsporidia evolved from within the Fungi. *Mol Biol Evol* 17:23–31
- Knoll AH (1992) The early evolution of Eukaryotes: a geological perspective. *Science* 256:622–627
- Kühn S, Lange M, Medlin LK (2000) Phylogenetic position of *Cryothecomonas* inferred from nuclear-encoded small subunit ribosomal RNA. *Protist* 151:337–345
- Larsen N, Olsen GJ, Maidak BL, McCaughey MJ, Overbeek R, Macke TJ, Marsh TL, Woese CR (1993) The ribosomal database project. *Nucleic Acids Res* 21:3021–3023
- Lee JJ, Pawlowski J, Debenay JP, Whittaker J, Banner F, Gooday AJ, Tendal O, Haynes J, Faber WW (2000) Phylum Granuloreticulosa. In: Lee JJ, Leedale GF, Bradbury P (eds) *The illustrated guide to the Protozoa*, 2nd ed. Allen Press, Lawrence, KS, pp 872–951
- López-García P, Rodríguez-Valera F, Moreira D (2002) Toward the monophyly of Haeckel's Radiolaria: 18S rRNA environmental data support the sisterhood of Polycystinea and Acantharea. *Mol Biol Evol* 19:118–121
- Moreira D, López-García P (2002) The molecular ecology of microbial Eukaryotes unveils a hidden world. *Trends Microbiol* 10:31–38
- Moreira D, Le Guyader H, Philippe H (2000) The origin of red algae: Implications for the evolution of chloroplasts. *Nature* 405:69–72
- Morin L (2000) Long branch attraction effects and the status of “basal Eukaryotes”: Phylogeny and structural analysis of the ribosomal RNA gene cluster of the free-living Diplomonad *Trepomonas agilis*. *J Eukaryot Microbiol* 47:167–177
- Neefs JM, Van de Peer Y, De Rijk P, Chapelle S, De Wachter R (1993) Compilation of small subunit RNA structures. *Nucleic Acids Res* 21:3025–3049
- Pawlowski J, Bolívar I, Guiard-Maffia J, Gouy M (1994) Phylogenetic position of Foraminifera inferred from LSU rDNA gene sequences. *Mol Biol Evol* 11:929–938
- Pawlowski J, Bolívar I, Fahrni JF, Cavalier-Smith T, Gouy M (1996) Early origin of Foraminifera suggested by SSU rDNA gene sequences. *Mol Biol Evol* 13:445–450
- Pawlowski J, Bolívar I, Fahrni JF, De Vargas C (1999) Molecular evidence that *Reticulomyxa filosa* is a freshwater naked Foraminifera. *J Eukaryot Microbiol* 46:612–617
- Philippe H, Adoutte A (1998) The molecular phylogeny of Eukaryota: Solid facts and uncertainties. In: Coombs G, Vickermann K, Sleight M, Warren A (eds) *Evolutionary relationships among Protozoa*. Chapman & Hall, London, pp 25–56
- Philippe H, Germot A (2000) Phylogeny of Eukaryotes based on ribosomal RNA: Long-branch attraction and models of sequence evolution. *Mol Biol Evol* 17:830–834
- Philippe H, Lopez P, Brinkmann H, Budin K, Germot A, Laurent J, Moreira D, Müller M, Le Guyader H (2000) Early branching or fast evolving Eukaryotes? An answer based on slowly evolving positions. *Proc R Soc Lond Ser B* 267:1213–1222
- Posada D, Crandall KA (1998) Modeltest: Testing the model of DNA substitution. *Bioinformatics* 14:817–818
- Robinson-Rechavi M, Huchon D (2000) RRTree: Relative-rate tests between groups of sequences on a phylogenetic tree. *Bioinformatics* 16:296–297
- Saitou N, Nei M (1987) The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Sogin ML (1991) Early evolution and the origin of Eukaryotes. *Curr Opin Genet Dev* 1:457–463
- Sogin ML (1997) History assignment: when was the mitochondrion founded? *Curr Opin Genet Dev* 7:792–799
- Sorhannus U (1996) Higher ribosomal RNA substitution rates in Bacillariophyta and Dasycladales than in Mollusca, Echinodermata and Actinistia-Tetrapoda. *Mol Biol Evol* 13:1032–1038
- Stechmann A, Cavalier-Smith T (2002) Rooting the Eukaryote tree by using a derived gene fusion. *Science* 297:89–91
- Stiller JW, Hall BD (1999) Long-branch attraction and the rDNA model of early eukaryotic evolution. *Mol Biol Evol* 16:1270–1279
- Swofford DL (1998) PAUP*, phylogenetic analysis using parsimony (*and other methods). Sinauer Associates, Sunderland, MA
- Swofford DL, Olsen GJ, Waddell PJ, Hillis DM (1996) Phylogenetic Inference. In: Hillis DM, Moritz C, Mable BK (eds) *Molecular systematics*. Sinauer, Sunderland, MA, pp 407–514
- Tamura K, Nei M (1993) Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol* 10:512–526
- Van de Peer Y, De Wachter R (1997) Evolutionary relationships among the eukaryotic crown taxa taking into account site-to-site rate variation in 18S rRNA. *J Mol Evol* 45:619–630
- Van de Peer Y, Baldauf SL, Doolittle WF, Meyer A (2000a) An updated and comprehensive rDNA phylogeny of (crown)

- Eukaryotes based on rate-calibrated evolutionary distances. *J Mol Evol* 51:565–576
- Van de Peer Y, Ben Ali A, Meyer A (2000b) Microsporidia: Accumulating evidence that a group of amitochondriate and suspectedly primitive Eukaryotes are just curious Fungi. *Gene* 246:1–8
- Vickerman K, Le Ray D, Hoef-Emden K, De Jonckheere J(2002) The soil flagellate *Proleptomonas faecicola*: Cell organisation and phylogeny suggest that the only described free-living Trypanosomatid is not a Kinetoplastid but has Cercomonad affinities. *Protist* 153:9–24
- Wheeler WC (1990) Nucleic acid sequence phylogeny and random outgroups. *Cladistics* 6:363–367
- Wuyts J, De Rijk P, Van de Peer Y, Pison G, Rousseeuw P, De Wachter R (2000) Comparative analysis of more than 3000 sequences reveals the existence of two pseudoknots in area V4 of eukaryotic small subunit ribosomal RNA. *Nucleic Acids Res* 28:4698–4708