



Article
scientifique

Revue de la
littérature

2024

Published
version

Open
Access

This is the published version of the publication, made available in accordance with the publisher's policy.

Exploring host–pathogen interactions in the *Dictyostelium*
discoideum–*Mycobacterium marinum* infection model of tuberculosis

Guallar Garrido, Sandra; Soldati, Thierry

How to cite

GUALLAR GARRIDO, Sandra, SOLDATI, Thierry. Exploring host–pathogen interactions in the *Dictyostelium discoideum*–*Mycobacterium marinum* infection model of tuberculosis. In: Disease models & mechanisms, 2024, vol. 17, n° 7, p. dmm050698. doi: 10.1242/dmm.050698

This publication URL: <https://archive-ouverte.unige.ch/unige:179081>

Publication DOI: [10.1242/dmm.050698](https://doi.org/10.1242/dmm.050698)

REVIEW

Exploring host–pathogen interactions in the *Dictyostelium discoideum*–*Mycobacterium marinum* infection model of tuberculosis

Sandra Guallar-Garrido* and Thierry Soldati

ABSTRACT

Mycobacterium tuberculosis is a pathogenic mycobacterium that causes tuberculosis. Tuberculosis is a significant global health concern that poses numerous clinical challenges, particularly in terms of finding effective treatments for patients. Throughout evolution, host immune cells have developed cell-autonomous defence strategies to restrain and eliminate mycobacteria. Concurrently, mycobacteria have evolved an array of virulence factors to counteract these host defences, resulting in a dynamic interaction between host and pathogen. Here, we review recent findings, including those arising from the use of the amoeba *Dictyostelium discoideum* as a model to investigate key mycobacterial infection pathways. *D. discoideum* serves as a scalable and genetically tractable model for human phagocytes, providing valuable insights into the intricate mechanisms of host–pathogen interactions. We also highlight certain similarities between *M. tuberculosis* and *Mycobacterium marinum*, and the use of *M. marinum* to more safely investigate mycobacteria in *D. discoideum*.

KEY WORDS: Autophagy, *Dictyostelium*, ESCRT, *Mycobacteria*, Phagocytes

Introduction

Mycobacterium tuberculosis (Mtb) causes tuberculosis (TB), a disease responsible for approximately 10 million cases and approximately 1.5 million annual deaths worldwide (WHO, 2023). Indeed, in 2022, TB ranked as the second most deadly infectious disease globally, following COVID-19 and surpassing HIV/AIDS (WHO, 2023). Mtb belongs to the genus *Mycobacterium*, which also includes a substantial group of nontuberculous mycobacteria of medical significance, such as *Mycobacterium marinum* (Mm), *Mycobacterium leprae*, *Mycobacterium abscessus* or *Mycobacterium avium* complex (reviewed by Sharma and Upadhyay, 2020).

TB presents a complex and persistent clinical challenge across multiple stages of the disease. Lung macrophages are initially infected by Mtb, which then serve as the primary replication site, significantly contributing to bacterial dissemination. Macrophages typically engulf, kill and digest pathogens via phagocytosis (see Glossary, Box 1) and xenophagy (Box 1) (Fig. 1). However, certain bacteria, including Mtb and Mm, adeptly manipulate phagocytic cells, creating permissive environments for their growth and

dissemination (Cardenal-Munoz et al., 2018). To better understand and to facilitate TB research, researchers study mycobacteria that are less harmful to humans, including Mm, which shares virulence factors with Mtb (Tobin and Ramakrishnan, 2008).

The complex interplay between mycobacteria and cell-autonomous defence mechanisms has been extensively studied in a variety of model systems, including in amoebae (Cardenal-Munoz et al., 2018), zebrafish (Ramakrishnan, 2013), *Drosophila* (Marshall and Dionne, 2022), mice (Li and Li, 2023), human primary bone marrow-derived macrophages (BMDMs) (Podinovskaia et al., 2013) and human induced pluripotent stem cell-derived macrophages (iPSDMs) (Bernard et al., 2021). Cell models, in particular, are advancing our understanding of the molecular, cellular and dynamic aspects of the interactions that occur between host cells and mycobacteria, such as Mm and Mtb, which we discuss here.

We also review here the use of the amoeba *Dictyostelium discoideum* (Dd) as a model to explore cell-autonomous defence mechanisms that occur in response to mycobacterial infections. Dd shares high levels of evolutionary conservation of host defence mechanisms with mammalian cells and, as such, has proven to be a powerful model for studying host–pathogen interactions and for identifying metabolic pathways relevant to macrophages, as detailed throughout the Review (Cardenal-Munoz et al., 2018; Dunn et al., 2018).

Mtb and Mm share conserved virulence strategies

Using Mm as an Mtb research model entails benefits, such as genetic resemblance and reduced risk to laboratory staff, but also several constraints, which are discussed below. Mm is a close genetic relative of Mtb, sharing significant genomic similarity (Stinear et al., 2008) and essential genes (Lefrançois et al., 2024). At the proteome level, Mtb and Mm share ~3000 orthologous proteins, displaying an average amino acid identity of 85% (Stinear et al., 2008).

Mm causes skin granulomatous infections in humans and TB-like infections in poikilotherms, such as frogs and fish, whereas Mtb infects humans, causing TB (Davis et al., 2002). Consequently, Mm presents itself as an ideal organism that could be used to explore the mechanisms of Mtb virulence in a safer and more efficient manner given the stringent safety measures that are required when conducting research using Mtb. However, evolutionary divergence between both mycobacteria species may lead to different host–pathogen interactions, immune evasion mechanisms or even antimicrobial drug responses, implying that findings in Mm might not always directly translate to Mtb.

The infection process orchestrated by Mtb and Mm in both macrophages and Dd is highly similar (Fig. 1). Both Mtb and Mm employ similar strategies to evade host cell defence mechanisms. These include the release of proteinaceous and lipidic virulence factors and the creation of a mycobacteria-containing vacuole (MCV, Box 1)

Department of Biochemistry, Faculty of Science, University of Geneva, 30 quai Ernest-Ansermet, Science II, 1211 Geneva-4, Switzerland.

*Author for correspondence (sandra.guallargarrido@unige.ch)

DOI: S.G.-G., 0000-0002-9666-5757; T.S., 0000-0002-2056-7931

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

Box 1. Glossary

Autophagy: a conserved intracellular degradation pathway that captures damaged cytoplasmic materials or cytosolic pathogens within an autophagosome – a double-membrane organelle, which fuses with lysosomes, forming a degradative environment known as the autolysosome, in which targeted material undergoes digestion.

DNA extracellular traps: structures released by immune cells, consisting of chromatin and antimicrobial proteins, to ensnare and neutralize pathogens during infections.

Endosomal sorting complex required for transport (ESCRT): an evolutionarily conserved multi-protein complex involved in membrane dynamics and repair that consists of ESCRT-0, ESCRT-I, ESCRT-II, ESCRT-III, Vps4 (AAA-ATPase) and accessory proteins such as ALIX.

Flotillins: membrane proteins associated with lipid rafts in cell membranes, playing roles in cellular processes such as signal transduction, endocytosis and membrane trafficking.

Galectins: soluble β -galactoside-binding receptors that sense self and non-self carbohydrates. Fifteen mammalian galectins act extracellularly and intracellularly.

Guanylate-binding proteins (GBPs): proteins belonging to an interferon γ -inducible subfamily of guanosine triphosphatases (GTPases) that are produced in several cell types, including macrophages, and play a role in the immune response.

Inflammasome system: a multiprotein complex that detects pathogen-associated or danger signals, triggering inflammation and the release of pro-inflammatory cytokines, crucial for immune responses.

iNOS: inducible nitric oxide synthase, an enzyme involved in the production of the antimycobacterial molecule nitric oxide (NO) from L-arginine.

Kil1: a Golgi sulfotransferase involved in the maturation of lysosomal enzymes.

Kil2: a putative magnesium pump involved in the inflammasome system.

Lysozymes: antimicrobial enzymes, including glycosidases, proteases, nucleases, lipases, phosphatases and sulfatases.

Mycobacteria-containing vacuole (MCV): a specialized compartment within host cells where mycobacteria reside after manipulating phagosomes, allowing them to evade host defences and manipulate cellular processes for survival and replication.

Necroptosis: a programmed form of cell death characterized by plasma membrane rupture and release of cellular contents, triggered by specific signalling pathways.

Pathogen-associated molecular patterns (PAMPs): conserved molecules found in pathogens, recognized by pattern recognition receptors, initiating immune responses.

Phagosome: cellular compartment formed by the engulfment of particles or microorganisms, serving to degrade and digest the ingested material via fusion with lysosomes.

Phagocytosis: cellular process in which specialized cells engulf and internalize pathogens or cellular debris into membrane-bound vesicles called phagosomes, facilitating their degradation and clearance.

Pyroptosis: a highly inflammatory form of programmed cell death initiated by inflammasome activation, leading to membrane rupture and release of pro-inflammatory cytokines.

Siderophores: molecules secreted by microorganisms to scavenge iron from the environment.

Stress granules: aggregations of non-translated messenger ribonucleo-proteins and diverse proteins, often regarded as membraneless organelles that can associate with membranes.

Target of rapamycin complex 1 (TORC1): a nutrient-sensitive kinase complex that modulates cellular responses based on nutrient availability.


Ubiquitination: a post-translational modification in which ubiquitin binds covalently to lysine residues in target proteins. Ubiquitination can trigger various cell responses, including protein degradation via the proteasome, autophagy and innate immune signalling.

Xenophagy: selective autophagy that targets cytosolic bacteria by forming autophagosomes, encapsulating the bacteria, and facilitating their eventual delivery to lysosomes.


to bypass phagolysosome maturation and to escape to the host cytosol. Notably, both species harbour the type VII secretion system ESX-1, which is encoded by the ‘region of difference 1’ (RD1) locus and is responsible for secreting the EsxA–EsxB dimer that is crucial for damaging the MCV (Osman et al., 2022; Smith et al., 2008). MCV damage favours escape of Mm and Mtb to the cytosol and release of Mtb DNA into the cytosol, prompting the host to produce type I interferons (Watson et al., 2015). It is worth noting that the attenuated *Mycobacterium bovis* bacillus Calmette–Guérin (BCG) strain lacks RD1 and is used as a vaccine against Mtb (Conrad et al., 2017; Schnettger et al., 2017). Moreover, Mtb and Mm in which RD1 is deleted (Δ RD1) exhibit restricted growth in amoebae (Lopez-Jimenez et al., 2018), mouse or human macrophages (Lewis et al., 2003), iPSDMs (Bernard et al., 2021) and human monocyte-derived macrophages (hMDMs) (Welin et al., 2011). Additionally, both Mm and Mtb possess the ESX-3 system, which secretes small virulence factors such as the dimer EsxG–EsxH, which can interact with the host ‘endosomal sorting complex required for transport’ (ESCRT) complex (Box 1) (Saelens et al., 2022), and the bacterial ESX-5 system (Abdallah et al., 2008). Moreover, Mtb and Mm contain non-essential secretion systems, such as SecA2 (Ge et al., 2022; Serene et al., 2024; van der Woude et al., 2014; Zulauf et al., 2018). It is important to note that although the infectious processes of both mycobacteria are highly similar, there are notable differences. Mm infection thrives at 25°C, contrasting with Mtb preference for 37°C, aligning with their niche-specific adaptation.

Mycobacterial species are characterized by their complex cell walls, which consist of a core layer made up of peptidoglycan, arabinogalactan and specific mycolic acids, the composition of which varies depending on the species (Chiaradia et al., 2017). Peptidoglycan and mycolic acids are significant pathogen-associated molecular patterns (PAMPs, Box 1) (Hossain and Norazmi, 2013); yet, limited research has been conducted on arabinogalactan due to the unavailability of arabinogalactan-deficient mycobacteria (Toyonaga et al., 2016). Recent studies have used chemically synthesized arabinogalactan to demonstrate its role as a virulence factor that interacts with host galectin-9 (LGALS9) (Box 1), exacerbating mycobacterial infections in both Mtb-infected severe combined immunodeficient (SCID) mice and in Mm-infected zebrafish (Wu et al., 2021). Mycobacteria species also produce various other lipids (Guallar-Garrido et al., 2022, 2021). Host cells recognize each lipid through specific receptors (Fig. 2), as reviewed recently by Zihad et al. (2023). For instance, trehalose 6,6'-dimycolate (TDM), present in all known mycobacteria species, can be recognized by host C-type lectin or scavenger receptors, triggering an inflammatory response (Ishikawa et al., 2009) and inhibiting phagosome (Box 1) maturation (Patin et al., 2017; Spargo et al., 1991). Lipoarabinomannan (LAM) produced by species such as Mtb or Mm is recognized by host mannose and DC-SIGN (also known as CD209) receptors, exerting an anti-inflammatory effect (Maeda et al., 2003). Additionally, LAM produced by Mtb can insert into host cell membrane rafts, modifying kinase activity and impeding phagosome maturation (Welin et al., 2008). Sulfoglycolipid-1 (SL-1), which is produced only by pathogenic mycobacteria, remodels host cell membranes, impacting their fluidity and modifying autophagy (Box 1) (Gilmore et al., 2012; Bah et al., 2020). Furthermore, phthiocerol dimycocerosates (PDIMs) insert into host membranes, modifying cholesterol-enriched domains and affecting pathogenesis in zebrafish (Cambier et al., 2020), host interaction in hMDMs (Augenstein et al., 2017), autophagy in human macrophages (Bah et al., 2020) and antibiotic resistance in nutrient-limited conditions *in vitro* (Block et al., 2023).

A Mammalian phagocyte



B Dictyostelium discoideum



Disease Models & Mechanisms

Fig. 1. See next page for legend.

Additionally, mycobacteria can synthesize intracytosolic lipid inclusions that serve as energy reserves (Foulon et al., 2022), even in the absence of external stressors (Campo-Perez et al., 2022; Fines et al., 2023 preprint; Barisch and Soldati, 2017). [Table 1](#)

summarizes the virulence factors of Mtb and Mm, showcasing their fundamental structural and component similarities for efficient pathogenesis and host interactions, despite inhabiting distinct ecological niches (Tobin and Ramakrishnan, 2008).

Fig. 1. Infection cycle of a pathogenic mycobacterium in mammalian phagocytes and in the amoeba *Dictyostelium discoideum*. Schematics of the phagocytosis of pathogenic *Mycobacterium tuberculosis* (Mtb) in a mammalian phagocyte (A) and *Mycobacterium marinum* (Mm) in the amoeba *Dictyostelium discoideum* (Dd) (B). Mycobacteria are recognized by host pathogen recognition receptors (PRRs) on the cell surface, then the host cells internalize the pathogens via phagocytic cups, which are supported by an F-actin scaffold and actin-related protein (ARP) 2/3 complexes. Early phagosomes undergo maturation, transitioning from Ras-related proteins 5 (Rab5) to 7 (Rab7), acquiring vacuolar ATPase (vATPase) to decrease luminal pH, lysosomal enzymes and lysosome-associated membrane proteins (LAMP/LmpA) through endosome fusion, and activating the NADPH oxidase (NOX) complex. In mammalian macrophages and Dd, ingested mycobacteria impede phagosome maturation by forming a mycobacteria-containing vacuole (MCV), within which they proliferate (Barisch et al., 2015; Cardenal-Muñoz et al., 2017) and induce membrane damage via virulence factors that can take advantage of host membrane microdomains, such as those containing flotillins/vacuolins. Alternatively, mycobacteria can undergo early exocytosis before phagosome maturation or replication within the MCV. The host endosomal sorting complex required for transport (ESCRT) machinery and autophagy can repair the damaged MCV membrane, and host factors, such as guanylate-binding protein (GBP), bind the MCV to limit mycobacterial growth. This ultimately leads to mycobacterial degradation upon lysosome fusion with the MCV. However, a dynamic state can be created that alternates between mycobacterial damage and host repair (Lopez-Jimenez et al., 2018; Mehra et al., 2013). Extensive damage to the MCV membrane can recruit the damage sensors galectin-3 (Gal3) in mammalian cells and discoidins (Dsc) in Dd, and can lead to cytosolic access for mycobacteria, facilitating their ubiquitination by E3 ubiquitin ligases (E3-Ub), recapture and subsequent degradation through xenophagy (Cardenal-Muñoz et al., 2017; Gutierrez and Enninga, 2022). Mycobacteria may win over host cell defences, resulting in cytosolic replication and dissemination via exocytosis, host cell lytic/non-lytic mechanisms, or by ejection in the case of Dd (Bo et al., 2023; Gerstenmaier et al., 2024, 2015; Hagedorn et al., 2009). These processes can lead to the formation of extracellular aggregates, which are then phagocytosed by newly recruited Dd (Hagedorn et al., 2009) and macrophages, significantly contributing to bacterial replication, growth, and dissemination (Toniolo et al., 2023). Abbreviations: Atg8, autophagy-related protein 8; LC3, microtubule-associated proteins 1A/1B light chain 3B (MAP1LC3B); Ub, ubiquitinated proteins.

Overall, the utilization of various infection models has been crucial in elucidating the behaviour and virulence factors of mycobacteria. Specifically, Dd provides experimental advantages to study host–pathogen interactions owing to its phagocytic capabilities and conserved innate immune pathways, as elaborated in the subsequent section.

Dd – a versatile phagocyte

Dd, a member of the Amoebozoa phylum, diverged from fungi and animals shortly after the separation of the phylum from plants (Eichinger et al., 2005). Over the past half-century, this soil amoeba has become a versatile model for studying the molecular and cellular mechanisms of cell-autonomous defence mechanisms. It has a haploid genome facilitating its genetic tractability, and low mutation rates compared to other eukaryotes (Gill and Chain, 2023). Dd is especially suited for investigating chemotaxis, cell motility, cell–cell interactions, phagocytosis, cell-autonomous immune defences and lipid-related host–pathogen interactions (Du et al., 2013), and for screening anti-infective compounds (Hanna et al., 2021).

Importantly, Dd utilizes diverse antibacterial mechanisms similar to those used by human phagocytes (Crespo-Yanez et al., 2023), making it a valuable tool for studying a variety of human pathogens, including mycobacteria (Butler et al., 2020; Lefrançois et al., 2024), *Legionella pneumophila*, *Vibrio cholera*, *Francisella noatunensis*, *Pseudomonas aeruginosa* and *Salmonella enterica*, as well as

yeasts and fungi like *Cryptococcus neoformans* and *Aspergillus fumigatus*, as reviewed by Cardenal-Munoz et al. (2018) and Dunn et al. (2018). It has also played a crucial role in screening anti-bacterial compounds against multiple microorganisms, including *Klebsiella pneumoniae* (Ifrid et al., 2022) and Mm (Hanna et al., 2021). Its use as a model host has provided insights into the impact of virulence factors produced by intracellular pathogenic mycobacteria, contributing to our understanding of host–pathogen interactions (Cardenal-Muñoz et al., 2017; Hüsler et al., 2023; Swart et al., 2018).

Beyond their immune functions, Dd cells display altruistic social behaviours. Upon starvation, the amoebae transition from a single-cell state to a multicellular slug that migrates photostatically and thermostatically to the soil surface to form a fruiting body (Kin et al., 2022). The stalk is formed of altruistically dying cells supporting a mass of spores that will later be released and germinate to initiate a new cycle (Bretschneider et al., 2016; Kin et al., 2022). Sentinel cells, which make up <1% of the multicellular slug, play a key altruistic role in protecting the slug from infection by releasing mitochondrially derived DNA extracellular traps (Box 1) to combat bacterial infections (Brock et al., 2016; Chen et al., 2007; Zhang et al., 2016b). Moreover, Dd can exclude pathogen-infected cells from early stages of multicellular development (mounds) but tolerates others, suggesting a potential microbiota-like role for some bacteria (Brock et al., 2018; Farinholt et al., 2019; Haselkorn et al., 2019; López-Jiménez et al., 2019 preprint; Nicolussi et al., 2018).

Protocols have been developed to analyse host–pathogen interactions with Dd (Arafah et al., 2013; Barisch et al., 2015), including analysis of the infection course at the single-cell level (Mottet et al., 2021), isolation and proteomic characterization of MCVs (Guého et al., 2023 preprint) and other bacteria-containing vacuoles (Manske et al., 2018; Schmolders et al., 2017), and assays for gene expression in both the host and pathogen at various infection stages (Kjellin et al., 2019; Lefrançois et al., 2023).

Although Dd shares similarities with animal macrophages, there are notable differences. As a single-celled organism, Dd amoebae possess only innate immune defences, with phagocytic receptors for various ligands and other cell-autonomous pathways (Dunn et al., 2018), as depicted in Fig. 2. Notably, Dd lacks a complex inflammasome system (Box 1) responsible for pro-inflammatory cytokine secretion (Cosson and Soldati, 2008), suggesting that Dd populations will behave differently towards (myco)bacterial infections compared to mammalian host cells. Moreover, as a multicellular organism, Mm-infected Dd slugs do not develop granulomas or TB-like disease (López-Jiménez et al., 2019 preprint), as observed in other model organisms such as zebrafish or mice, precluding the study of granuloma formation and mycobacteria dissemination. Consequently, the evolutionary distance between amoebae and mammalian cells means that findings from Dd studies may not always directly translate to human diseases.

With its strengths and limitations, Dd remains valuable and is arguably the simplest model for studying mycobacteria pathogenesis and host interactions, particularly at the cellular and molecular levels, as detailed in the following sections.

First host–pathogen contact

The initial interaction between mycobacteria and host, together with subsequent events within the host cell, greatly shapes the infection progression. In this section, we compare and contrast key insights into the initial events that take place between mycobacteria and Dd or host macrophages.




Fig. 2. Key receptors that recognize and uptake mycobacteria in mammalian phagocytes and *Dictyostelium discoideum*. (A) Phagocytes, including macrophages, dendritic cells and neutrophils, of the innate immune system can detect pathogens via pathogen recognition receptors (PRRs). Among the most relevant PRRs in mammalian cells during *Mycobacterium tuberculosis* (Mtb) infection are: Toll-like receptors (TLRs) formed by leucine-rich repeats (LRRs) and Toll/interleukin-1 receptor (TIR) domains; C-type lectins [dectin-1 and mannose receptor (MR)]; class B scavenger receptor family (CD36 and LIMP-2); the integrin CR3, which can be found at the plasma membrane and in endosomal membranes; and the cytosolic nucleotide-binding oligomerization domain receptors (NODs) (Aqdas et al., 2021; Goyal et al., 2016; Gunther and Seyfert, 2018). There are several types of TLRs in mammalian phagocytes. (B) In terms of direct orthologs of these PRRs in *Dictyostelium discoideum* (Dd), only class B scavenger receptors and cytosolic proteins containing the TIR domain have been identified and studied in the context of *Mycobacterium marinum* (Mm) infection (Chen et al., 2007; Sattler et al., 2018). Dd harbours lysosomal membrane proteins A (LmpA) and B (LmpB) that serve as scavenger receptors, akin to LIMP-2 and CD36 in mammals, respectively. Moreover, Dd expresses two cytosolic proteins containing the TIR domain, TirA and TirB (Li et al., 2009). TirA is crucial for effective phagocytosis of Gram-negative bacteria (Chen et al., 2007; Zhang et al., 2016b). Dd also has 45 LRR transmembrane proteins without a cytosolic TIR domain, which have been described but not further studied. Dd also possesses three orthologs of C-type lectin receptors (MR), but their functions necessitate further examination (Cosson and Soldati, 2008). Dd also expresses 'similar to integrin- β A' (SibA), which shares characteristics with mammalian CR3 and is involved in adhesion and phagocytosis. Dd features additional receptors among which Far1, a Venus-trap G protein-coupled receptor (GPCR), is implicated in binding bacterial pathogen-associated molecular patterns (PAMPs) (Xu et al., 2021).

Receptors involved in mycobacteria recognition and uptake

Among the most relevant pathogen recognition receptors in mammalian cells during Mtb infection are Toll-like receptors (TLRs), C-type lectins, class B scavenger receptors and the cytosolic nucleotide-binding oligomerization domain receptors (NODs) (Fig. 2) (Aqdas et al., 2021; Goyal et al., 2016; Gunther and Seyfert, 2018). In terms of direct orthologs of these in Dd (Fig. 2), only class B scavenger receptors and cytosolic proteins containing the toll/interleukin 1 receptor (TIR) domain have been identified and studied (Chen et al., 2007; Sattler et al., 2018).

TLRs, located at the plasma membrane (e.g. TLR1, TLR2, TLR4 and TLR6) or within intracellular endosomal compartments (TLR8 and TLR9), trigger signalling pathways upon ligand interaction. When the leucine-rich repeats (LRRs) of TLRs interact with their respective ligands, the cytoplasmic TIR domain triggers signalling to recruit adaptor proteins (Varshney et al., 2022). Dd lacks TLR orthologs, but it does have 45 LRR transmembrane proteins without a cytosolic TIR domain, which have been described but not further studied (Cosson and Soldati, 2008). However, there are two cytosolic proteins in Dd that contain TIR domains, TirA and TirB (Li et al., 2009). TirA expression is upregulated after *L. pneumophila* infection and is essential for the efficient phagocytosis of Gram-negative bacteria (Chen et al., 2007; Zhang et al., 2016b). However, little is known during mycobacterial infection.

In macrophages, C-type lectin receptors, such as mannose receptors or dectin-1 (also known as CLEC7A), are involved in

recognizing specific carbohydrates, such as TDM, and β -glucans present in the mycobacterial cell wall, respectively. They initiate the host immune response and mannose receptors also facilitate mycobacteria internalization (Stamm et al., 2015). In Dd, three orthologs of these receptors exist, although their roles require further study (Cosson and Soldati, 2008). In mammals, three different scavenger receptors are crucial during Mtb infection: CD36, MARCO and class A scavenger receptors (SRA) (Stamm et al., 2015). Dd does not have MARCO or SRA, but three class B scavenger receptors are present. LmpA/LmpC and LmpB are functional counterparts of LIMP-2 (also known as SCARB2) and CD36 in mammals, respectively. LmpB is primarily located in lipid rafts at the plasma membrane and early phagosomes, and its absence is linked to reduced mycobacteria uptake. LmpA is predominantly present in endosomes and lysosomes, and its absence results in decreased acidification and proteolysis in phagosomes, resembling the function of LIMP-2 (Sattler et al., 2018).

CR3 is an integrin heterodimer composed of CD18 (ITGB2) and CD11B (ITGAM), with roles in chemotaxis and phagocytosis of *Mycobacterium kansasii*, *Mycobacterium smegmatis* or Mtb in both neutrophils and macrophages. Following CR3 activation, its CD18 cytoplasmic tail interacts with the F-actin cytoskeleton to facilitate phagocytosis (Smirnov et al., 2023). In Dd, 'similar to integrin- β A' (SibA) shares characteristics with the integrin- β chain of CR3 and is involved in adhesion and phagocytosis

Table 1. Virulence factors used by *Mycobacterium tuberculosis* and *Mycobacterium marinum* to evade intracellular killing by phagocytes

Virulence factor	Abbreviation	Mycobacteria species	Outcome	Cell type	Reference
Phthiocerol dimycocerosates	PDIM	Mtb	Phagosomal membrane damage Phagosomal escape Blocked autophagy, limited acidification of LC3-positive structures	hMDMs THP-1 PBMCs, THP-1	Augenstreich et al., 2017 Quigley et al., 2017 Bah et al., 2020
Sulfoglycolipid-1	SL-1	Mm	Phagosomal permeabilization Spread into host membranes	THP-1 Zebrafish	Osman et al., 2022 Cambier et al., 2020
		Mtb	Enhanced autophagy for a protective niche Blocked autophagy through TLR2 antagonism	THP-1 PBMCs, THP-1	Mishra et al., 2019 Bah et al., 2020
Trehalose 6,6'-dimycolate Lipoarabinomannan	TDM LAM	Mtb Mtb	Alters phagosome maturation Incorporation into membrane, block of phagosomal maturation	BMDMs, RAW 264.7 hMDMs	Patin et al., 2017 Welin et al., 2008
1-tuberculosinyladenosine PE family protein	1-TbAd PE6	Mtb Mtb	Phagosomal escape Reduced conversion of LC3BI to LC3BII, accumulation of p62, binding to iron	MDDCs RAW 264.7	Buter et al., 2019 Sharma et al., 2021
	PE_PGRS20	Mtb	Ca ²⁺ homeostasis Blocked autophagy initiation by direct interaction with Rab1A protein	THP-1 THP-1, BMDMs, RAW 264.7	Medha et al., 2023 Strong et al., 2021
	PE_PGRS30	Mtb	Inhibition of phagosome-lysosome fusion	J774, THP-1	Iantomasi et al., 2012
	PE_PGRS47	Mtb	Blocked autophagy initiation by direct interaction with Rab1A protein Decreased acidification and lysosomal fusion	THP-1, BMDMs, RAW 264.7 RAW 264.7, BMDMs	Strong et al., 2021 Saini et al., 2016
	PE_PGRS (MMAR_0242)	Mm	Inhibition of lysosomal fusion	Ac, J774, zebrafish	Singh et al., 2016
Early secretory antigenic target 6 kDa	ESAT-6 (EsxA)	Mtb Mtb/Mm Mm	Inhibition of autophagosome formation Inhibition of phagosome maturation Impaired cytosolic translocation Phagosomal membrane damage Escape from the MCV	RAW 264.7 BMDMs RAW 264.7, THP-1 J774, THP-1, zebrafish J774.A1, RAW 264.7, BMDMs J774	Zhang et al., 2012 Xu et al., 2007 Zhang et al., 2016a Osman et al., 2022 Smith et al., 2008 Tan et al., 2006
			Inhibition of phagolysosomal fusion Autophagy manipulation ESCRT and autophagy activation	Dd Dd	Cardenal-Muñoz et al., 2017 Lopez-Jimenez et al., 2018
ESAT-6-like protein EsxH	EsxH	Mtb	Inhibition of ESCRT-dependent trafficking of receptors to the lysosome Disruption of ESCRT function and impairment of phagosome maturation	BMDMs RAW 264.7, HEK293	Mittal et al., 2018 Mehra et al., 2013
ESAT-6-like protein EsxG	EsxG	Mtb	Inhibition of ESCRT-dependent trafficking of receptors to the lysosome	BMDMs	Mittal et al., 2018
ESAT-6 secretion-associated protein B	EspB EspBM	Mtb Mm	Inhibition of autophagosome formation Required for virulence and growth	ANA-1 BMDMs	Huang and Bao, 2016 McLaughlin et al., 2007
Protein kinase G	PknG	Mtb	Blocked autophagy by interaction with RAB14 Release of ubiquitin from UbcH7 as an isopeptidase before attaching it to the substrate Inhibition of Mtb delivery to autophagolysosomes	HeLa, U937 HEK293T, U937 BMDMs, RAW 264.7	Ge et al., 2022 Wang et al., 2021 Zulauf et al., 2018
		Mm	Inhibition of phagosomal maturation	Zebrafish	van der Woude et al., 2014
Protein tyrosine kinase A	PtkA	Mtb	Inhibition of phagosome acidification	THP-1	Wong et al., 2018

Continued

Table 1. Continued

Virulence factor	Abbreviation	Mycobacteria species	Outcome	Cell type	Reference
Protein tyrosine phosphatase A	PtpA	Mtb	Block of TAB3, suppression of NF-κB via ubiquitin interference	HEK293T, U937	Wang et al., 2015
			Anti-apoptotic activity	THP-1	Poirier et al., 2014
			Interaction with vATPase, dephosphorylation of VPS33B and subsequent exclusion of vATPase from the phagosome	THP-1	Wong et al., 2011
			Inhibition of phagosome–lysosome fusion	THP-1	Bach et al., 2008
			Induction of ferroptosis	U937	Qiang et al., 2023
Protein tyrosine phosphatase B	PtpB	Mm	Reduction of MCV acidification by blocking vATPase recruitment, bacterial escape	Ac, Dd	Koliwer-Brandl et al., 2019
			Bacterial escape	Ac, Dd	Koliwer-Brandl et al., 2019
Secreted acid phosphatase M	SapM	Mtb	Inhibition of Mtb delivery to autophagolysosomes	BMDMs, RAW 264.7	Zulauf et al., 2018
Enhanced intracellular survival protein	EIS	Mtb	Blocked phagosomal maturation	THP-1	Puri et al., 2013
			Inhibition of phagosome–lysosome fusion	THP-1	Saikolappan et al., 2012
			Inhibition of phagosome–late endosome fusion via PI3P hydrolysis	RAW 264.7	Vergne et al., 2014
			Bacterial escape	Ac, Dd	Koliwer-Brandl et al., 2019
			Rapamycin-induced autophagy	THP-1	Duan et al., 2016
Nucleoside diphosphate kinase	NdK	Mtb	Modulation of autophagy in a redox-dependent manner	BMDMs, RAW 264.7	Shin et al., 2010
			Attenuation of NADPH oxidase	RAW 264.7	Sun et al., 2013
Lipoprotein	LprI	Mtb	Inhibition of phagosome–lysosome fusion through inactivation of Rab5 and Rab7	RAW 264.7	Sun et al., 2010
			Lysozyme Inhibitor	MDMs, THP-1, RAW 264.7	Sethi et al., 2016
Superoxide dismutase	SOD	Mtb	Reduction of free radicals	C57BL/6	Edwards et al., 2001
Zinc transporter	ZntA, ZntB	Mm	Zn ²⁺ efflux by the transporter from the MCV	Dd	Hanna et al., 2021
Capsular polysaccharide synthesis protein A	CpsA	Mtb	Blocked activity of NADPH oxidase	BMDMs, RAW 264.7	Koster et al., 2017

Abbreviations: Ac, *Acanthamoeba castellanii*; BMDCs, bone marrow-derived dendritic cells; BMDMs, bone marrow-derived macrophages; Dd, *Dictyostelium discoideum*; ESCRT, endosomal sorting complex required for transport; hMDMs, human monocyte-derived macrophages; MCV, mycobacteria-containing vacuole; MDDCs, monocyte-derived dendritic cells; Mm, *Mycobacterium marinum*; Mtb, *Mycobacterium tuberculosis*; PBMCs, peripheral blood mononuclear cells; PI3P, phosphatidylinositol 3-phosphate; vATPase, vacuolar ATPase.

(Cosson and Soldati, 2008). SibB, SibC, SibD and SibE present in Dd have not yet been studied in detail. In Dd, a Venus-trap G protein-coupled receptor, Far1, is also implicated in binding bacterial PAMPs, such as folate and lipopolysaccharides, and serves as a phagocytic receptor (Xu et al., 2021).

Remarkably, amoebae have evolved a plethora of specialized and redundant receptors to engulf a variety of bacteria and interact with various surfaces. Therefore, when single receptor genes are genetically inactivated in Dd, their loss does not significantly affect cellular phenotypes due to the functional redundancy and compensatory mechanisms that exist within extended receptor families.

Actin rearrangement

Phagocytes and Dd reorganize their actin cytoskeleton to engulf mycobacteria (Fig. 1) (Song et al., 2018). Indeed, maintaining the integrity of the actin cytoskeleton is crucial for *M. smegmatis* entry into human macrophages (Dutta et al., 2022). In Dd, the regulation of small GTPases, such as Ras and Rac, through the multidomain

protein RGBARG (RCC1, RhoGEF, BAR and RasGAP-containing protein) is responsible for generating large macropinosomes that facilitate the engulfment of objects with complex shapes, such as mycobacteria (Buckley et al., 2020).

Following phagocytosis, actin facilitates the fusion of early endosomes with phagosomes. However, pathogenic mycobacteria, such as Mm, Mtb and *M. avium*, can disrupt the F-actin network of the host cell, preventing phagosome acidification and maturation, a phenomenon that is not observed with non-pathogenic mycobacteria, such as *M. smegmatis* (Guerin and de Chastellier, 2000). Similarly, in the Dd–Mm model, Mm hinders the actin nucleation-promoting activity of the WASH complex, thereby favouring phagosome maturation arrest and MCV biogenesis (Kolonko et al., 2014).

Remodelling of membrane identity for MCV maturation

Remodelling of phosphatidylinositol phosphates (PIPs) in host endomembranes is evident in both macrophages and Dd infected with mycobacteria, showcasing the conserved role of PIPs during

the phagocytic process (Fig. 3). Phosphatidylinositol 3-phosphate (PI3P) is a crucial regulator of phagosome maturation (Dickson and Hille, 2019; Dormann et al., 2004). After phagocytes and Dd ingest bacteria, there is an initial increase in phosphatidylinositol (3,4,5)-trisphosphate [PI(3,4,5)P₃] at the engulfment site, which is rapidly converted into phosphatidylinositol (3,4)-bisphosphate [PI(3,4)P₂] and PI3P, which is important for recruiting PI3P-binding proteins, such as early endosome antigen 1 (EEA1), FYVE-type zinc finger-containing PIP kinase (PIKfyve), PROPPINs and hepatocyte growth factor-regulated tyrosine kinase substrate (Hrs) (Lawe et al., 2002; Schnettger et al., 2017; Tornero-Ecija et al., 2022; Vines et al., 2023). PI3P is finally

converted by PIKfyve to phosphatidylinositol (3,5)-bisphosphate [PI(3,5)P₂], which drives the accumulation of Rab7 and other lysosomal machinery components on phagosomes in animal cells and Dd (Vines et al., 2023).

During infection, Mtb and Mm secrete three phosphatases: protein tyrosine phosphatase A (PtpA) and B (PtpB) and PI3P acid phosphatase M (SapM), which collectively impede phagosome maturation and promote cytosol escape from the MCV (Bach et al., 2008; Koliwer-Brandl et al., 2019; Puri et al., 2013; Saikolappan et al., 2012; Vergne et al., 2014; Wong et al., 2011). Notably, BCG also secretes SapM via the SecA2 secretion system (Xander et al., 2024).

A Mammalian phagocyte

B *Dictyostelium discoideum*




Fig. 3. Mammalian phagocyte and *Dictyostelium discoideum* responses to mycobacteria. Host cell phagosomes harbour an array of mechanisms to eliminate bacteria in general, including avirulent or attenuated mycobacteria. To survive and proliferate, virulent mycobacteria must disarm this bactericidal compartment and tailor a permissive mycobacteria-containing vacuole (MCV). Phagosome mechanisms of mammalian phagocytes are shown on the left (A), and those of *Dictyostelium discoideum* (Dd) on the right (B). In both mammalian phagocytes and Dd, phagosomal vacuolar ATPase (vATPase) induces proton influx to acidify phagosomes. Concurrently, NADPH oxidase 2 (NOX2) in mammalian phagocytes and NADPH oxidase (NOX) A/B/C in Dd generates reactive oxygen species (ROS), including H_2O_2 . Expression of inducible nitric oxide synthase (iNOS) is exclusive to mammalian phagocytes. When lysosomes fuse with phagosomes, their enzymes (e.g. lysozyme and hydrolases) selectively digest components of the mycobacteria cell wall. Mammals and Dd also both elevate the concentration of copper (Cu^+), magnesium (Mg^{2+}) and zinc (Zn^{2+}) ions to poison intracellular pathogens. Copper is assimilated into the cell by copper transport protein 1 (CTR1) in mammalian phagocytes and SLC31 copper transporters in Dd. In mammals, it is transported through the cytosol by ATOX1 and incorporated into phagosomes and MCVs via ATP7A. In Dd, how Cu^+ reaches the phagosomes remains unknown but Atp7a facilitates its translocation there. In macrophages, Kil2 facilitates Mg^{2+} accumulation in phagosomes, whereas in Dd, Mg^{2+} enters the cytosol via activation of Far1, a G protein-coupled receptor, which also recognizes bacterial lipopolysaccharides (LPS). ZIP transporters transport extracellular Zn^{2+} into macrophages and Dd, and zinc transporters (ZnTs) incorporate it into phagosomes and MCVs. Both mammalian phagocytes and Dd can also limit bacterial access to iron (Fe^{2+}) through NRAMP1/Nramp1. Abbreviations: Mm, *Mycobacterium marinum*; Mtb, *Mycobacterium tuberculosis*; NADP^+ , nicotinamide adenine dinucleotide phosphate (oxidized form); NADPH, nicotinamide adenine dinucleotide phosphate (reduced form).

Rab GTPases – orchestrators of phagosome maturation

Rab GTPases play a pivotal role in orchestrating vesicle trafficking. In both Dd and mammals, PI3P on the MCV facilitates the transition from Rab5 to Rab7, which is essential for vesicular trafficking and for phagosome maturation. In Dd, Rab5A has been identified on the Mm MCV (Barisch et al., 2015) and on *Legionella*-containing vacuoles (Hoffmann et al., 2014). Also, in Dd, the pleckstrin homology (PH) domain-containing protein PripA forms a complex with TbcA, which also promotes the conversion from Rab5A to Rab7A and contributes to phagosome maturation (Tu et al., 2022).

In macrophages infected with Mtb, Rab8A is phosphorylated by the LRR kinase 2 (LRKK2) and is recruited to early endolysosomes, leading to the recruitment of ESCRT components (Herbst et al., 2020). Although data are lacking for mycobacteria-infected Dd, LrrkA, the likely homolog of LRKK2 in Dd, influences intraphagosomal killing of *K. pneumoniae* (Bodinier et al., 2021).

Rab20, an interferon γ (IFN- γ or IFNG)-inducible GTPase, associates rapidly with phagosomes in mycobacteria-infected macrophages to maintain MCV integrity and reduces the cytosolic translocation of mycobacteria (Egami and Araki, 2012; Schnettger et al., 2017; Seto et al., 2011). Notably, Rab20, Rab5C and Rab11B are upregulated in sputum samples from patients with active TB. However, Mtb can trigger Rab20 dissociation, disrupting phagosome maturation and ensuring its survival (Schnettger et al., 2017). In Dd, no Rab20 ortholog has been identified.

PE_PGRS proteins represent some of many virulence factors produced by mycobacteria. Mtb deploys virulence factors that regulate membrane trafficking (Chai et al., 2020), such as PE_PGRS20 and PE_PGRS47, which directly bind to Rab1A, blocking autophagy initiation (Strong et al., 2021). Moreover, the nucleoside diphosphate kinase (NdK) associated with the inactivation of both Rab5 and Rab7 further contributes to Mtb evasion mechanisms (Sun et al., 2010). Mycobacteria have developed numerous other virulence factors to counteract the host defence mechanisms, as elaborated below.

Host defence mechanisms and mycobacterial counterattacks

The phagosome, which initially houses engulfed mycobacteria before evolving into an MCV, acts as a time bomb that is armed with various mechanisms to eliminate the pathogen (Fig. 3). However, mycobacteria have evolved various strategies to counteract these host defence mechanisms, as we discuss in this section.

Chemical warfare – pH acidification, ROS and RNS

The maintenance of an acidic pH in phagosomes is an important defence mechanism against infection. Macrophage phagosomes typically have a pH of 4.5–5, whereas Dd phagosomes have a pH lower than 3.5 (Marchetti et al., 2009). A recent study suggests that the pH of phagosomes can also fluctuate in response to the nature of the cargo and external stimuli (Foote et al., 2019).

The vacuolar ATPase (vATPase) plays a crucial role in the acidification of mammalian and Dd phagosomes by pumping protons in their lumen (Soldati and Neyrolles, 2012). Moreover, neutrophils and macrophages possess the H⁺ channel Hv1 (or HVCN1) for further phagosome acidification (El Chemaly and Demarex, 2012), but it has not been yet described in Dd. Proton accumulation is counteracted by chloride influx and cation efflux of the host (Soldati and Neyrolles, 2012). Remarkably, Mtb sheds the antacid 1-tuberculosynyladenosine (1-TbAd) to arrest lysosomal acidification (Bedard et al., 2023; Buter et al., 2019).

NADPH oxidase 2 (NOX2, also known as gp91^{phox} or CYBB) generates the superoxide anion (O₂^{•−}), which is released into the phagosomal lumen and is transformed into other reactive oxygen species (ROS). In mammals, the NADPH oxidase (NOX) complex consists of the catalytic transmembrane protein NOX2 and the regulatory subunit p22^{phox} (also known as CYBA). In Dd, there are three functional homologs of NOX catalytic subunits: NoxA and NoxB, which are NOX2 homologs, and NoxC, which is a NOX5 homolog. Dd also encodes a regulatory subunit, CybA (p22^{phox} in mammals) and the cytosolic activating factor NcfA (p67^{phox} or NCF2 in mammals) (Zhang et al., 2016b). Although the major source of ROS is NOX2, it can also be produced by the respiratory electron transport chain of mitochondria (Dan Dunn et al., 2015) and by peroxisomes (Huang et al., 2023). ROS contribute to better control of Mtb in macrophages (Chandra et al., 2020). Furthermore, peroxisomal ROS limits the cytosolic growth of Mtb in macrophages derived from Mtb-infected human IPSDMs (Pellegrino et al., 2023). In Dd, NoxA is crucial for eliminating *K. pneumoniae* (Crespo-Yanez et al., 2023) and for lysing *P. aeruginosa* (Ayadi et al., 2024). However, it is unknown whether ROS play a role in Dd cell-autonomous defences during mycobacterial infection, necessitating further investigation.

Increased ROS levels impact both pathogens and host cells, leading to the generation of ROS-detoxifying enzymes by hosts, including superoxide dismutases (SODs), catalases and peroxiredoxins. Mtb has also evolved mechanisms to counteract ROS production by producing SODs (Edwards et al., 2001), by inhibiting ROS production (Pellegrino et al., 2023), by reducing NOX2 expression (Lv et al., 2017) or by modulating host NOX2 activity (Koster et al., 2017; Mo et al., 2022). In response to oxidative stress, mycobacterial SodA, mycolic acids, KatG and SodC may help mitigate the toxicity of extracellular ROS (Tyagi et al., 2015). Additional protective strategies include the use of mycobacterial cytosolic reducing buffers, such as mycothiol and thioredoxins, to maintain the redox environment within the bacteria (Pacl et al., 2018). Moreover, Mtb possesses a type I NADH dehydrogenase that antagonizes phagosomal NOX2 activity (Miller et al., 2010), and Mtb also secretes CpsA, which interferes with NOX activity, thereby reducing ROS concentration in phagosomes and impeding its clearance (Koster et al., 2017).

Neutrophils and macrophages also release extracellular traps in response to pathogens, including in response to Mtb, which affects its course of infection (Garcia-Bengoa et al., 2023). In multicellular Dd slugs infected with *K. pneumoniae* or exposed to lipopolysaccharide, sentinel cells can produce extracellular traps, which is dependent on ROS produced by NoxA, NoxB and NoxC (Zhang et al., 2016b). However, further research is required to determine the role of extracellular traps during mycobacterial infection in Dd.

Reactive nitrogen species (RNS), such as nitric oxide (NO), and inducible NO synthase (iNOS, also known as NOS2, Box 1) are crucial components of the innate immune responses against mycobacterial infections, as reviewed by Yang et al. (2009) and Jamaati et al. (2017). Mtb-infected mouse and human macrophages induce iNOS production to significantly different levels *in vitro*. Indeed, human macrophages generate lower levels of NO than mouse macrophages, which complicates our understanding of iNOS in Mtb control (Jung et al., 2013). Furthermore, NO can direct macrophages to form multinucleated giant cells, which create a permissive environment for mycobacterial persistence (Gharun et al., 2017). It is worth noting that Dd has no recognizable NO synthase, precluding the study of NO during mycobacterial infections.

Lysosomal enzymes

The process of phagosome maturation into a degradative and bactericidal milieu is conserved between mammalian cells and Dd (Boulais et al., 2010). The lysosome fuses with the phagosome following Rab7 interaction with members of the homotypic fusion and protein sorting (HOPS) complex, such as Vps18 (Jani et al., 2023 preprint), which then functions as a tethering complex involved in vesicle trafficking. Following this, lysosomal enzymes are delivered to phagosomes to target specific components of the cell wall and membranes of the pathogen (Trivedi et al., 2020).

Mtb employs multiple evasion strategies at this stage. It upregulates the lipoprotein LprL to neutralize lysozyme (Box 1) activity in peritoneal and monocyte-derived macrophages (Sethi et al., 2016), and it expresses PE_PGRS proteins (Iantomasi et al., 2012; Saini et al., 2016) and EsxA (also known as ESAT-6) (Xu et al., 2007; Tan et al., 2006) to inhibit phagosome maturation. Additionally, Mtb exploits lysosome-poor monocyte-derived cells for persistence *in vivo* (Zheng et al., 2024).

In the Dd model, the membrane-permeabilizing proteins AlyL and BpiC, which target peptidoglycans and lipopolysaccharides, respectively, are effective against *K. pneumoniae* (Crespo-Yanez et al., 2023), but it is not yet known if these are effective against mycobacteria. Furthermore, in Dd, Kil1 (Box 1) and Kil2 (Box 1) play vital roles in bacterial digestion within phagosomes. Kil1 delivers proteases (Bodinier et al., 2020), whereas Kil2, activated by folate, enhances magnesium ion transfer to the phagosomal lumen, improving lysosomal enzyme efficiency. Both Kil1 and Kil2 contribute significantly to *K. pneumoniae* digestion (Crespo-Yanez et al., 2023), although their roles in mycobacterial infection remains unexplored.

Metal transporters

The regulation of essential divalent metals, such as zinc, copper, iron and magnesium, as enzyme cofactors, is crucial for both hosts and pathogens. Host cells employ strategies to either poison intracellular pathogens or to deprive them of essential micronutrients.

Zinc is an abundant micronutrient that is crucial for regulating gene expression, cell processes, immune responses and/or antioxidant defences, among other roles in the host. Indeed, about 10% of the human proteome present with zinc-binding motifs (reviewed by Maret, 2013). In eukaryotes, ZIP family transporters facilitate extracellular zinc entry into the cell cytosol, whereas the zinc transporter (ZnT) proteins export zinc outside the cell or to the lumen of endocytic and secretory organelles. In Dd, seven ZIP transporters (ZpIA-G) and four ZnT transporters (ZntA-D) have been identified (reviewed by Dunn et al., 2018). Upon Mm infection of Dd cells, the ZntA and ZntB efflux pumps are recruited to the MCV, increasing the zinc concentration in MCVs and restricting mycobacteria growth (Barisch et al., 2018; Hanna et al., 2021), as also observed in macrophages infected with Mtb (Botella et al., 2011; Neyrolles et al., 2013). Mycobacteria counteract zinc poisoning with metal efflux pumps, including CtpC in Mtb and Mm, and CtpG in BCG (Boudehen et al., 2022; Chen et al., 2022; Hanna et al., 2021).

Copper, a redox-active metal, undergoes cycles between the Cu⁺ and Cu²⁺ ion states in the MCV under physiological conditions. Mammalian copper transport protein 1 (CTR1) pumps copper into the cytosol, where it binds ATOX1, delivering it to ATP7A present on the MCV membrane (reviewed by Neyrolles et al., 2015). Dd expresses six copper transporters, including three SLC31 copper transporters and three P-type Cu-ATPases, one sharing homology with human ATP7A (Buracco et al., 2018). In mammalian phagocytes and Dd, the recruitment of ATP7A to phagosomes

might enhance copper pumping into the phagosomal lumen, whereas p80, a predicted copper transporter homolog of CTR1, might play a role in copper influx to the cytosol (Buracco et al., 2018; Neyrolles et al., 2015). Mycobacteria deploy defences against copper stress. They express Cu⁺-binding metallothionein (MymT), copper transport protein B (MctB) and copper (I) transporting P_{1B}-type ATPases (CtpV and CtpB) (reviewed by Botella et al., 2012; Darwin, 2015; Leon-Torres et al., 2020). Experimental use of copper chelators on Mtb-infected macrophages reduced bacterial load, indicating a potential role for copper in Mtb intracellular growth (Libardo et al., 2018; Shah et al., 2016; Speer et al., 2013). However, copper fluctuations during Dd growth did not affect *Legionella* infection (Buracco et al., 2018), and similar research is needed for mycobacteria.

Nutritional immunity refers to host strategies that impede pathogen growth by limiting metal availability. In Mtb-infected macrophages, the host protein NRAMP1 diminishes the availability of iron by redirecting its storage from the phagolysosome to the cytosol (reviewed by Murdoch and Skaar, 2022). NRAMP1 is expressed both by macrophages and Dd and appears to restrict the growth of mycobacteria (Medapati et al., 2017; Peracino et al., 2006). Mammals also produce NRAMP2 (also known as SLC11A2), but Dd Nramp2 is more akin to protist and fungal Nramp proteins (Peracino et al., 2013). Dd Nramp2 restricts *Francisella* growth (Brenz et al., 2017), whereas data for mycobacteria are still lacking. The iron response of mycobacteria involves siderophores (Box 1), which facilitate their growth in both Dd and mouse macrophages (Knobloch et al., 2020). In the case of Mtb and Mm, distinct siderophores, mycobactin and carboxymycobactin, have been identified. For a comprehensive review of the relevance of iron to TB pathogenesis, see Rodriguez et al. (2022).

Host metabolism

Target of rapamycin complex 1 (TORC1) (Box 1) inhibits autophagy, promoting animal and Dd cell growth by boosting ribosome biogenesis and protein translation in nutrient-rich conditions. However, under conditions of low nutrients, TORC1 is inhibited, and autophagy provides the metabolites and energy required to sustain essential functions in both mammalian cells and Dd (Cardenal-Muñoz et al., 2017; Linares et al., 2013).

Upon Mm infection, the mammalian target of rapamycin (mTOR) kinase is inhibited, resulting in a host-protective effect by enhancing autophagy and glycolysis in Dd and zebrafish larvae that lack adaptive immunity, relying solely on innate responses (Cardenal-Muñoz et al., 2018; Pagán et al., 2022). Similar responses have been observed in Mm- and Mtb-infected THP-1 macrophages (Pagán et al., 2022). Moreover, in mammalian cells, the interaction of stress granule (Box 1) proteins, such as NUFIP2 or G3BP1, with GABA receptor-associated proteins (GABARAPs) ensures the inactivation of mTORC1 via the Ragulator–Rag system (Jia et al., 2022). However, mTOR deficiency can also lead to a significant innate susceptibility to mycobacteria, leading to the death of infected macrophages through (or due to) elevating mitochondrial energy metabolism driven by glycolysis in response to infection (Pagán et al., 2022).

Experiments in Dd have shown that Mm blocks autophagic flux, resulting in the accumulation of membranes and cytoplasmic material in the MCV, potentially supporting bacterial survival within this niche (Cardenal-Muñoz et al., 2017).

Host resilience

During infection, the slightly acidified MCV milieu activates the membranolytic factor EsxA of Mtb and Mm (Bao et al., 2021). This,

coupled with other virulence factors, such as PDIMs, contribute to MCV damage (Augenstreich et al., 2017; Bussi et al., 2023; also reviewed by Augenstreich and Briken, 2020; Chandra et al., 2022). As observed in Dd, the membranolytic activity of these mycobacterial virulence factors benefits from membrane microdomains that contain sterols and vacuolins, which are homologs of mammalian lipid raft-associated flotillins (Box 1) (Bosmani et al., 2021 preprint). Similarly, Mtb can also cause lysosomal damage, triggering protease leakage into the cytosol and leading to mitochondrial disruption (Bhattacharyya et al., 2023; Bussi et al., 2022; Radulovic et al., 2022). Such damage interferes with cellular functions and activates immune responses. As a result, cells employ various mechanisms to recognize and limit damage to their membranes caused by intracellular pathogens. Although the characteristics that determine which repair pathway responds to membrane damage are still poorly understood, the extent of the damage appears to be crucial for activating repair mechanisms. Minor damage to membranes is repaired by the ESCRT system (Radulovic et al., 2018) or through membrane contact sites (Radulovic et al., 2022). More extensive damage to host cell membranes induces the activation of autophagy (Schnettger et al., 2017).

In this section, we review and discuss the initial events that follow mycobacterial-induced host membrane damage, highlighting the mechanisms employed by host cells to repair the damage and how these repair machineries are coordinated (Fig. 1).

Sensing mycobacteria-triggered damage

Host galectin-3 (LGALS3), galectin-8 (LGALS8) and galectin-9 play vital roles in recognizing host glycolipids and glycoproteins on the luminal leaflet of the MCV that become exposed to the cytosol after Mtb-induced membrane damage (Bell et al., 2021).

ESX1-dependent damage triggered by Mtb induces the recruitment of galectin-3 in iPSDMs and THP-1-infected cells (Fig. 1) (Augenstreich et al., 2017; Beckwith et al., 2020). Additionally, in bone marrow-derived macrophages, galectin-9 binds to the cytosolically exposed arabinogalactan of the Mtb cell wall (Morrison et al., 2023; Wu et al., 2021). In macrophages, galectin-8 might play a direct role in the repair and clearance of Mtb-induced MCV damage (Jani et al., 2023 preprint), possibly due to its role in recruiting the autophagy machinery (Bell et al., 2021). Overall, although galectins are involved in damage sensing, they do not seem to limit mycobacteria growth (Morrison et al., 2023). Dd lacks galectin orthologs but possesses functionally homologous discoidins (DscA, DscC, DscD and DscE), emphasizing the evolutionary adaptations in response to mycobacterial infections in different hosts (Aragao et al., 2008; Mathieu et al., 2010). Cytosolic discoidins recognize glycosylated Mm lipids or proteins exposed to the cytosol after MCV rupture (Fig. 1). Specific ligands for discoidins in the Mm cell wall and the implications of their recognition are still under investigation (López-Jimenez, 2017).

Ubiquitin as a 'repair-me' and 'eat-me' signal

Ubiquitination (Box 1) is a process conserved between animal cells (Madiraju et al., 2022) and Dd (Pergolizzi et al., 2019; Raykov et al., 2023; Xiong et al., 2023) and is important in Mtb-infected human macrophages and Mm-infected Dd. Smurf1 transfers lysine (K) 48-linked ubiquitin chains that serve as a signal for degradation by the proteasome, and parkin proteins transfer K63-linked ubiquitin chains recognized by autophagy adapters (Franco et al., 2017; Manzanillo et al., 2013). Additionally, essential E3 ubiquitin ligases such as tumour necrosis factor α (TNF- α) receptor-associated factors (TRAFs) and tripartite motif-containing proteins (TRIMs) also contribute to infection outcomes.

In humans, seven TRAFs have been characterized (TRAF1-7). TRAF6, a RING-type E3 ligase, influences various host immune defence functions, such as the transcription of TNF- α (Kim et al., 2022b) and transforming growth factor β (TGF- β , encoded by *TGFB*) (Landstrom, 2010), the induction of autophagy (Kim et al., 2022a) or the maturation of autophagosomes under oxidative stress (Wang et al., 2022).

In Dd, over 40 TRAF-like proteins have been predicted, with 16 of them presenting RING, zinc finger and TRAF domains akin to those of mammalian TRAF2, TRAF3, TRAF5 and TRAF6 (Dunn et al., 2018). In particular, the TRAF E3 ubiquitin ligase TraFE (a TRAF6 ortholog) plays a pivotal role during Mm infection in Dd. Specifically, TraFE is proposed to act as a coordinator between ESCRT and autophagy pathways through TraFE-mediated K63 ubiquitination of yet unknown target(s) (Raykov et al., 2023).

TRIM proteins are a conserved ubiquitin ligase family that have diverse roles in immune responses or autophagy. The expression of 20 TRIM genes in patients with active TB was reported to be decreased compared to that in patients with latent TB or healthy donors, linking these genes to the pathogenesis of TB and highlighting their potential utility as TB biomarkers (Chen et al., 2018). Mammalian TRIM proteins have been linked to Mtb (TRIM16, TRIM22, TRIM27 and 32), are shown to induce autophagy (TRIM16, TRIM22 and TRIM32) (Chauhan et al., 2016; Lou et al., 2018; Romagnoli et al., 2023) and are counterintuitively associated with Mtb growth (TRIM14, TRIM25, TRIM36 and TRIM56) (Hoffpauir et al., 2020). Dd possesses a single TRIM protein (Dunn et al., 2018) with homology to human TRIM37, associated with autophagy and viral restriction (Gu et al., 2023). However, the specific role of Dd TRIM in mycobacterial infection remains understudied (Raykov, 2021).

Influx of extracellular Ca^{2+}

In eukaryotes, efficient membrane repair often relies on the influx of Ca^{2+} (Gronski et al., 2009; Yuan et al., 2001), triggering the accumulation of calcium-binding proteins at the wound site to support cytoskeletal reorganization and the assembly of signalling molecules, as reviewed by Cooper and McNeil (2015).

During Mtb phagocytosis, intracellular Ca^{2+} levels rise in response to opsonized or heat-killed Mtb, whereas in response to live Mtb, intracellular Ca^{2+} levels decrease in macrophages, affecting proteins such as calmodulin and phosphorylated Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII) (Jayachandran et al., 2007). In Mtb-infected macrophages, intracellular Ca^{2+} signalling is crucial for membrane trafficking, which involves LRRK2, Rab8 and ESCRT machinery recruitment at damaged endolysosomes (Herbst et al., 2020).

Upon sterile damage in Dd, accumulation of actin filaments at the wound site relies on Ca^{2+} influx, which is crucial for repair (Talukder et al., 2020). However, no study is available relating calcium and mycobacteria infection in the amoeba model.

Remarkably, mycobacterial virulence factors, such as mannose-capped LAM (ManLAM) (Rojas et al., 2000), PE6 (Medha et al., 2023), the secreted protein PE_PGRS1 that contains seven Ca^{2+} binding domains (Yu et al., 2023), and the calcium P-type ATPase CtpF (Maya-Hoyos et al., 2022), thwart host intracellular Ca^{2+} increase to ensure bacteria survival.

Membrane-damage repair mechanisms

Host cells respond to membrane damage through coordinated repair strategies that are influenced by the extent and characteristics of the damage. In this subsection, we discuss primary repair mechanisms in mammalian macrophages and Dd and shed light on

recent insights into the membrane damage that is induced by mycobacteria.

Small membrane damage in the MCV of Mtb-infected macrophages (Philips et al., 2008) and of Mm-infected Dd (Lopez-Jimenez et al., 2018) is primarily repaired by ESCRT to ensure bacterial containment, highlighting the conservation of this repair pathway (Fig. 1). In Mm-infected Dd, ESCRT proteins localize to the MCV, forming distinct patches or rings. In response, Mtb employs a countermeasure by secreting EsxG and EsxH factors through the ESX-3 type VII secretion system to hinder the recruitment of the ESCRT-III machinery to sites of MCV damage (Mittal et al., 2018).

ESCRT is not limited to endolysosomal repair; it also contributes to plasma membrane repair. In Mtb-infected THP-1 cells, damage-induced calcium influx triggers the recruitment of ALG2 to the plasma membrane (Beckwith et al., 2020). In TR146 mammalian cells infected with *Candida albicans*, ALG2 then interacts with ALG2-interacting protein X (ALIX, also known as PDCD6IP) and with CHMP proteins of the ESCRT-III complex, facilitating plasma membrane repair (Westman et al., 2022). This phenomenon is similar in Dd, where PefA, one of the two penta-EF hand homologs of mammalian ALG2, and ALIX are recruited to the site of plasma membrane damage in a calcium-dependent manner (Talukder et al., 2020).

Autophagy is active in both mammals and Dd, which possesses orthologs of most autophagy-related mammalian proteins, as reviewed by Calvo-Garrido et al. (2010), Cosson and Soldati (2008) and Mesquita et al. (2017). In Mm-infected Dd, an increased number of Atg8- and Atg18-positive structures, along with the recruitment of autophagy markers, such as ubiquitin and p62, are observed soon after infection (Cardenal-Muñoz et al., 2017).

Recent findings propose that mammalian ATG8 proteins bind to ESCRT components to maintain the integrity of autophagosomes (Javed et al., 2023). Autophagy-mediated membrane repair is exemplified by the presence of Vps32 (also known as CHMP4), Atg8 and the autophagy adaptor p62 (SQSTM1) at pathogen-containing vacuoles in mammalian cells infected with *Salmonella* (Kreibich et al., 2015) or *C. albicans* (Lapaquette et al., 2022), or in Dd infected with Mm (Lopez-Jimenez et al., 2018; Raykov et al., 2023) or *A. fumigatus* (Ferling et al., 2020).

In the Dd–Mm model, the ESX-1 secretion system of Mm induces a robust repair response involving both the ESCRT-III and autophagy pathways (Lopez-Jimenez et al., 2018; Raykov et al., 2023). The coordination of both pathways by TrafE has been demonstrated in the Dd–Mm model (Raykov et al., 2023), and it is thus plausible that a similar scenario occurs in Mtb-infected mammalian cells.

Recent research in mammalian cells has revealed additional cellular mechanisms that mend damaged membranes, notably involving membrane–contact sites and the formation of stress granules (Bussi et al., 2023). Following lysosomal damage, phosphatidylinositol 4-kinase type 2- α (PI4K2A) orchestrates the synthesis of phosphatidylinositol 4-phosphate (PI4P), resulting in its excessive accumulation on the injured vacuole. This buildup triggers the recruitment of oxysterol-binding protein (OSBP), an endoplasmic reticulum (ER)–Golgi protein, and OSBP-related proteins (ORPs) 9–11 (also known as OSBPL9–11), giving rise to extensive ER–lysosomal membrane contact sites. OSBP plays a crucial role in transferring cholesterol and phosphatidylserine from the ER to damaged lysosomes, reciprocated by PI4P, thereby regulating PI4P levels. Likewise, ORPs, with their lipid binding and transport capabilities, facilitate PI4P-driven phosphatidylserine transfer from the ER to damaged lysosomes, mirroring the functions of ORP1L (an isoform of ORP1, also known as

OSBPL1A), ORP5 (OSBPL5) and ORP8 (OSBPL8) (Chung et al., 2015; Radulovic et al., 2022; Tan and Finkel, 2022). Notably, OSBP is recruited to Mtb and Mm-containing vacuoles in an ESX-1-dependent manner. In Dd, OSBP8 is also recruited early during infection and is present on ER tubules forming contact sites with the MCV. OSBP8 depletion negatively impacts Dd cell viability and enhances Mm growth (Anand et al., 2023).

In human macrophages infected with Mtb, stress granules and other condensates rapidly nucleate nearby damaged MCVs or endolysosomes, acting as a plug to stabilize ruptured membranes. The complete engulfment of the droplet inside the compartment aids in its effective repair, either spontaneously or facilitated by the ESCRT machinery (Bussi et al., 2023). Notably, mammalian ATG8 proteins can interact with stress granule proteins, influencing stress granules and mTOR responses to Mtb damage (Jia et al., 2022), indicative of a coordinated cellular response. Although Dd induces protein aggregation under stress, the specific role of these protein condensates as potential patches upon mycobacteria-induced membrane damage remains unknown.

Mycobacteria restriction

Extensive research has been conducted on xenophagy, a specific form of autophagy that targets intracellular pathogens, to understand its role in controlling Mtb infection and in limiting its growth in macrophages (Gutierrez et al., 2004). This phenomenon has been observed in various models, including in Dd–Mm, BMDM–Mm and iPSDM–Mtb, demonstrating its conservation between amoebae and mammals (Bernard et al., 2021; Collins et al., 2009; Lopez-Jimenez et al., 2018; Simeone et al., 2012). Xenophagy activation involves galectin recruitment and ubiquitination of the bacteria at damaged MCVs (Fig. 1). It can also be initiated by ubiquitinated bacteria in the cytosol or be enhanced by cGAS–STING-dependent signalling, which recognize foreign DNA in the cytosol (Bernard et al., 2021; Watson et al., 2015, 2012).

The deletion of autophagy-related genes in mouse macrophages *in vivo* is linked to increased levels of cytosolic Mtb and Mm, leading to rapid necrotic cell death (Feng et al., 2024; Golovkine et al., 2023). Specifically, knocking out autophagy-related genes, such as ATG7, ATG14 and ATG16, increased Mtb growth and subsequent macrophage cell death. Although xenophagy triggers a type I interferon immune response in mammals to restrict mycobacteria growth, some authors propose that mycobacterial control is mainly achieved by promoting phagosome maturation and by activating the autophagy machinery (Schnettger et al., 2017). Accordingly, Atg8a (an LC3 family protein commonly used as autophagy reporter) is observed at MCVs in the Dd–Mm model (Cardenal-Muñoz et al., 2017; Lopez-Jimenez et al., 2018; Raykov et al., 2023) and in mammalian macrophages infected with Mtb (Bernard et al., 2021). Additionally, the autophagy regulator DRAM1 forms puncta near mycobacteria, resulting in the colocalization of DRAM1, LC3 and Mm in zebrafish and macrophages (Banducci-Karp et al., 2023). However, deciphering the specific contribution of autophagy-related genes to mycobacteria control poses challenges, particularly given the contrasting findings related to the knockout of the ATG5-coding gene (Castillo et al., 2012; Golovkine et al., 2023; Kimmey et al., 2015; Kinsella et al., 2023; Watson et al., 2012). Overall, the consensus in the field is that autophagy is the primary pathway that restricts cytosolic mycobacteria, with ATG5 exhibiting additional and unique functions beyond autophagy (Deretic and Wang, 2023).

In the context of TB, guanylate-binding proteins (GBPs, Box 1) are associated with susceptibility to bacterial infection and host

response (Esterhuyse et al., 2015; Kim et al., 2012), as they bind to MCVs and limit the growth of intracellular pathogens, such as BCG (Marinho et al., 2020). However, their ability to restrict the growth of Mtb does not extend to virulent Mtb expressing the ESX-1 secretion system in infected mice (Olive et al., 2023). Conversely, Mtb secretes virulence factors, such as PE31, that induce GBP1 production in macrophages and reduce macrophage apoptosis (Ali et al., 2020). The role of the single Dd homolog of human GBPs remains understudied (Raykov, 2021).

Despite host efforts to control and eliminate mycobacteria, whether contained within the MCV or in the cytosol, virulent mycobacteria can also win, proliferate and disseminate. This last phase of the productive infection cycle will be presented in the following section.

Mycobacteria triumph and dissemination

The intracellular localization of mycobacteria cycles between the MCV and the host cytosol in a dynamic manner, reflecting the competing processes of mycobacteria-triggered damage and host repair machineries in Dd (Cardenal-Muñoz et al., 2017; Lopez-Jimenez et al., 2018; Raykov et al., 2023) as well as in mouse macrophages (Schnettger et al., 2017) and iPSDMs (Bernard et al., 2021). Host cells employ diverse mechanisms to constrain and eradicate mycobacteria. For instance, Dd initially expels intracellular bacteria by exocytosis, ejection or by host lysis; infected cells are then excluded from multicellular aggregates by a collective effort, thereby ensuring germ-free spores (López-Jiménez et al., 2019 preprint).

Despite these host defence responses, mycobacteria can persist and propagate within cells, leading to their dissemination to other cells or organisms (Fig. 1). The rupture of the phagolysosome or the cytosolic sensing of bacterial DNA activates various cell death processes that are crucial for Mtb spread (Ruan et al., 2024). In some cases, cell death can even be induced by extracellular contact with mycobacterial aggregates (Toniolo et al., 2023). These host cell death processes include non-lytic and lytic processes, such as apoptosis in mouse macrophages and hMDMs (Aguilo et al., 2013; Augenstreich et al., 2017), necrosis in BMDMs (Lee et al., 2006), necroptosis (Box 1) in mice (Pajuelo et al., 2018; Zhao et al., 2017), pyroptosis (Box 1) in THP-1 cells and peripheral blood mononuclear cells (Beckwith et al., 2020; Golovkine et al., 2023), as well as other mechanisms mediated by interferons in BMDMs (Zhang et al., 2021) and by TNF signalling in mammalian cells or zebrafish (Roca et al., 2019).

In Dd, an additional mechanism of mycobacterial dissemination involves Mm egress from the cell via a regulated process that balances host cell integrity with infection spread (Hagedorn et al., 2009). Ejection, leading to host plasma membrane damage, is controlled by an autophagosome structure in Dd, whereas Mm uses a barrel-shaped F-actin structure (ejectosome) for egress (Fig. 1). ESCRT and autophagic proteins at the distal pole shield the host cell from lysis during Mm egress (Gerstenmaier et al., 2015), with Vps4 completing membrane sealing upon bacterial exit. Despite the unknown localization mechanism of Vps4 (Gerstenmaier et al., 2024), the dependence of its recruitment on TrafE at MCV damage sites in Dd (Raykov et al., 2023) suggests some level of coordination with the response at the ejectosome site.

Conclusion

The natural ability of Dd as a bacterial predator and its reliance on only cell-autonomous defences makes it crucial for exploring innate immunity against various microorganisms. Specifically, the synergy

between Dd and Mm as a model for investigating host–pathogen interactions presents a unique opportunity. This is primarily due to the similarities in virulence factors between Mm and Mtb, as well as their infection processes in Dd and macrophages, respectively. Although Dd offers experimental advantages due to its conserved innate immune pathways, it is also essential to consider its evolutionary distance from macrophages.

To trigger initial phagocytosis of (myco)bacteria, Dd exhibits a myriad of receptors, although their genetic study is complicated by functional redundancy. Nonetheless, cytosolic proteins such as TirA are recognized for their significance in sensing pathogens other than mycobacteria. Additionally, although lectin receptors and integrin-like receptors are present in Dd, further investigation is necessary to delineate their roles during (myco)bacterial infections. After phagocytosis, mycobacteria transiently reside within phagosomes, which usually mature into the MCV, the ultimate bactericidal machine to eradicate and infection. Dd has been instrumental in elucidating the specific role of PI(3,5)P2 in Rab7 accumulation and lysosomal biogenesis. However, certain important mammalian Rabs, such as Rab20, remain unidentified in Dd.

Both mammalian phagocytes and Dd employ mechanisms such as vATPase accumulation and ROS production against pathogens. However, research gaps persist in Dd, including the identification of additional acidification channels (Hv1 H⁺ channel) and the specific role of ROS during mycobacterial infection. Although neutrophils, macrophages and Dd release extracellular traps, their involvement in response to pathogens is only understood in mammalian cells.

Phagosomal maturation involves the delivery of lysosomal enzymes into the phagosome in both macrophages and Dd. Although several lysosomal enzymes (AlyL, BPiC, Kil1 and Kil2) have been implicated in Dd infected with *K. pneumoniae*, their specific roles in the context of mycobacterial infection require further investigation. Other antibacterial mechanisms employed by Dd and macrophages consist of regulating the presence of essential metals to poison pathogens or limit their availability, with zinc and copper being particularly noteworthy in Mtb-infected macrophages. However, copper fluctuations were not considered important in *Legionella*-infected Dd, and further research is required on mycobacterial infection.

Both Mtb and Mm convert the phagosome into a more friendly MCV, in part by causing damage to the membrane surrounding the mycobacteria. Galectins are implicated in damage sensing triggered by mycobacteria in mammals, yet specific ligands of discoidins (galectin homologs) in Dd remain under intense research. Ubiquitination is also crucial during infection, with the E3 ubiquitin ligase TrafE demonstrated to be essential in coordinating repair mechanisms in Dd. Despite some similarities, this precise pathway has not yet been described in mammals. However, and reciprocally, other ubiquitin ligases such as TRIMs, implicated in mycobacterial infections in mammals, deserve further investigation in the Dd model.

Repair mechanisms, including ESCRT, autophagy, membrane contact sites and stress granules, are conserved in eukaryotes. Although ESCRT and autophagy are described in mammals and Dd in response to mycobacterial damage, the precise roles of membrane contact sites and stress granules in the Dd–Mm model remain to be elucidated.

Overall, the Dd–Mm model facilitates the elucidation of critical mechanisms in a safe and 3R (Replacement, Reduction and Refinement)-compliant manner, aiming to advance the development of effective therapies against TB infection.

Acknowledgements

The authors thank the funders for their support.

Competing interests

The authors declare no competing or financial interests.

Funding

This work was funded by the Swiss National Science Foundation research (SNSF, Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung) grants 310030_169386 and 310030_188813 to T.S. S.G.-G. is a recipient of the SNSF Swiss Postdoctoral Fellowship (TMPFP3_217291).

References

Abdallah, A. M., Savage, N. D., Van Zon, M., Wilson, L., Vandenbroucke-Grauls, C. M., Van Der Wel, N. N., Ottenhoff, T. H. and Bitter, W. (2008). The ESX-5 secretion system of *Mycobacterium marinum* modulates the macrophage response. *J. Immunol.* **181**, 7166–7175. doi:10.4049/jimmunol.181.10.7166

Aguilo, J. I., Alonso, H., Uranga, S., Marinova, D., Arbues, A., De Martino, A., Anel, A., Monzon, M., Badiola, J., Pardo, J. et al. (2013). ESX-1-induced apoptosis is involved in cell-to-cell spread of *Mycobacterium tuberculosis*. *Cell. Microbiol.* **15**, 1994–2005. doi:10.1111/cmi.12169

Ali, M. K., Zhen, G., Nzungize, L., Stojkoska, A., Duan, X., Li, C., Duan, W., Xu, J. and Xie, J. (2020). *Mycobacterium tuberculosis* PE31 (Rv3477) attenuates host cell apoptosis and promotes recombinant *M. smegmatis* intracellular survival via up-regulating GTPase guanylate binding protein-1. *Front. Cell Infect. Microbiol.* **10**, 40. doi:10.3389/fcimb.2020.00040

Anand, A., Mazur, A. C., Rosell-Arevalo, P., Franzkoch, R., Breitsprecher, L., Listian, S. A., Huttel, S. V., Muller, D., Schafer, D. G., Vormittag, S. et al. (2023). ER-dependent membrane repair of mycobacteria-induced vacuole damage. *mBio* **14**, e0094323. doi:10.1128/mbio.00943-23

Aqdas, M., Singh, S., Amir, M., Maurya, S. K., Pahari, S. and Agrewala, J. N. (2021). Cumulative signaling through NOD-2 and TLR-4 eliminates the *Mycobacterium tuberculosis* concealed inside the mesenchymal stem cells. *Front. Cell Infect. Microbiol.* **11**, 669168. doi:10.3389/fcimb.2021.669168

Arafah, S., Kicka, S., Trofimov, V., Hagedorn, M., Andreu, N., Wiles, S., Robertson, B. and Soldati, T. (2013). Setting up and monitoring an infection of *Dictyostelium discoideum* with mycobacteria. *Methods Mol. Biol.* **983**, 403–417. doi:10.1007/978-1-62703-302-2_22

Aragao, K. S., Satre, M., Imbert, A. and Varrot, A. (2008). Structure determination of Discoidin II from *Dictyostelium discoideum* and carbohydrate binding properties of the lectin domain. *Proteins* **73**, 43–52. doi:10.1002/prot.22038

Augenstein, J. and Briken, V. (2020). Host cell targets of released lipid and secreted protein effectors of *Mycobacterium tuberculosis*. *Front. Cell Infect. Microbiol.* **10**, 595029. doi:10.3389/fcimb.2020.595029

Augenstein, J., Arbues, A., Simeone, R., Haanappel, E., Wegener, A., Sayes, F., Le Chevalier, F., Chalut, C., Malaga, W., Guilhot, C. et al. (2017). ESX-1 and phthiocerol dimycocerosates of *Mycobacterium tuberculosis* act in concert to cause phagosomal rupture and host cell apoptosis. *Cell. Microbiol.* **19**, e12726. doi:10.1111/cmi.12726

Ayadi, I., Lamrabet, O., Munoz-Ruiz, R., Jauslin, T., Guilhen, C. and Cosson, P. (2024). Extracellular and intracellular destruction of *Pseudomonas aeruginosa* by *Dictyostelium discoideum* phagocytes mobilize different antibacterial mechanisms. *Mol. Microbiol.* **121**, 69–84. doi:10.1111/mmi.15197

Bach, H., Papavasasundaram, K. G., Wong, D., Hmama, Z. and Av-Gay, Y. (2008). *Mycobacterium tuberculosis* virulence is mediated by PtpA dephosphorylation of human vacuolar protein sorting 33B. *Cell Host Microbe* **3**, 316–322. doi:10.1016/j.chom.2008.03.008

Bah, A., Sanicas, M., Nigou, J., Guilhot, C., Astarie-Dequeker, C. and Vergne, I. (2020). The lipid virulence factors of *Mycobacterium tuberculosis* exert multilayered control over autophagy-related pathways in infected human macrophages. *Cells* **9**, 666. doi:10.3390/cells9030666

Banducci-Karp, A., Xie, J., Engels, S. A. G., Sarantaris, C., van Hage, P., Varela, M., Meijer, A. H. and van der Vaart, M. (2022). DRAM1 promotes lysosomal delivery of *Mycobacterium marinum* in macrophages. *Cells* **12**, 828. doi:10.3390/cells12060828

Bao, Y., Wang, L. and Sun, J. (2021). A small protein but with diverse roles: a review of EsxA in mycobacterium-host interaction. *Cells* **10**, 1645. doi:10.3390/cells10071645

Barisch, C. and Soldati, T. (2017). *Mycobacterium marinum* degrades both triacylglycerols and phospholipids from its *Dictyostelium* host to synthesise its own triacylglycerols and generate lipid inclusions. *PLoS Pathog.* **13**, e1006095. doi:10.1371/journal.ppat.1006095

Barisch, C., Lopez-Jimenez, A. T. and Soldati, T. (2015). Live imaging of *Mycobacterium marinum* infection in *Dictyostelium discoideum*. *Methods Mol. Biol.* **1285**, 369–385. doi:10.1007/978-1-4939-2450-9_23

Barisch, C., Kalinina, V., Lefrançois, L. H., Appiah, J., López-Jiménez, A. T. and Soldati, T. (2018). Localization of all four ZnT zinc transporters in *Dictyostelium* and impact of ZntA and ZntB knockout on bacteria killing. *J. Cell Sci.* **131**, jcs222000. doi:10.1242/jcs.222000

Beckwith, K. S., Beckwith, M. S., Ullmann, S., Saetra, R. S., Kim, H., Marstad, A., Asberg, S. E., Strand, T. A., Haug, M., Niederweis, M. et al. (2020). Plasma membrane damage causes NLRP3 activation and pyroptosis during

Mycobacterium tuberculosis infection. *Nat. Commun.* **11**, 2270. doi:10.1038/s41467-020-16143-6

Bedard, M., Van Der Niet, S., Bernard, E. M., Babunovic, G., Cheng, T. Y., Aylan, B., Grootemaat, A. E., Raman, S., Botella, L., Ishikawa, E. et al. (2023). A terpene nucleoside from *M. tuberculosis* induces lysosomal lipid storage in foamy macrophages. *J. Clin. Invest.* **133**, e161944. doi:10.1172/JCI161944

Bell, S. L., Lopez, K. L., Cox, J. S., Patrick, K. L. and Watson, R. O. (2021). Galectin-8 senses phagosomal damage and recruits selective autophagy adapter TAX1BP1 to control *Mycobacterium tuberculosis* infection in macrophages. *mBio* **12**, e0187120.

Bernard, E. M., Fearn, A., Bussi, C., Santucci, P., Peddie, C. J., Lai, R. J., Collinson, L. M. and Gutierrez, M. G. (2021). *M. tuberculosis* infection of human iPSC-derived macrophages reveals complex membrane dynamics during xenophagy evasion. *J. Cell Sci.* **134**, jcs252973. doi:10.1242/jcs.252973

Bhattacharyya, M., Dhar, R., Basu, S., Das, A., Reynolds, D. M. and Dutta, T. K. (2023). Molecular evaluation of the metabolism of estrogenic di(2-ethylhexyl) phthalate in *Mycobacterium* sp. *Microb. Cell Fact.* **22**, 82. doi:10.1186/s12934-023-02096-0

Block, A. M., Namugenyi, S. B., Palani, N. P., Brokaw, A. M., Zhang, L., Beckman, K. B. and Tischler, A. D. (2023). *Mycobacterium tuberculosis* requires the outer membrane lipid phthiocerol dimycocerosate for starvation-induced antibiotic tolerance. *mSystems* **8**, e0069922. doi:10.1128/mSystems.00699-22

Bo, H., Moure, U. A. E., Yang, Y., Pan, J., Li, L., Wang, M., Ke, X. and Cui, H. (2023). *Mycobacterium tuberculosis*-macrophage interaction: molecular updates. *Front. Cell Infect. Microbiol.* **13**, 1062963. doi:10.3389/fcimb.2023.1062963

Bodinier, R., Leiba, J., Sabra, A., Jauslin, T. N., Lamrabet, O., Guilhen, C., Marchetti, A., Iwade, Y., Kawata, T., Lima, W. C. et al. (2020). LrrkA, a kinase with leucine-rich repeats, links folate sensing with Kif2 activity and intracellular killing. *Cell. Microbiol.* **22**, e13129. doi:10.1111/cmi.13129

Bodinier, R., Sabra, A., Leiba, J., Marchetti, A., Lamrabet, O., Ayadi, I., Filić, V., Kawata, T., Weber, I. and Cosson, P. (2021). Role of LrrkA in the control of phagocytosis and cell motility in *Dictyostelium discoideum*. *Front. Cell Dev. Biol.* **9**, 629200. doi:10.3389/fcell.2021.629200

Bosmani, C., Perret, A., Leuba, F., Guého, A. and Hanna, N. (2021). Disruption of vacuolin microdomains in the host *Dictyostelium discoideum* 1 increases resistance to *Mycobacterium marinum*-induced membrane 2 damage and infection 3. *bioRxiv* 2021.11.16.468763. doi:10.1101/2021.11.16.468763

Botella, H., Peyron, P., Levillain, F., Poincloux, R., Poquet, Y., Brandli, I., Wang, C., Tailleux, L., Tilleul, S., Charriere, G. M. et al. (2011). Mycobacterial p(1)-type ATPases mediate resistance to zinc poisoning in human macrophages. *Cell Host Microbe* **10**, 248–259. doi:10.1016/j.chom.2011.08.006

Botella, H., Stadthagen, G., Lugo-Villarino, G., De Chastellier, C. and Neyrolles, O. (2012). Metallobiology of host-pathogen interactions: an intoxicating new insight. *Trends Microbiol.* **20**, 106–112. doi:10.1016/j.tim.2012.01.005

Boudehen, Y. M., Faucher, M., Marechal, X., Miras, R., Rech, J., Rombouts, Y., Seneque, O., Wallat, M., Demange, P., Bouet, J. Y. et al. (2022). Mycobacterial resistance to zinc poisoning requires assembly of P-ATPase-containing membrane metal efflux platforms. *Nat. Commun.* **13**, 4731. doi:10.1038/s41467-022-32085-7

Boulais, J., Trost, M., Landry, C. R., Dieckmann, R., Levy, E. D., Soldati, T., Michnick, S. W., Thibault, P. and Desjardins, M. (2010). Molecular characterization of the evolution of phagosomes. *Mol. Syst. Biol.* **6**, 423. doi:10.1038/msb.2010.80

Brenz, Y., Ohnzeit, D., Winther-Larsen, H. C. and Hagedorn, M. (2017). Nramp1 and NrampB contribute to resistance against *Francisella* in *Dictyostelium*. *Front. Cell Infect. Microbiol.* **7**, 282. doi:10.3389/fcimb.2017.00282

Bretschneider, T., Othmer, H. G. and Weijer, C. J. (2016). Progress and perspectives in signal transduction, actin dynamics, and movement at the cell and tissue level: lessons from *Dictyostelium*. *Interface Focus* **6**, 20160047. doi:10.1098/rsfs.2016.0047

Brock, D. A., Callison, W. E., Strassmann, J. E. and Queller, D. C. (2016). Sentinel cells, symbiotic bacteria and toxin resistance in the social amoeba *Dictyostelium discoideum*. *Proc. Biol. Sci.* **283**, 20152727. doi:10.1098/rspb.2015.2727

Brock, D. A., Haselkorn, T. S., Garcia, J. R., Bashir, U., Douglas, T. E., Galloway, J., Brodie, F., Queller, D. C. and Strassmann, J. E. (2018). Diversity of free-living environmental bacteria and their interactions with a bacterivorous amoeba. *Front. Cell Infect. Microbiol.* **8**, 411. doi:10.3389/fcimb.2018.00411

Buckley, C. M., Pots, H., Gueho, A., Vines, J. H., Munn, C. J., Phillips, B. A., Gilsbach, B., Traynor, D., Nikolaev, A., Soldati, T. et al. (2020). Coordinated Ras and Rac activity shapes macropinocytic cups and enables phagocytosis of geometrically diverse bacteria. *Curr. Biol.* **30**, 2912–2926.e5. doi:10.1016/j.cub.2020.05.049

Buracco, S., Peracino, B., Andreini, C., Bracco, E. and Bozzaro, S. (2018). Differential effects of iron, zinc, and copper on *Dictyostelium discoideum* cell growth and resistance to *Legionella pneumophila*. *Front. Cell Infect. Microbiol.* **7**, 536. doi:10.3389/fcimb.2017.00536

Bussi, C., Heunis, T., Pellegrino, E., Bernard, E. M., Bah, N., Dos Santos, M. S., Santucci, P., Aylan, B., Rodgers, A., Fearn, A. et al. (2022). Lysosomal

- damage drives mitochondrial proteome remodelling and reprograms macrophage immunometabolism. *Nat. Commun.* **13**, 7338. doi:10.1038/s41467-022-34632-8
- Bussi, C., Mangiarotti, A., Vanhille-Campos, C., Aylan, B., Pellegrino, E., Athanasiadi, N., Fearn, A., Rodgers, A., Franzmann, T. M., Saric, A. et al. (2023). Stress granules plug and stabilize damaged endolysosomal membranes. *Nature* **623**, 1062-1069. doi:10.1038/s41586-023-06726-w
- Buter, J., Cheng, T. Y., Ghanem, M., Grootemaat, A. E., Raman, S., Feng, X., Plantijn, A. R., Ennis, T., Wang, J., Cotton, R. N. et al. (2019). *Mycobacterium tuberculosis* releases an antacid that remodels phagosomes. *Nat. Chem. Biol.* **15**, 889-899. doi:10.1038/s41589-019-0336-0
- Butler, R. E., Smith, A. A., Mendum, T. A., Chandran, A., Wu, H., Lefrancois, L., Chambers, M., Soldati, T. and Stewart, G. R. (2020). *Mycobacterium bovis* uses the ESX-1 Type VII secretion system to escape predation by the soil-dwelling amoeba *Dictyostelium discoideum*. *ISME J.* **14**, 919-930. doi:10.1038/s41396-019-0572-z
- Calvo-Garrido, J., Carilla-Latorre, S., Kubohara, Y., Santos-Rodrigo, N., Mesquita, A., Soldati, T., Golstein, P. and Escalante, R. (2010). Autophagy in *Dictyostelium*: genes and pathways, cell death and infection. *Autophagy* **6**, 686-701. doi:10.4161/auto.6.6.12513
- Cambier, C. J., Banik, S. M., Buonomo, J. A. and Bertozzi, C. R. (2020). Spreading of a mycobacterial cell-surface lipid into host epithelial membranes promotes infectivity. *Elife* **9**, e60648. doi:10.7554/eLife.60648
- Campo-Perez, V., Gualar-Garrido, S., Luquin, M., Sanchez-Chardi, A. and Julian, E. (2022). The high plasticity of nonpathogenic *Mycobacterium brumae* induces rapid changes in its lipid profile during pellicle maturation: the potential of this bacterium as a versatile cell factory for lipid compounds of therapeutic interest. *Int. J. Mol. Sci.* **23**, 13609. doi:10.3390/ijms232113609
- Cardenal-Muñoz, E., Arafah, S., López-Jiménez, A. T., Kicka, S., Falaise, A., Bach, F., Schaad, O., King, J. S., Hagedorn, M. and Soldati, T. (2017). *Mycobacterium marinum* antagonistically induces an autophagic response while repressing the autophagic flux in a TORC1- and ESX-1-dependent manner. *PLoS Pathog.* **13**, e1006344. doi:10.1371/journal.ppat.1006344
- Cardenal-Munoz, E., Barisch, C., Lefrancois, L. H., Lopez-Jimenez, A. T. and Soldati, T. (2018). When Dicty met Myco, a (not so) romantic story about one amoeba and its intracellular pathogen. *Front. Cell Infect. Microbiol.* **7**, 529. doi:10.3389/fcimb.2017.00529
- Castillo, E. F., Dekonenko, A., Arko-Mensah, J., Mandell, M. A., Dupont, N., Jiang, S., Delgado-Vargas, M., Timmins, G. S., Bhattacharya, D., Yang, H. et al. (2012). Autophagy protects against active tuberculosis by suppressing bacterial burden and inflammation. *Proc. Natl. Acad. Sci. USA* **109**, E3168-E3176. doi:10.1073/pnas.1210500109
- Chai, Q., Wang, L., Liu, C. H. and Ge, B. (2020). New insights into the evasion of host innate immunity by *Mycobacterium tuberculosis*. *Cell. Mol. Immunol.* **17**, 901-913. doi:10.1038/s41423-020-0502-z
- Chandra, P., He, L., Zimmerman, M., Yang, G., Koster, S., Ouimet, M., Wang, H., Moore, K. J., Dartois, V., Schilling, J. D. et al. (2020). Inhibition of fatty acid oxidation promotes macrophage control of *Mycobacterium tuberculosis*. *mBio* **11**, e01139-20. doi:10.1128/mBio.01139-20
- Chandra, P., Grigsby, S. J. and Philips, J. A. (2022). Immune evasion and provocation by *Mycobacterium tuberculosis*. *Nat. Rev. Microbiol.* **20**, 750-766. doi:10.1038/s41579-022-00763-4
- Chauhan, S., Kumar, S., Jain, A., Ponpuak, M., Mudd, M. H., Kimura, T., Choi, S. W., Peters, R., Mandell, M., Bruun, J. A. et al. (2016). TRIMs and galectins globally cooperate and TRIM16 and galectin-3 co-direct autophagy in endomembrane damage homeostasis. *Dev. Cell* **39**, 13-27. doi:10.1016/j.devcel.2016.08.003
- Chen, G., Zhuchenko, O. and Kuspa, A. (2007). Immune-like phagocyte activity in the social amoeba. *Science* **317**, 678-681. doi:10.1126/science.1143991
- Chen, Y., Cao, S., Sun, Y. and Li, C. (2018). Gene expression profiling of the TRIM protein family reveals potential biomarkers for indicating tuberculosis status. *Microb. Pathog.* **114**, 385-392. doi:10.1016/j.micpath.2017.12.008
- Chen, L., Li, X., Xu, P. and He, Z. G. (2022). A novel zinc exporter CtpG enhances resistance to zinc toxicity and survival in *Mycobacterium bovis*. *Microbiol. Spectr.* **10**, e0145621. doi:10.1128/spectrum.01456-21
- Chiaradia, L., Lefebvre, C., Parra, J., Marcoux, J., Bulet-Schiltz, O., Etienne, G., Tropis, M. and Daffe, M. (2017). Dissecting the mycobacterial cell envelope and defining the composition of the native mycomembrane. *Sci. Rep.* **7**, 12807. doi:10.1038/s41598-017-12718-4
- Chung, J., Torta, F., Masai, K., Lucast, L., Czapl, H., Tanner, L. B., Narayanaswamy, P., Wenk, M. R., Nakatsu, F. and De Camilli, P. (2015). PI4P/phosphatidylserine countertransport at ORP5- and ORP8-mediated ER-plasma membrane contacts. *Science* **349**, 428-432. doi:10.1126/science.aab1370
- Collins, C. A., De Maziere, A., Van Dijk, S., Carlsson, F., Klumperman, J. and Brown, E. J. (2009). Atg5-independent sequestration of ubiquitinated mycobacteria. *PLoS Pathog.* **5**, e1000430. doi:10.1371/journal.ppat.1000430
- Conrad, W. H., Osman, M. M., Shanahan, J. K., Chu, F., Takaki, K. K., Cameron, J., Hopkinson-Woolley, D., Brosch, R. and Ramakrishnan, L. (2017). Mycobacterial ESX-1 secretion system mediates host cell lysis through bacterium contact-dependent gross membrane disruptions. *Proc. Natl. Acad. Sci. USA* **114**, 1371-1376. doi:10.1073/pnas.1620133114
- Cooper, S. T. and McNeil, P. L. (2015). Membrane repair: mechanisms and pathophysiology. *Physiol. Rev.* **95**, 1205-1240. doi:10.1152/physrev.00037.2014
- Cosson, P. and Soldati, T. (2008). Eat, kill or die: when amoeba meets bacteria. *Curr. Opin. Microbiol.* **11**, 271-276. doi:10.1016/j.mib.2008.05.005
- Crespo-Yanez, X., Oddy, J., Lamrabet, O., Jauslin, T., Marchetti, A. and Cosson, P. (2023). Sequential action of antibacterial effectors in *Dictyostelium discoideum* phagosomes. *Mol. Microbiol.* **119**, 74-85. doi:10.1111/mmi.15004
- Dan Dunn, J., Alvarez, L. A., Zhang, X. and Soldati, T. (2015). Reactive oxygen species and mitochondria: a nexus of cellular homeostasis. *Redox Biol.* **6**, 472-485. doi:10.1016/j.redox.2015.09.005
- Darwin, K. H. (2015). *Mycobacterium tuberculosis* and copper: a newly appreciated defense against an old foe? *J. Biol. Chem.* **290**, 18962-18966. doi:10.1074/jbc.R115.640193
- Davis, J. M., Clay, H., Lewis, J. L., Ghori, N., Herbomel, P. and Ramakrishnan, L. (2002). Real-time visualization of mycobacterium-macrophage interactions leading to initiation of granuloma formation in zebrafish embryos. *Immunity* **17**, 693-702. doi:10.1016/S1074-7613(02)00475-2
- Deretic, V. and Wang, F. (2023). Autophagy is part of the answer to tuberculosis. *Nat. Microbiol.* **8**, 762-763. doi:10.1038/s41564-023-01373-3
- Dickson, E. J. and Hille, B. (2019). Understanding phosphoinositides: rare, dynamic, and essential membrane phospholipids. *Biochem. J.* **476**, 1-23. doi:10.1042/BCJ20180022
- Dormann, D., Weijer, G., Dowler, S. and Weijer, C. J. (2004). In vivo analysis of 3-phosphoinositide dynamics during *Dictyostelium* phagocytosis and chemotaxis. *J. Cell Sci.* **117**, 6497-6509. doi:10.1242/jcs.01579
- Du, X., Barisch, C., Paschke, P., Herrfurth, C., Bertinetti, O., Pawolleck, N., Otto, H., Ruhling, H., Feussner, I., Herberg, F. W. et al. (2013). *Dictyostelium* lipid droplets host novel proteins. *Eukaryot. Cell* **12**, 1517-1529. doi:10.1128/EC.00182-13
- Duan, L., Yi, M., Chen, J., Li, S. and Chen, W. (2016). *Mycobacterium tuberculosis* EIS gene inhibits macrophage autophagy through up-regulation of IL-10 by increasing the acetylation of histone H3. *Biochem. Biophys. Res. Commun.* **473**, 1229-1234. doi:10.1016/j.bbrc.2016.04.045
- Dunn, J. D., Bosmani, C., Barisch, C., Raykov, L., Lefrancois, L. H., Cardenal-Munoz, E., Lopez-Jimenez, A. T. and Soldati, T. (2018). Eat prey, live: *Dictyostelium discoideum* as a model for cell-autonomous defenses. *Front. Immunol.* **8**, 1906. doi:10.3389/fimmu.2017.01906
- Dutta, A., Mukku, R. P., Kumar, G. A., Jafurulla, M., Raghunand, T. R. and Chattopadhyay, A. (2022). Integrity of the actin cytoskeleton of host macrophages is necessary for mycobacterial entry. *J. Membr. Biol.* **255**, 623-632. doi:10.1007/s00232-022-00217-1
- Edwards, K. M., Cynamon, M. H., Voladri, R. K., Hager, C. C., Destefano, M. S., Tham, K. T., Lakey, D. L., Bochan, M. R. and Kernodle, D. S. (2001). Iron-cofactored superoxide dismutase inhibits responses to *Mycobacterium tuberculosis*. *Am. J. Respir. Crit. Care. Med.* **164**, 2213-2219. doi:10.1164/ajrccm.164.12.2106093
- Egami, Y. and Araki, N. (2012). Rab20 regulates phagosome maturation in RAW264 macrophages during Fc gamma receptor-mediated phagocytosis. *PLoS One* **7**, e35663. doi:10.1371/journal.pone.0035663
- Eichinger, L., Pachebat, J. A., Glockner, G., Rajandream, M. A., Sugang, R., Berriman, M., Song, J., Olsen, R., Szafranski, K., Xu, Q. et al. (2005). The genome of the social amoeba *Dictyostelium discoideum*. *Nature* **435**, 43-57. doi:10.1038/nature03481
- El Chemaly, A. and Demareux, N. (2012). Do Hv1 proton channels regulate the ionic and redox homeostasis of phagosomes? *Mol. Cell. Endocrinol.* **353**, 82-87. doi:10.1016/j.mce.2011.10.005
- Esterhuysen, M. M., Weiner, J., 3rd, Caron, E., Loxton, A. G., Iannaccone, M., Wagman, C., Saikali, P., Stanley, K., Wolski, W. E., Mollenkopf, H. J. et al. (2015). Epigenetics and proteomics join transcriptomics in the quest for tuberculosis biomarkers. *mBio* **6**, e01187-e01115. doi:10.1128/mBio.01187-15
- Farinholt, T., Dinh, C. and Kuspa, A. (2019). Microbiome management in the social amoeba *Dictyostelium discoideum* compared to humans. *Int. J. Dev. Biol.* **63**, 447-450. doi:10.1387/ijdb.190240ak
- Feng, S., Mcnehan, M. E., Kinsella, R. L., Sur Chowdhury, C., Chavez, S. M., Naik, S. K., McKee, S. R., Van Winkle, J. A., Dubey, N., Samuels, A. et al. (2024). Autophagy promotes efficient T cell responses to restrict high-dose *Mycobacterium tuberculosis* infection in mice. *Nat. Microbiol.* **9**, 684-697. doi:10.1038/s41564-024-01608-x
- Ferling, I., Dunn, J. D., Ferling, A., Soldati, T. and Hillmann, F. (2020). Conidial melanin of the human-pathogenic fungus *Aspergillus fumigatus* disrupts cell autonomous defenses in amoebae. *mBio* **11**, e00862-20. doi:10.1128/mBio.00862-20
- Fines, D. M., Schichnes, D., Knight, M., Anaya-Sanchez, A., Thuong, N., Cox, J. and Stanley, S. A. (2023). Mycobacterial formation of intracellular lipid inclusions is a dynamic process associated with rapid replication. *bioRxiv* 2023.08.10.552809. doi:10.1101/2023.08.10.552809

- Footte, J. R., Patel, A. A., Yona, S. and Segal, A. W. (2019). Variations in the phagosomal environment of human neutrophils and mononuclear phagocyte subsets. *Front. Immunol.* **10**, 188. doi:10.3389/fimmu.2019.00188
- Foulon, M., Listian, S. A., Soldati, T. and Barisch, C. (2022). Chapter 6 - Conserved mechanisms drive host-lipid access, import, and utilization in *Mycobacterium tuberculosis* and *M. marinum*. Academic Press. doi:10.1016/B978-0-323-91948-7.00011-7
- Franco, L. H., Nair, V. R., Scharn, C. R., Xavier, R. J., Torrealba, J. R., Shiloh, M. U. and Levine, B. (2017). The ubiquitin ligase Smurf1 functions in selective autophagy of *Mycobacterium tuberculosis* and anti-tuberculous host defense. *Cell Host Microbe* **21**, 59-72. doi:10.1016/j.chom.2016.11.002
- Garcia-Bengoa, M., Meurer, M., Goethe, R., Singh, M., Reljic, R. and Von Kockritz-Blickwede, M. (2023). Role of phagocyte extracellular traps during *Mycobacterium tuberculosis* infections and tuberculosis disease processes. *Front. Microbiol.* **14**, 983299. doi:10.3389/fmicb.2023.983299
- Ge, P., Lei, Z., Yu, Y., Lu, Z., Qiang, L., Chai, Q., Zhang, Y., Zhao, D., Li, B., Pang, Y. et al. (2022). *M. tuberculosis* PknG manipulates host autophagy flux to promote pathogen intracellular survival. *Autophagy* **18**, 576-594. doi:10.1080/15548627.2021.1938912
- Gerstenmaier, L., Pilla, R., Herrmann, L., Herrmann, H., Prado, M., Villafano, G. J., Kolonko, M., Reimer, R., Soldati, T., King, J. S. et al. (2015). The autophagic machinery ensures nonlytic transmission of mycobacteria. *Proc. Natl. Acad. Sci. USA* **112**, E687-E692. doi:10.1073/pnas.1423318112
- Gerstenmaier, L., Colasanti, O., Behrens, H., Kolonko, M., Hammann, C. and Hagedorn, M. (2024). Recruitment of both the ESCRT and autophagic machineries to ejecting *Mycobacterium marinum*. *Mol. Microbiol.* **121**, 385-393. doi:10.1111/mmi.15075
- Gharun, K., Senges, J., Seidl, M., Losslein, A., Kolter, J., Lohrmann, F., Fliegau, M., Elgizouli, M., Alber, M., Vavra, M. et al. (2017). Mycobacteria exploit nitric oxide-induced transformation of macrophages into permissive giant cells. *EMBO Rep.* **18**, 2144-2159. doi:10.15252/embr.201744121
- Gill, S. E. and Chain, F. J. J. (2023). Very low rates of spontaneous gene deletions and gene duplications in *Dictyostelium discoideum*. *J. Mol. Evol.* **91**, 24-32. doi:10.1007/s00239-022-10081-1
- Gilmore, S. A., Schelle, M. W., Holsclaw, C. M., Leigh, C. D., Jain, M., Cox, J. S., Leary, J. A. and Bertozzi, C. R. (2012). Sulfolipid-1 biosynthesis restricts *Mycobacterium tuberculosis* growth in human macrophages. *ACS Chem. Biol.* **7**, 863-870. doi:10.1021/cb200311s
- Golovkine, G. R., Roberts, A. W., Morrison, H. M., Rivera-Lugo, R., McCall, R. M., Nilsson, H., Garellis, H. E., Repasy, T., Cronce, M., Budzik, J. et al. (2023). Autophagy restricts *Mycobacterium tuberculosis* during acute infection in mice. *Nat. Microbiol.* **8**, 819-832. doi:10.1038/s41564-023-01354-6
- Goyal, S., Klassert, T. E. and Slevogt, H. (2016). C-type lectin receptors in tuberculosis: what we know. *Med. Microbiol. Immunol.* **205**, 513-535. doi:10.1007/s00430-016-0470-1
- Gronski, M. A., Kinchen, J. M., Juncadella, I. J., Franc, N. C. and Ravichandran, K. S. (2009). An essential role for calcium flux in phagocytes for apoptotic cell engulfment and the anti-inflammatory response. *Cell Death Differ.* **16**, 1323-1331. doi:10.1038/cdd.2009.55
- Gu, W., Zhang, J., Li, Q., Zhang, Y., Lin, X., Wu, B., Yin, Q., Sun, J., Lu, Y., Sun, X. et al. (2023). The TRIM37 variants in Mulibrey nanism patients paralyze follicular helper T cell differentiation. *Cell Discov.* **9**, 82. doi:10.1038/s41421-023-00561-z
- Guallar-Garrido, S., Luquin, M. and Julian, E. (2021). Analysis of the lipid composition of mycobacteria by thin layer chromatography. *J. Vis. Exp.* **170**, e62368. doi:10.3791/62368
- Guallar-Garrido, S., Campo-Pérez, V., Pérez-Trujillo, M., Cabrera, C., Sensenich, J., Sánchez-Chardi, A., Rabanal, R. M., Gómez-Mora, E., Noguera-Ortega, E., Luquin, M. et al. (2022). Mycobacterial surface characters remodeled by growth conditions drive different tumor-infiltrating cells and systemic IFN- γ /IL-17 release in bladder cancer treatment. *Oncoimmunology* **11**, 2051845. doi:10.1080/2162402X.2022.2051845
- Guého, A., Bosmani, C., Nitschke, J. and Soldati, T. (2023). Proteomic characterization of the *Mycobacterium marinum*-containing vacuole in *Dictyostelium discoideum*. *bioRxiv* 592717. doi:10.1101/592717
- Guerin, I. and De Chastellier, C. (2000). Pathogenic mycobacteria disrupt the macrophage actin filament network. *Infect. Immun.* **68**, 2655-2662. doi:10.1128/IAI.68.5.2655-2662.2000
- Gunther, J. and Seyfert, H. M. (2018). The first line of defence: insights into mechanisms and relevance of phagocytosis in epithelial cells. *Semin. Immunopathol.* **40**, 555-565. doi:10.1007/s00281-018-0701-1
- Gutierrez, M. G. and Enninga, J. (2022). Intracellular niche switching as host subversion strategy of bacterial pathogens. *Curr. Opin. Cell Biol.* **76**, 102081. doi:10.1016/j.ccb.2022.102081
- Gutierrez, M. G., Master, S. S., Singh, S. B., Taylor, G. A., Colombo, M. I. and Deretic, V. (2004). Autophagy is a defense mechanism inhibiting BCG and *Mycobacterium tuberculosis* survival in infected macrophages. *Cell* **119**, 753-766. doi:10.1016/j.cell.2004.11.038
- Hagedorn, M., Rohde, K. H., Russell, D. G. and Soldati, T. (2009). Infection by tubercular mycobacteria is spread by nonlytic ejection from their amoeba hosts. *Science* **323**, 1729-1733. doi:10.1126/science.1169381
- Hanna, N., Koliwer-Brandl, H., Lefrancois, L. H., Kalinina, V., Cardenal-Munoz, E., Appiah, J., Leuba, F., Gueho, A., Hilbi, H., Soldati, T. et al. (2021). Zn²⁺ intoxication of *Mycobacterium marinum* during *Dictyostelium discoideum* infection is counteracted by induction of the pathogen Zn²⁺ exporter CtpC. *mBio* **12**, e01313-20. doi:10.1128/mBio.01313-20
- Haselkorn, T. S., Disalvo, S., Miller, J. W., Bashir, U., Brock, D. A., Queller, D. C. and Strassmann, J. E. (2019). The specificity of *Burkholderia* symbionts in the social amoeba farming symbiosis: prevalence, species, genetic and phenotypic diversity. *Mol. Ecol.* **28**, 847-862. doi:10.1111/mec.14982
- Herbst, S., Campbell, P., Harvey, J., Bernard, E. M., Papayannopoulos, V., Wood, N. W., Morris, H. R. and Gutierrez, M. G. (2020). LRRK 2 activation controls the repair of damaged endomembranes in macrophages. *EMBO J.* **39**, e104494. doi:10.15252/emboj.2020104494
- Hoffmann, C., Finsel, I., Otto, A., Pfaffinger, G., Rothmeier, E., Hecker, M., Becher, D. and Hilbi, H. (2014). Functional analysis of novel Rab GTPases identified in the proteome of purified *Legionella*-containing vacuoles from macrophages. *Cell. Microbiol.* **16**, 1034-1052. doi:10.1111/cmi.12235
- Hoffpauir, C. T., Bell, S. L., West, K. O., Jing, T., Wagner, A. R., Torres-Odio, S., Cox, J. S., West, A. P., Li, P., Patrick, K. L. et al. (2020). TRIM14 is a key regulator of the type I IFN response during *Mycobacterium tuberculosis* infection. *J. Immunol.* **205**, 153-167. doi:10.4049/jimmunol.1901511
- Hossain, M. M. and Norazmi, M. N. (2013). Pattern recognition receptors and cytokines in *Mycobacterium tuberculosis* infection—the double-edged sword? *Biomed. Res. Int.* **2013**, 179174. doi:10.1155/2013/179174
- Huang, D. and Bao, L. (2016). *Mycobacterium tuberculosis* EspB protein suppresses interferon- γ -induced autophagy in murine macrophages. *J. Microbiol. Immunol. Infect.* **49**, 859-865. doi:10.1016/j.jmii.2014.11.008
- Huang, F., Cai, F., Dahabieh, M. S., Gunawardena, K., Talebi, A., Dehairs, J., El-Turk, F., Park, J. Y., Li, M., Goncalves, C. et al. (2023). Peroxisome disruption alters lipid metabolism and potentiates antitumor response with MAPK-targeted therapy in melanoma. *J. Clin. Invest.* **133**, e166644. doi:10.1172/JCI166644
- Hüsler, D., Stauffer, P., Keller, B., Böck, D., Steiner, T., Ostrzinski, A., Vormittag, S., Striednig, B., Swart, A. L., Letourneur, F. et al. (2023). The large GTPase Sey1/atlastin mediates lipid droplet- and FadL-dependent intracellular fatty acid metabolism of *Legionella pneumophila*. *Elife* **12**, e85142. doi:10.7554/eLife.85142
- Iantomasi, R., Sali, M., Cascioferro, A., Palucci, I., Zumbo, A., Soldini, S., Rocca, S., Greco, E., Maulucci, G., De Spirito, M. et al. (2012). PE_{PGRS30} is required for the full virulence of *Mycobacterium tuberculosis*. *Cell. Microbiol.* **14**, 356-367. doi:10.1111/j.1462-5822.2011.01721.x
- Ifrid, E., Ouertatani-Sakouhi, H., Jauslin, T., Kicka, S., Chiriano, G., Harrison, C. F., Hilbi, H., Scapozza, L., Soldati, T. and Cosson, P. (2022). 5-ethyl-2'-deoxyuridine fragilizes *Klebsiella pneumoniae* outer wall and facilitates intracellular killing by phagocytic cells. *PLoS One* **17**, e0269093. doi:10.1371/journal.pone.0269093
- Ishikawa, E., Ishikawa, T., Morita, Y. S., Toyonaga, K., Yamada, H., Takeuchi, O., Kinoshita, T., Akira, S., Yoshikai, Y. and Yamasaki, S. (2009). Direct recognition of the mycobacterial glycolipid, trehalose dimycolate, by C-type lectin Mincle. *J. Exp. Med.* **206**, 2879-2888. doi:10.1084/jem.20091750
- Jamaati, H., Mortaz, E., Pajouhi, Z., Folkerts, G., Movassaghi, M., Moloudizargari, M., Adcock, I. M. and Garssen, J. (2017). Nitric oxide in the pathogenesis and treatment of tuberculosis. *Front. Microbiol.* **8**, 2008. doi:10.3389/fmicb.2017.02008
- Jani, C., Marsh, A., Uchil, P., Jain, N., Baskir, Z. R., Glover, O. T., Root, D. E., Doench, J. G. and Barczak, A. K. (2023). Vps18 contributes to phagosome membrane integrity in *Mycobacterium tuberculosis*-infected macrophages. *bioRxiv* 2023.10.01.560397. doi:10.1101/2023.10.01.560397
- Javed, R., Jain, A., Duque, T., Hendrix, E., Paddar, M. A., Khan, S., Claude-Taupin, A., Jia, J., Allers, L., Wang, F. et al. (2023). Mammalian ATG8 proteins maintain autophagosomal membrane integrity through ESCRTs. *EMBO J.* **42**, e112845. doi:10.15252/emboj.2022112845
- Jayachandran, R., Sundaramurthy, V., Combaluzier, B., Mueller, P., Korf, H., Huygen, K., Miyazaki, T., Albrecht, I., Massner, J. and Pieters, J. (2007). Survival of mycobacteria in macrophages is mediated by coronin 1-dependent activation of calcineurin. *Cell* **130**, 37-50. doi:10.1016/j.cell.2007.04.043
- Jia, J., Wang, F., Bhujabal, Z., Peters, R., Mudd, M., Duque, T., Allers, L., Javed, R., Salemi, M., Behrends, C. et al. (2022). Stress granules and mTOR are regulated by membrane atg8ylation during lysosomal damage. *J. Cell Biol.* **221**, e202207091. doi:10.1083/jcb.202207091
- Jung, J. Y., Madan-Lala, R., Georgieva, M., Rengarajan, J., Sohaskey, C. D., Bange, F. C. and Robinson, C. M. (2013). The intracellular environment of human macrophages that produce nitric oxide promotes growth of mycobacteria. *Infect. Immun.* **81**, 3198-3209. doi:10.1128/IAI.00611-13
- Kim, B. H., Shenoy, A. R., Kumar, P., Bradfield, C. J. and Macmicking, J. D. (2012). IFN-inducible GTPases in host cell defense. *Cell Host Microbe* **12**, 432-444. doi:10.1016/j.chom.2012.09.007
- Kim, J. Y., Kim, M. J., Lee, J. S., Son, J., Kim, D. H., Lee, J. S., Jeong, S. K., Chun, E. and Lee, K. Y. (2022a). Stratifin (SFN) regulates lung cancer progression via nucleating the Vps34-BECN1-TRAF6 complex for autophagy induction. *Clin. Transl. Med.* **12**, e896. doi:10.1002/ctm2.896

- Kim, S. H., Baek, S. I., Jung, J., Lee, E. S., Na, Y., Hwang, B. Y., Roh, Y. S., Hong, J. T., Han, S. B. and Kim, Y. (2022b). Chemical inhibition of TRAF6-TAK1 axis as therapeutic strategy of endotoxin-induced liver disease. *Biomed. Pharmacother.* **155**, 113688. doi:10.1016/j.biopha.2022.113688
- Kimmey, J. M., Huynh, J. P., Weiss, L. A., Park, S., Kambal, A., Debnath, J., Virgin, H. W. and Stallings, C. L. (2015). Unique role for ATG5 in neutrophil-mediated immunopathology during *M. tuberculosis* infection. *Nature* **528**, 565-569. doi:10.1038/nature16451
- Kin, K., Chen, Z. H., Forbes, G. and Schaap, P. (2022). Evolution of a novel cell type in *Dictyostelia* required gene duplication of a cudA-like transcription factor. *Curr. Biol.* **32**, 428-437.e4. doi:10.1016/j.cub.2021.11.047
- Kinsella, R. L., Kimmey, J. M., Smirnov, A., Woodson, R., Gaggioli, M. R., Chavez, S. M., Kreamealmeyer, D. and Stallings, C. L. (2023). Autophagy prevents early proinflammatory responses and neutrophil recruitment during *Mycobacterium tuberculosis* infection without affecting pathogen burden in macrophages. *PLoS Biol.* **21**, e3002159. doi:10.1371/journal.pbio.3002159
- Kjellin, J., Pranting, M., Bach, F., Vaid, R., Edelbroek, B., Li, Z., Hoepfner, M. P., Grabherr, M., Isberg, R. R., Hagedorn, M. et al. (2019). Investigation of the host transcriptional response to intracellular bacterial infection using *Dictyostelium discoideum* as a host model. *BMC Genomics* **20**, 961. doi:10.1186/s12864-019-6269-x
- Knobloch, P., Koliwer-Brandl, H., Arnold, F. M., Hanna, N., Gonda, I., Adenau, S., Personnic, N., Barisch, C., Seeger, M. A., Soldati, T. et al. (2020). *Mycobacterium marinum* produces distinct mycobactin and carboxymycobactin siderophores to promote growth in broth and phagocytes. *Cell. Microbiol.* **22**, e13163. doi:10.1111/cmi.13163
- Koliwer-Brandl, H., Knobloch, P., Barisch, C., Welin, A., Hanna, N., Soldati, T. and Hilbi, H. (2019). Distinct *Mycobacterium marinum* phosphatases determine pathogen vacuole phosphoinositide pattern, phagosome maturation, and escape to the cytosol. *Cell. Microbiol.* **21**, e13008. doi:10.1111/cmi.13008
- Kolonko, M., Geffken, A. C., Blumer, T., Hagens, K., Schaible, U. E. and Hagedorn, M. (2014). WASH-driven actin polymerization is required for efficient mycobacterial phagosome maturation arrest. *Cell. Microbiol.* **16**, 232-246. doi:10.1111/cmi.12217
- Koster, S., Upadhyay, S., Chandra, P., Papavinasundaram, K., Yang, G., Hassan, A., Grigsby, S. J., Mittal, E., Park, H. S., Jones, V. et al. (2017). *Mycobacterium tuberculosis* is protected from NADPH oxidase and LC3-associated phagocytosis by the LCP protein CpsA. *Proc. Natl. Acad. Sci. USA* **114**, E8711-E8720. doi:10.1073/pnas.1707792114
- Kreibich, S., Emmenlauer, M., Fredlund, J., Ramo, P., Munz, C., Dehio, C., Enninga, J. and Hardt, W. D. (2015). Autophagy proteins promote repair of endosomal membranes damaged by the salmonella type three secretion system 1. *Cell Host Microbe* **18**, 527-537. doi:10.1016/j.chom.2015.10.015
- Landstrom, M. (2010). The TAK1-TRAF6 signalling pathway. *Int. J. Biochem. Cell Biol.* **42**, 585-589. doi:10.1016/j.biocel.2009.12.023
- Lapaquette, P., Ducreux, A., Basmaciyan, L., Paradis, T., Bon, F., Bataille, A., Winckler, P., Hube, B., D'entfert, C., Escalante, A. et al. (2022). Membrane protective role of autophagic machinery during infection of epithelial cells by *Candida albicans*. *Gut Microbes* **14**, 2004798. doi:10.1080/19490976.2021.2004798
- Lawe, D. C., Chawla, A., Merithew, E., Dumas, J., Carrington, W., Fogarty, K., Lifshitz, L., Tuft, R., Lambright, D. and Corvera, S. (2002). Sequential roles for phosphatidylinositol 3-phosphate and Rab5 in tethering and fusion of early endosomes via their interaction with EEA1. *J. Biol. Chem.* **277**, 8611-8617. doi:10.1074/jbc.M109239200
- Lee, J., Remold, H. G., Jeong, M. H. and Kornfeld, H. (2006). Macrophage apoptosis in response to high intracellular burden of *Mycobacterium tuberculosis* is mediated by a novel caspase-independent pathway. *J. Immunol.* **176**, 4267-4274. doi:10.4049/jimmunol.176.7.4267
- Lefrançois, L. H., Nitschke, J., Wu, H., Panis, G., Prados, J., Butler, R. E., Mendum, T. A., Hanna, N., Stewart, G. R. and Soldati, T. (2024). Temporal genome-wide fitness analysis of *Mycobacterium marinum* during infection reveals the genetic requirement for virulence and survival in amoebae and microglial cells. *mSystems* **9**, e0132623. doi:10.1128/msystems.01326-23
- Leon-Torres, A., Arango, E., Castillo, E. and Soto, C. Y. (2020). CtpB is a plasma membrane copper (I) transporting P-type ATPase of *Mycobacterium tuberculosis*. *Biol. Res.* **53**, 6. doi:10.1186/s40659-020-00274-7
- Lewis, K. N., Liao, R., Guinn, K. M., Hickey, M. J., Smith, S., Behr, M. A. and Sherman, D. R. (2003). Deletion of RD1 from *Mycobacterium tuberculosis* mimics bacille Calmette-Guerin attenuation. *J. Infect. Dis.* **187**, 117-123. doi:10.1086/345862
- Li, H. and Li, H. (2023). Animal models of tuberculosis. In: *Vaccines for Neglected Pathogens: Strategies, Achievements and Challenges: Focus on Leprosy, Leishmaniasis, Melioidosis and Tuberculosis* (ed. M. Christodoulides), pp. 139-170. Cham: Springer International Publishing. doi:10.1007/978-3-031-24355-4_7
- Li, Z., Dugan, A. S., Bloomfield, G., Skelton, J., Ivens, A., Losick, V. and Isberg, R. R. (2009). The amoebal MAP kinase response to *Legionella pneumophila* is regulated by DupA. *Cell Host Microbe* **6**, 253-267. doi:10.1016/j.chom.2009.08.005
- Libardo, M. D. J., De, L. A., Fuente-Nunez, C., Anand, K., Krishnamoorthy, G., Kaiser, P., Pringle, S. C., Dietz, C., Pierce, S., Smith, M. B. et al. (2018). Phagosomal copper-promoted oxidative attack on intracellular *Mycobacterium tuberculosis*. *ACS Infect. Dis.* **4**, 1623-1634. doi:10.1021/acscinfdis.8b00171
- Linares, J. F., Duran, A., Yajima, T., Pasparakis, M., Moscat, J. and Diaz-Meco, M. T. (2013). K63 polyubiquitination and activation of mTOR by the p62-TRAF6 complex in nutrient-activated cells. *Mol. Cell* **51**, 283-296. doi:10.1016/j.molcel.2013.06.020
- López-Jiménez, A. (2017). Fate of intracellular mycobacteria and host response to vacuolar escape. *PhD thesis*, University of Geneva. doi:10.13097/archive-ouverte/unige:101548
- Lopez-Jimenez, A. T., Cardenal-Munoz, E., Leuba, F., Gerstenmaier, L., Barisch, C., Hagedorn, M., King, J. S. and Soldati, T. (2018). The ESCRT and autophagy machineries cooperate to repair ESX-1-dependent damage at the *Mycobacterium*-containing vacuole but have opposite impact on containing the infection. *PLoS Pathog.* **14**, e1007501. doi:10.1371/journal.ppat.1007501
- López-Jiménez, A. T., Hagedorn, M., Delincé, M. J., McKinney, J. and Soldati, T. (2019). The developmental cycle of *Dictyostelium discoideum* ensures curing of a mycobacterial infection at both cell-autonomous level and by collaborative exclusion. *bioRxiv* 586263. doi:10.1101/586263
- Lou, J., Wang, Y., Zheng, X. and Qiu, W. (2018). TRIM22 regulates macrophage autophagy and enhances *Mycobacterium tuberculosis* clearance by targeting the nuclear factor-multiplicity kappaB/Beclin 1 pathway. *J. Cell. Biochem.* **119**, 8971-8980. doi:10.1002/jcb.27153
- Ly, J., He, X., Wang, H., Wang, Z., Kelly, G. T., Wang, X., Chen, Y., Wang, T. and Qian, Z. (2017). TLR4-NOX2 axis regulates the phagocytosis and killing of *Mycobacterium tuberculosis* by macrophages. *BMC Pulm. Med.* **17**, 194. doi:10.1186/s12890-017-0517-0
- Madiraju, C., Novack, J. P., Reed, J. C. and Matsuzawa, S. I. (2022). K63 ubiquitination in immune signaling. *Trends Immunol.* **43**, 148-162. doi:10.1016/j.it.2021.12.005
- Maeda, N., Nigou, J., Herrmann, J. L., Jackson, M., Amara, A., Lagrange, P. H., Puzo, G., Gicquel, B. and Neyrolles, O. (2003). The cell surface receptor DC-SIGN discriminates between *Mycobacterium* species through selective recognition of the mannose caps on lipoarabinomannan. *J. Biol. Chem.* **278**, 5513-5516. doi:10.1074/jbc.C200586200
- Manske, C., Finsel, I., Hoffmann, C. and Hilbi, H. (2018). Analysis of *Legionella* metabolism by pathogen vacuole proteomics. *Methods Mol. Biol.* **1841**, 59-76. doi:10.1007/978-1-4939-8695-8_6
- Manzanillo, P. S., Ayres, J. S., Watson, R. O., Collins, A. C., Souza, G., Rae, C. S., Schneider, D. S., Nakamura, K., Shiloh, M. U. and Cox, J. S. (2013). The ubiquitin ligase parkin mediates resistance to intracellular pathogens. *Nature* **501**, 512-516. doi:10.1038/nature12566
- Marchetti, A., Lelong, E. and Cosson, P. (2009). A measure of endosomal pH by flow cytometry in *Dictyostelium*. *BMC Res. Notes* **2**, 7. doi:10.1186/1756-0500-2-7
- Maret, W. (2013). Zinc biochemistry: from a single zinc enzyme to a key element of life. *Adv. Nutr.* **4**, 82-91. doi:10.3945/an.112.003038
- Marinho, F. V., Fahel, J. S., De Araujo, A., Diniz, L. T. S., Gomes, M. T. R., Resende, D. P., Junqueira-Kipnis, A. P. and Oliveira, S. C. (2020). Guanylate binding proteins contained in the murine chromosome 3 are important to control mycobacterial infection. *J. Leukoc. Biol.* **108**, 1279-1291. doi:10.1002/JLB.4MA0620-526RR
- Marshall, E. K. P. and Dionne, M. S. (2022). *Drosophila* versus *Mycobacteria*: a model for mycobacterial host-pathogen interactions. *Mol. Microbiol.* **117**, 600-609. doi:10.1111/mmi.14819
- Mathieu, S. V., Aragao, K. S., Imbert, A. and Varrot, A. (2010). Discoidin I from *Dictyostelium discoideum* and interactions with oligosaccharides: specificity, affinity, crystal structures, and comparison with discoidin II. *J. Mol. Biol.* **400**, 540-554. doi:10.1016/j.jmb.2010.05.042
- Maya-Hoyos, M., Mata-Espinosa, D., Lopez-Torres, M. O., Tovar-Vazquez, B., Barrios-Payan, J., Leon-Contreras, J. C., Ocampo, M., Hernandez-Pando, R. and Soto, C. Y. (2022). The ctpF gene encoding a calcium P-type ATPase of the plasma membrane contributes to full virulence of *Mycobacterium tuberculosis*. *Int. J. Mol. Sci.* **23**, 6015. doi:10.3390/ijms23116015
- McLaughlin, B., Chon, J. S., Macgurn, J. A., Carlsson, F., Cheng, T. L., Cox, J. S. and Brown, E. J. (2007). A mycobacterium ESX-1-secreted virulence factor with unique requirements for export. *PLoS Pathog.* **3**, e105. doi:10.1371/journal.ppat.0030105
- Medapati, R. V., Suvvari, S., Godi, S. and Gangiseti, P. (2017). NRAMP1 and VDR gene polymorphisms in susceptibility to pulmonary tuberculosis among Andhra Pradesh population in India: a case-control study. *BMC Pulm. Med.* **17**, 89. doi:10.1186/s12890-017-0431-5
- Medha, P., Bhatt, P., Sharma, S. and Sharma, M. and Sharma, M. (2023). Role of C-terminal domain of *Mycobacterium tuberculosis* PE6 (Rv0335c) protein in host mitochondrial stress and macrophage apoptosis. *Apoptosis* **28**, 136-165. doi:10.1007/s10495-022-01778-1
- Mehra, A., Zahra, A., Thompson, V., Sirisaengtaksin, N., Wells, A., Porto, M., Koster, S., Penberthy, K., Kubota, Y., Dricot, A. et al. (2013). *Mycobacterium*

- tuberculosis* type VII secreted effector EsxH targets host ESCRT to impair trafficking. *PLoS Pathog.* **9**, e1003734. doi:10.1371/journal.ppat.1003734
- Mesquita, A., Cardenal-Munoz, E., Dominguez, E., Munoz-Braceras, S., Nunez-Corcuer, B., Phillips, B. A., Tabara, L. C., Xiong, Q., Coria, R., Eichinger, L. et al. (2017). Autophagy in *Dictyostelium*: mechanisms, regulation and disease in a simple biomedical model. *Autophagy* **13**, 24–40. doi:10.1080/15548627.2016.1226737
- Miller, J. L., Velmurugan, K., Cowan, M. J. and Briken, V. (2010). The type I NADH dehydrogenase of *Mycobacterium tuberculosis* counters phagosomal NOX2 activity to inhibit TNF- α -mediated host cell apoptosis. *PLoS Pathog.* **6**, e1000864. doi:10.1371/journal.ppat.1000864
- Mishra, M., Adhyapak, P., Dadhich, R. and Kapoor, S. (2019). Dynamic remodeling of the host cell membrane by virulent mycobacterial sulfolipid-1. *Sci. Rep.* **9**, 12844. doi:10.1038/s41598-019-49343-2
- Mittal, E., Skowrya, M. L., Uwase, G., Tinaztepe, E., Mehra, A., Koster, S., Hanson, P. I. and Phillips, J. A. (2018). *Mycobacterium tuberculosis* type VII secretion system effectors differentially impact the ESCRT endomembrane damage response. *mBio* **9**, e01765-18. doi:10.1128/mBio.01765-18
- Mo, S., Guo, J., Ye, T., Zhang, X., Zeng, J., Xu, Y., Peng, B., Dai, Y., Xiao, W., Zhang, P. et al. (2022). *Mycobacterium tuberculosis* utilizes host histamine receptor H1 to modulate reactive oxygen species production and phagosome maturation via the p38MAPK-NOX2 axis. *mBio* **13**, e0200422.
- Morrison, H. M., Craft, J., Rivera-Lugo, R., Johnson, J. R., Golovkine, G. R., Bell, S. L., Dodd, C. E., Van Dis, E., Beatty, W. L., Margolis, S. R. et al. (2023). Deficiency in Galectin-3, -8, and -9 impairs immunity to chronic *Mycobacterium tuberculosis* infection but not acute infection with multiple intracellular pathogens. *PLoS Pathog.* **19**, e1011088. doi:10.1371/journal.ppat.1011088
- Mottet, M., Bosmani, C., Hanna, N., Nitschke, J., Lefrancois, L. H. and Soldati, T. (2021). Novel single-cell and high-throughput microscopy techniques to monitor *Dictyostelium discoideum*-*Mycobacterium marinum* infection dynamics. *Methods Mol. Biol.* **2314**, 183–203. doi:10.1007/978-1-0716-1460-0_7
- Murdoch, C. C. and Skaar, E. P. (2022). Nutritional immunity: the battle for nutrient metals at the host-pathogen interface. *Nat. Rev. Microbiol.* **20**, 657–670. doi:10.1038/s41579-022-00745-6
- Neyrolles, O., Mintz, E. and Catty, P. (2013). Zinc and copper toxicity in host defense against pathogens: *Mycobacterium tuberculosis* as a model example of an emerging paradigm. *Front. Cell Infect. Microbiol.* **3**, 89. doi:10.3389/fcimb.2013.00089
- Neyrolles, O., Wolschendorf, F., Mitra, A. and Niederweis, M. (2015). *Mycobacteria*, metals, and the macrophage. *Immunol. Rev.* **264**, 249–263. doi:10.1111/immr.12265
- Nicolussi, A., Dunn, J. D., Mlynek, G., Bellei, M., Zamocky, M., Battistuzzi, G., Djinoic-Carugo, K., Furtmuller, P. G., Soldati, T. and Obinger, C. (2018). Secreted heme peroxidase from *Dictyostelium discoideum*: insights into catalysis, structure, and biological role. *J. Biol. Chem.* **293**, 1330–1345. doi:10.1074/jbc.RA117.000463
- Olive, A. J., Smith, C. M., Baer, C. E., Coers, J. and Sasseti, C. M. (2023). *Mycobacterium tuberculosis* evasion of guanylate binding protein-mediated host defense in mice requires the ESX1 secretion system. *Int. J. Mol. Sci.* **24**, 2861. doi:10.3390/ijms24032861
- Osman, M. M., Shanahan, J. K., Chu, F., Takaki, K. K., Pinckert, M. L., Pagan, A. J., Brosch, R., Conrad, W. H. and Ramakrishnan, L. (2022). The C terminus of the mycobacterium ESX-1 secretion system substrate ESAT-6 is required for phagosomal membrane damage and virulence. *Proc. Natl. Acad. Sci. USA* **119**, e2122161119. doi:10.1073/pnas.2122161119
- Paci, H. T., Reddy, V. P., Saini, V., Chinta, K. C. and Steyn, A. J. C. (2018). Host-pathogen redox dynamics modulate *Mycobacterium tuberculosis* pathogenesis. *Pathog. Dis.* **76**, fty036. doi:10.1093/femspd/fty036
- Pagán, A. J., Lee, L. J., Edwards-Hicks, J., Moens, C. B., Tobin, D. M., Busch-Nentwich, E. M., Pearce, E. L. and Ramakrishnan, L. (2022). mTOR-regulated mitochondrial metabolism limits mycobacterium-induced cytotoxicity. *Cell* **185**, 3720–3738.e13. doi:10.1016/j.cell.2022.08.018
- Pajuelo, D., Gonzalez-Juarbe, N., Tak, U., Sun, J., Orihuela, C. J. and Niederweis, M. (2018). NAD⁺ depletion triggers macrophage necroptosis, a cell death pathway exploited by *Mycobacterium tuberculosis*. *Cell Rep.* **24**, 429–440. doi:10.1016/j.celrep.2018.06.042
- Patin, E. C., Geffken, A. C., Willcocks, S., Leschczyk, C., Haas, A., Nimmerjahn, F., Lang, R., Ward, T. H. and Schaible, U. E. (2017). Trehalose dimycolate interferes with Fc γ mediated phagosome maturation through Mincle, SHP-1 and Fc γ mediated RILB signalling. *PLoS One* **12**, e0174973. doi:10.1371/journal.pone.0174973
- Pellegrino, E., Aylan, B., Bussi, C., Fearn, A., Bernard, E. M., Athanasiadi, N., Santucci, P., Botella, L. and Gutierrez, M. G. (2023). Peroxisomal ROS control cytosolic *Mycobacterium tuberculosis* replication in human macrophages. *J. Cell Biol.* **222**, e202303066. doi:10.1083/jcb.202303066
- Peracino, B., Wagner, C., Balest, A., Balbo, A., Pergolizzi, B., Noegel, A. A., Steinert, M. and Bozzaro, S. (2006). Function and mechanism of action of *Dictyostelium* Nramp1 (Slc11a1) in bacterial infection. *Traffic* **7**, 22–38. doi:10.1111/j.1600-0854.2005.00356.x
- Peracino, B., Buracco, S. and Bozzaro, S. (2013). The Nramp (Slc11) proteins regulate development, resistance to pathogenic bacteria and iron homeostasis in *Dictyostelium discoideum*. *J. Cell Sci.* **126**, 301–311. doi:10.1242/jcs.116210
- Pergolizzi, B., Bozzaro, S. and Bracco, E. (2019). *Dictyostelium* as model for studying ubiquitination and deubiquitination. *Int. J. Dev. Biol.* **63**, 529–539. doi:10.1387/jdb.190260eb
- Philips, J. A., Porto, M. C., Wang, H., Rubin, E. J. and Perrimon, N. (2008). ESCRT factors restrict mycobacterial growth. *Proc. Natl. Acad. Sci. USA* **105**, 3070–3075. doi:10.1073/pnas.0707206105
- Podinovskaia, M., Lee, W., Caldwell, S. and Russell, D. G. (2013). Infection of macrophages with *Mycobacterium tuberculosis* induces global modifications to phagosomal function. *Cell. Microbiol.* **15**, 843–859. doi:10.1111/cmi.12092
- Poirier, V., Bach, H. and Av-Gay, Y. (2014). *Mycobacterium tuberculosis* promotes anti-apoptotic activity of the macrophage by PtpA protein-dependent dephosphorylation of host GSK3 α . *J. Biol. Chem.* **289**, 29376–29385. doi:10.1074/jbc.M114.582502
- Puri, R. V., Reddy, P. V. and Tyagi, A. K. (2013). Secreted acid phosphatase (SapM) of *Mycobacterium tuberculosis* is indispensable for arresting phagosomal maturation and growth of the pathogen in guinea pig tissues. *PLoS One* **8**, e70514. doi:10.1371/journal.pone.0070514
- Qiang, L., Zhang, Y., Lei, Z., Lu, Z., Tan, S., Ge, P., Chai, Q., Zhao, M., Zhang, X., Li, B. et al. (2023). A mycobacterial effector promotes ferroptosis-dependent pathogenicity and dissemination. *Nat. Commun.* **14**, 1430. doi:10.1038/s41467-023-37148-x
- Quigley, J., Hughitt, V. K., Velikovskiy, C. A., Mariuzza, R. A., El-Sayed, N. M. and Briken, V. (2017). The cell wall lipid PDIM contributes to phagosomal escape and host cell exit of *Mycobacterium tuberculosis*. *mBio* **8**, e00148-17. doi:10.1128/mBio.00148-17
- Radulovic, M., Schink, K. O., Wenzel, E. M., Nähse, V., Bongiovanni, A., Lafont, F. and Stenmark, H. (2018). ESCRT-mediated lysosome repair precedes lysophagy and promotes cell survival. *EMBO J.* **37**, e99753. doi:10.15252/embj.201899753
- Radulovic, M., Wenzel, E. M., Gilani, S., Holland, L. K., Lystad, A. H., Phuyal, S., Oikkonen, V. M., Brech, A., Jaattela, M., Maeda, K. et al. (2022). Cholesterol transfer via endoplasmic reticulum contacts mediates lysosome damage repair. *EMBO J.* **41**, e112677. doi:10.15252/embj.2022112677
- Ramakrishnan, L. (2013). The zebrafish guide to tuberculosis immunity and treatment. *Cold Spring Harb. Symp. Quant. Biol.* **78**, 179–192. doi:10.1101/sqb.2013.78.023283
- Raykov, L. (2021). Identification and characterization of *Dictyostelium discoideum* conserved response factors involved in pathogen detection and stress signal transduction. *PhD thesis*, University of Geneva. doi:10.13097/archive-ouverte/unige:155688
- Raykov, L., Mottet, M., Nitschke, J. and Soldati, T. (2023). A TRAF-like E3 ubiquitin ligase TrafE coordinates ESCRT and autophagy in endolysosomal damage response and cell-autonomous immunity to *Mycobacterium marinum*. *Elife* **12**, e85727. doi:10.7554/eLife.85727
- Roca, F. J., Whitworth, L. J., Redmond, S., Jones, A. A. and Ramakrishnan, L. (2019). TNF induces pathogenic programmed macrophage necrosis in tuberculosis through a mitochondrial-lysosomal-endoplasmic reticulum circuit. *Cell* **178**, 1344–1361.e11. doi:10.1016/j.cell.2019.08.004
- Rodriguez, G. M., Sharma, N., Biswas, A. and Sharma, N. (2022). The iron response of *Mycobacterium tuberculosis* and its implications for tuberculosis pathogenesis and novel therapeutics. *Front. Cell Infect. Microbiol.* **12**, 876667. doi:10.3389/fcimb.2022.876667
- Rojas, M., Garcia, L. F., Nigou, J., Puzo, G. and Olivier, M. (2000). Mannosylated lipoarabinomannan antagonizes *Mycobacterium tuberculosis*-induced macrophage apoptosis by altering Ca²⁺-dependent cell signaling. *J. Infect. Dis.* **182**, 240–251. doi:10.1086/315676
- Romagnoli, A., Di Rienzo, M., Petruccioli, E., Fusco, C., Palucci, I., Micale, L., Mazza, T., Delogu, G., Merla, G., Goletti, D. et al. (2023). The ubiquitin ligase TRIM32 promotes the autophagic response to *Mycobacterium tuberculosis* infection in macrophages. *Cell Death Dis.* **14**, 505. doi:10.1038/s41419-023-06026-1
- Ruan, H., Lyu, M., Lai, H., Niu, L., Zhao, Z., Liu, T., Lei, S. and Ying, B. (2024). Host-pathogen dialogues in different cell death modes during *Mycobacterium tuberculosis* infection. *Interdiscip. Med.* **2**, e20230044. doi:10.1002/INMD.20230044
- Saelens, J. W., Sweeney, M. I., Viswanathan, G., Xet-Mull, A. M., Jurcic Smith, K. L., Sisk, D. M., Hu, D. D., Cronin, R. M., Hughes, E. J., Brewer, W. J. et al. (2022). An ancestral mycobacterial effector promotes dissemination of infection. *Cell* **185**, 4507–4525.e18. doi:10.1016/j.cell.2022.10.019
- Saikalappan, S., Estrella, J., Sasindran, S. J., Khan, A., Armitage, L. Y., Jagannath, C. and Dhandayuthapani, S. (2012). The fbpA/sapM double knock out strain of *Mycobacterium tuberculosis* is highly attenuated and immunogenic in macrophages. *PLoS One* **7**, e36198. doi:10.1371/journal.pone.0036198
- Saini, N. K., Baena, A., Ng, T. W., Venkataswamy, M. M., Kennedy, S. C., Kunnath-Velayudhan, S., Carreno, L. J., Xu, J., Chan, J., Larsen, M. H. et al. (2016). Suppression of autophagy and antigen presentation by *Mycobacterium*

- tuberculosis PE_PGRS47. *Nat. Microbiol.* **1**, 16133. doi:10.1038/nmicrobiol.2016.133
- Sattler, N., Bosmani, C., Barisch, C., Gueho, A., Gopaldass, N., Dias, M., Leuba, F., Bruckert, F., Cosson, P. and Soldati, T. (2018). Functions of the *Dictyostelium* LIMP-2 and CD36 homologues in bacteria uptake, phagolysosome biogenesis and host cell defence. *J. Cell Sci.* **131**, jcs218040. doi:10.1242/jcs.218040
- Schmolders, J., Manske, C., Otto, A., Hoffmann, C., Steiner, B., Welin, A., Becher, D. and Hilbi, H. (2017). Comparative proteomics of purified pathogen vacuoles correlates intracellular replication of *Legionella pneumophila* with the small GTPase ras-related protein 1 (Rap1). *Mol. Cell. Proteomics* **16**, 622-641. doi:10.1074/mcp.M116.063453
- Schnettger, L., Rodgers, A., Repnik, U., Lai, R. P., Pei, G., Verdoes, M., Wilkinson, R. J., Young, D. B. and Gutierrez, M. G. (2017). A Rab20-dependent membrane trafficking pathway controls *M. tuberculosis* replication by regulating phagosome spaciousness and integrity. *Cell Host Microbe* **21**, 619-628.e5. doi:10.1016/j.chom.2017.04.004
- Serene, L. G., Webber, K., Champion, P. A. and Schorey, J. S. (2024). *Mycobacterium tuberculosis* SecA2-dependent activation of host Rig-I/MAVs signaling is not conserved in *Mycobacterium marinum*. *PLoS One* **19**, e0281564. doi:10.1371/journal.pone.0281564
- Sethi, D., Mahajan, S., Singh, C., Lama, A., Hade, M. D., Gupta, P. and Dikshit, K. L. (2016). Lipoprotein Lprl of *Mycobacterium tuberculosis* acts as a lysozyme inhibitor. *J. Biol. Chem.* **291**, 2938-2953. doi:10.1074/jbc.M115.662593
- Seto, S., Tsujimura, K. and Koide, Y. (2011). Rab GTPases regulating phagosome maturation are differentially recruited to mycobacterial phagosomes. *Traffic* **12**, 407-420. doi:10.1111/j.1600-0854.2011.01165.x
- Shah, S., Dalecki, A. G., Malalasekera, A. P., Crawford, C. L., Michalek, S. M., Kutsch, O., Sun, J., Bossmann, S. H. and Wolschendorf, F. (2016). 8-hydroxyquinolines are boosting agents of copper-related toxicity in *Mycobacterium tuberculosis*. *Antimicrob. Agents Chemother.* **60**, 5765-5776. doi:10.1128/AAC.00325-16
- Sharma, S. K. and Upadhyay, V. (2020). Epidemiology, diagnosis & treatment of non-tuberculous mycobacterial diseases. *Indian J. Med. Res.* **152**, 185-226. doi:10.4103/ijmr.IJMR_902_20
- Sharma, N., Shariq, M., Quadir, N., Singh, J., Sheikh, J. A., Hasnain, S. E. and Ehtesham, N. Z. (2021). *Mycobacterium tuberculosis* protein PE6 (Rv0335c), a novel TLR4 agonist, evokes an inflammatory response and modulates the cell death pathways in macrophages to enhance intracellular survival. *Front. Immunol.* **12**, 696491. doi:10.3389/fimmu.2021.696491
- Shin, D. M., Jeon, B. Y., Lee, H. M., Jin, H. S., Yuk, J. M., Song, C. H., Lee, S. H., Lee, Z. W., Cho, S. N., Kim, J. M. et al. (2010). *Mycobacterium tuberculosis* Eis regulates autophagy, inflammation, and cell death through redox-dependent signaling. *PLoS Pathog.* **6**, e1001230. doi:10.1371/journal.ppat.1001230
- Simeone, R., Bobard, A., Lippmann, J., Bitter, W., Majlessi, L., Brosch, R. and Enninga, J. (2012). Phagosomal rupture by *Mycobacterium tuberculosis* results in toxicity and host cell death. *PLoS Pathog.* **8**, e1002507. doi:10.1371/journal.ppat.1002507
- Singh, V. K., Berry, L., Bernut, A., Singh, S., Carrere-Kremer, S., Viljoen, A., Ailabaud, L., Majlessi, L., Brosch, R., Chaturvedi, V. et al. (2016). A unique PE_PGRS protein inhibiting host cell cytosolic defenses and sustaining full virulence of *Mycobacterium marinum* in multiple hosts. *Cell. Microbiol.* **18**, 1489-1507. doi:10.1111/cmi.12606
- Smirnov, A., Daily, K. P., Gray, M. C., Ragland, S. A., Werner, L. M., Brittany Johnson, M., Eby, J. C., Hewlett, E. L., Taylor, R. P. and Criss, A. K. (2023). Phagocytosis via complement receptor 3 enables microbes to evade killing by neutrophils. *J. Leukoc. Biol.* **114**, 1-20. doi:10.1093/leuko/qiad028
- Smith, J., Manoranjan, J., Pan, M., Bohsali, A., Xu, J., Liu, J., McDonald, K. L., Szyk, A., Laronde-Leblanc, N. and Gao, L. Y. (2008). Evidence for pore formation in host cell membranes by ESX-1-secreted ESAT-6 and its role in *Mycobacterium marinum* escape from the vacuole. *Infect. Immun.* **76**, 5478-5487. doi:10.1128/IAI.00614-08
- Soldati, T. and Neyrolles, O. (2012). Mycobacteria and the intraphagosomal environment: take it with a pinch of salt(s)!. *Traffic* **13**, 1042-1052. doi:10.1111/j.1600-0854.2012.01358.x
- Song, O. R., Queval, C. J., Iantomasi, R., Delorme, V., Marion, S., Veyron-Churlet, R., Werkmeister, E., Popoff, M., Ricard, I., Jouny, S. et al. (2018). ArfGAP1 restricts *Mycobacterium tuberculosis* entry by controlling the actin cytoskeleton. *EMBO Rep.* **19**, 29-42. doi:10.15252/embr.201744371
- Spargo, B. J., Crowe, L. M., Ionedo, T., Beaman, B. L. and Crowe, J. H. (1991). Cord factor (alpha,alpha-trehalose 6,6'-dimycolate) inhibits fusion between phospholipid vesicles. *Proc. Natl. Acad. Sci. USA* **88**, 737-740. doi:10.1073/pnas.88.3.737
- Speer, A., Shrestha, T. B., Bossmann, S. H., Basaraba, R. J., Harber, G. J., Michalek, S. M., Niederweis, M., Kutsch, O. and Wolschendorf, F. (2013). Copper-boosting compounds: a novel concept for antimycobacterial drug discovery. *Antimicrob. Agents Chemother.* **57**, 1089-1091. doi:10.1128/AAC.01781-12
- Stamm, C. E., Collins, A. C. and Shiloh, M. U. (2015). Sensing of *Mycobacterium tuberculosis* and consequences to both host and bacillus. *Immunol. Rev.* **264**, 204-219. doi:10.1111/imr.12263
- Stinear, T. P., Seemann, T., Harrison, P. F., Jenkin, G. A., Davies, J. K., Johnson, P. D., Abdallah, Z., Arrowsmith, C., Chillingworth, T., Churcher, C. et al. (2008). Insights from the complete genome sequence of *Mycobacterium marinum* on the evolution of *Mycobacterium tuberculosis*. *Genome Res.* **18**, 729-741. doi:10.1101/gr.075069.107
- Strong, E. J., Ng, T. W., Porcelli, S. A. and Lee, S. (2021). *Mycobacterium tuberculosis* PE_PGRS20 and PE_PGRS47 proteins inhibit autophagy by interaction with Rab1A. *mSphere* **6**, e0054921. doi:10.1128/mSphere.00549-21
- Sun, J., Wang, X., Lau, A., Liao, T. Y., Bucci, C. and Hmama, Z. (2010). Mycobacterial nucleoside diphosphate kinase blocks phagosome maturation in murine RAW 264.7 macrophages. *PLoS One* **5**, e8769. doi:10.1371/journal.pone.0008769
- Sun, J., Singh, V., Lau, A., Stokes, R. W., Obregon-Henao, A., Orme, I. M., Wong, D., Av-Gay, Y. and Hmama, Z. (2013). *Mycobacterium tuberculosis* nucleoside diphosphate kinase inactivates small GTPases leading to evasion of innate immunity. *PLoS Pathog.* **9**, e1003499. doi:10.1371/journal.ppat.1003499
- Swart, A. L., Harrison, C. F., Eichinger, L., Steinert, M. and Hilbi, H. (2018). *Acanthamoeba* and *Dictyostelium* as cellular models for *Legionella* infection. *Front. Cell Infect. Microbiol.* **8**, 61. doi:10.3389/fcimb.2018.00061
- Talukder, M. S. U., Pervin, M. S., Tanvir, M. I. O., Fujimoto, K., Tanaka, M., Itoh, G. and Yumura, S. (2020). Ca²⁺-calmodulin dependent wound repair in *Dictyostelium* cell membrane. *Cells* **9**, 1058. doi:10.3390/cells9041058
- Tan, J. X. and Finkel, T. (2022). A phosphoinositide signalling pathway mediates rapid lysosomal repair. *Nature* **609**, 815-821. doi:10.1038/s41586-022-05164-4
- Tan, T., Lee, W. L., Alexander, D. C., Grinstein, S. and Liu, J. (2006). The ESAT-6/CFP-10 secretion system of *Mycobacterium marinum* modulates phagosome maturation. *Cell. Microbiol.* **8**, 1417-1429. doi:10.1111/j.1462-5822.2006.00721.x
- Tobin, D. M. and Ramakrishnan, L. (2008). Comparative pathogenesis of *Mycobacterium marinum* and *Mycobacterium tuberculosis*. *Cell. Microbiol.* **10**, 1027-1039. doi:10.1111/j.1462-5822.2008.01133.x
- Toniolo, C., Dhar, N. and McKinney, J. D. (2023). Uptake-independent killing of macrophages by extracellular *Mycobacterium tuberculosis* aggregates. *EMBO J.* **42**, e113490. doi:10.15252/embj.2023113490
- Tornero-Ecija, A., Tabara, L. C., Bueno-Arribas, M., Anton-Esteban, L., Navarro-Gomez, C., Sanchez, I., Vincent, O. and Escalante, R. (2022). A *Dictyostelium* model for BPAN disease reveals a functional relationship between the WDR45/WIP14 homolog Wdr45l and Vmp1 in the regulation of autophagy-associated PtdIns3P and ER stress. *Autophagy* **18**, 661-677. doi:10.1080/15548627.2021.1953262
- Toyonaga, K., Torigoe, S., Motomura, Y., Kamichi, T., Hayashi, J. M., Morita, Y. S., Noguchi, N., Chuma, Y., Kiyohara, H., Matsuo, K. et al. (2016). C-type lectin receptor DCAR recognizes mycobacterial phosphatidyl-inositol mannosides to promote a Th1 response during infection. *Immunity* **45**, 1245-1257. doi:10.1016/j.immuni.2016.10.012
- Trivedi, P. C., Bartlett, J. J. and Pulini, T. (2020). Lysosomal biology and function: modern view of cellular debris bin. *Cells* **9**, 1131. doi:10.3390/cells9051131
- Tu, H., Wang, Z., Yuan, Y., Miao, X., Li, D., Guo, H., Yang, Y. and Cai, H. (2022). The PripA-TbcA complex-centered Rab GAP cascade facilitates macropinosome maturation in *Dictyostelium*. *Nat. Commun.* **13**, 1787. doi:10.1038/s41467-022-29503-1
- Tyagi, P., Dharmaraja, A. T., Bhaskar, A., Chakrapani, H. and Singh, A. (2015). *Mycobacterium tuberculosis* has diminished capacity to counteract redox stress induced by elevated levels of endogenous superoxide. *Free Radic. Biol. Med.* **84**, 344-354. doi:10.1016/j.freeradbiomed.2015.03.008
- Van Der Woude, A. D., Stoop, E. J., Stieff, M., Wang, S., Ummels, R., Van Stempvoort, G., Piersma, S. R., Cascioferro, A., Jimenez, C. R., Houben, E. N. et al. (2014). Analysis of SecA2-dependent substrates in *Mycobacterium marinum* identifies protein kinase G (PknG) as a virulence effector. *Cell. Microbiol.* **16**, 280-295. doi:10.1111/cmi.12221
- Varshney, D., Singh, S., Sinha, E., Mohanty, K. K., Kumar, S., Kumar Barik, S., Patil, S. A. and Katara, P. (2022). Systematic review and meta-analysis of human Toll-like receptors genetic polymorphisms for susceptibility to tuberculosis infection. *Cytokine* **152**, 155791. doi:10.1016/j.cyto.2021.155791
- Vergne, I., Gilleron, M. and Nigou, J. (2014). Manipulation of the endocytic pathway and phagocyte functions by *Mycobacterium tuberculosis* lipoarabinomannan. *Front. Cell Infect. Microbiol.* **4**, 187. doi:10.3389/fcimb.2014.00187
- Vines, J. H., Maib, H., Buckley, C. M., Gueho, A., Soldati, T., Murray, D. and King, J. S. (2023). A PI(3,5)P2 reporter reveals PIKfyve activity and dynamics on macropinosomes and phagosomes. *J. Cell Biol.* **222**, e202209077. doi:10.1083/jcb.202209077
- Wang, J., Li, B. X., Ge, P. P., Li, J., Wang, Q., Gao, G. F., Qiu, X. B. and Liu, C. H. (2015). *Mycobacterium tuberculosis* suppresses innate immunity by coopting the host ubiquitin system. *Nat. Immunol.* **16**, 237-245. doi:10.1038/ni.3096
- Wang, J., Ge, P., Lei, Z., Lu, Z., Qiang, L., Chai, Q., Zhang, Y., Zhao, D., Li, B., Su, J. et al. (2021). *Mycobacterium tuberculosis* protein kinase G acts as an unusual

- ubiquitinating enzyme to impair host immunity. *EMBO Rep.* **22**, e52175. doi:10.15252/embr.202052175
- Wang, Y. T., Liu, T. Y., Shen, C. H., Lin, S. Y., Hung, C. C., Hsu, L. C. and Chen, G. C. (2022). K48/K63-linked polyubiquitination of ATG9A by TRAF6 E3 ligase regulates oxidative stress-induced autophagy. *Cell Rep.* **38**, 110354-110354. doi:10.1016/j.celrep.2022.110354
- Watson, R. O., Manzanillo, P. S. and Cox, J. S. (2012). Extracellular *M. tuberculosis* DNA targets bacteria for autophagy by activating the host DNA-sensing pathway. *Cell* **150**, 803-815. doi:10.1016/j.cell.2012.06.040
- Watson, R. O., Bell, S. L., Macduff, D. A., Kimmey, J. M., Diner, E. J., Olivas, J., Vance, R. E., Stallings, C. L., Virgin, H. W. and Cox, J. S. (2015). The cytosolic sensor cGAS detects *Mycobacterium tuberculosis* DNA to induce type I interferons and activate autophagy. *Cell Host Microbe* **17**, 811-819. doi:10.1016/j.chom.2015.05.004
- Welin, A., Winberg, M. E., Abdalla, H., Sarndahl, E., Rasmusson, B., Stendahl, O. and Lerm, M. (2008). Incorporation of *Mycobacterium tuberculosis* lipoarabinomannan into macrophage membrane rafts is a prerequisite for the phagosomal maturation block. *Infect. Immun.* **76**, 2882-2887. doi:10.1128/IAI.01549-07
- Welin, A., Eklund, D., Stendahl, O. and Lerm, M. (2011). Human macrophages infected with a high burden of ESAT-6-expressing *M. tuberculosis* undergo caspase-1- and cathepsin B-independent necrosis. *PLoS One* **6**, e20302. doi:10.1371/journal.pone.0020302
- Westman, J., Plumb, J., Licht, A., Yang, M., Allert, S., Naglik, J. R., Hube, B., Grinstein, S. and Maxson, M. E. (2022). Calcium-dependent ESCRT recruitment and lysosome exocytosis maintain epithelial integrity during *Candida albicans* invasion. *Cell Rep.* **38**, 110187. doi:10.1016/j.celrep.2021.110187
- WHO. (2023). Global Tuberculosis Report. World Health Organization. <https://www.who.int/publications/i/item/9789240083851>
- Wong, D., Bach, H., Sun, J., Hmama, Z. and Av-Gay, Y. (2011). *Mycobacterium tuberculosis* protein tyrosine phosphatase (PtpA) excludes host vacuolar-H⁺-ATPase to inhibit phagosomal acidification. *Proc. Natl. Acad. Sci. USA* **108**, 19371-19376. doi:10.1073/pnas.1109201108
- Wong, D., Li, W., Chao, J. D., Zhou, P., Narula, G., Tsui, C., Ko, M., Xie, J., Martinez-Frailes, C. and Av-Gay, Y. (2018). Protein tyrosine kinase, PtkA, is required for *Mycobacterium tuberculosis* growth in macrophages. *Sci. Rep.* **8**, 155. doi:10.1038/s41598-017-18547-9
- Wu, X., Wu, Y., Zheng, R., Tang, F., Qin, L., Lai, D., Zhang, L., Chen, L., Yan, B., Yang, H. et al. (2021). Sensing of mycobacterial arabinogalactan by galectin-9 exacerbates mycobacterial infection. *EMBO Rep.* **22**, e51678. doi:10.15252/embr.202051678
- Xander, C., Rajagopalan, S., Jacobs, W. R. Jr and Braunstein, M. (2024). The SapM phosphatase can arrest phagosome maturation in an ESX-1 independent manner in *Mycobacterium tuberculosis* and BCG. *Infect. Immun.* [Epub] e0021724. doi:10.1128/iai.00217-24
- Xiong, Q., Feng, R., Fischer, S., Karow, M., Stumpf, M., Meßling, S., Nitz, L., Müller, S., Clemen, C. S., Song, N. et al. (2023). Proteasomes of autophagy-deficient cells exhibit alterations in regulatory proteins and a marked reduction in activity. *Cells* **12**, 1514. doi:10.3390/cells12111514
- Xu, J., Laine, O., Masciocchi, M., Manoranjan, J., Smith, J., Du, S. J., Edwards, N., Zhu, X., Fenselau, C. and Gao, L. Y. (2007). A unique *Mycobacterium* ESX-1 protein co-secretes with CFP-10/ESAT-6 and is necessary for inhibiting phagosome maturation. *Mol. Microbiol.* **66**, 787-800. doi:10.1111/j.1365-2958.2007.05959.x
- Xu, X., Pan, M. and Jin, T. (2021). How phagocytes acquired the capability of hunting and removing pathogens from a human body: lessons learned from chemotaxis and phagocytosis of *Dictyostelium discoideum*. *Front. Cell Dev. Biol.* **9**, 724940. doi:10.3389/fcell.2021.724940
- Yang, C. S., Yuk, J. M. and Jo, E. K. (2009). The role of nitric oxide in mycobacterial infections. *Immune Netw.* **9**, 46-52. doi:10.4110/in.2009.9.2.46
- Yu, X., Huang, Y., Li, Y., Li, T., Yan, S., Ai, X., Lv, X., Fan, L. and Xie, J. (2023). *Mycobacterium tuberculosis* PE_PGRS1 promotes mycobacteria intracellular survival via reducing the concentration of intracellular free Ca²⁺ and suppressing endoplasmic reticulum stress. *Mol. Immunol.* **154**, 24-32. doi:10.1016/j.molimm.2022.12.007
- Yuan, A., Siu, C. H. and Chia, C. P. (2001). Calcium requirement for efficient phagocytosis by *Dictyostelium discoideum*. *Cell Calcium* **29**, 229-238. doi:10.1054/ceca.2000.0184
- Zhang, L., Zhang, H., Zhao, Y., Mao, F., Wu, J., Bai, B., Xu, Z., Jiang, Y. and Shi, C. (2012). Effects of *Mycobacterium tuberculosis* ESAT-6/CFP-10 fusion protein on the autophagy function of mouse macrophages. *DNA Cell Biol.* **31**, 171-179. doi:10.1089/dna.2011.1290
- Zhang, Q., Wang, D., Jiang, G., Liu, W., Deng, Q., Li, X., Qian, W., Ouellet, H. and Sun, J. (2016a). EsxA membrane-permeabilizing activity plays a key role in mycobacterial cytosolic translocation and virulence: effects of single-residue mutations at glutamine 5. *Sci. Rep.* **6**, 32618. doi:10.1038/srep32618
- Zhang, X., Zhuchenko, O., Kuspa, A. and Soldati, T. (2016b). Social amoebae trap and kill bacteria by casting DNA nets. *Nat. Commun.* **7**, 10938. doi:10.1038/ncomms10938
- Zhang, L., Jiang, X., Pfau, D., Ling, Y. and Nathan, C. F. (2021). Type I interferon signaling mediates *Mycobacterium tuberculosis*-induced macrophage death. *J. Exp. Med.* **218**, e20200887. doi:10.1084/jem.20200887
- Zhao, X., Khan, N., Gan, H., Tzelepis, F., Nishimura, T., Park, S. Y., Divangahi, M. and Remold, H. G. (2017). Bcl-x(L) mediates RIPK3-dependent necrosis in *M. tuberculosis*-infected macrophages. *Mucosal Immunol.* **10**, 1553-1568. doi:10.1038/mi.2017.12
- Zheng, W., Chang, I. C., Limberis, J., Budzik, J. M., Zha, B. S., Howard, Z., Chen, L. and Ernst, J. D. (2024). *Mycobacterium tuberculosis* resides in lysosome-poor monocyte-derived lung cells during chronic infection. *PLoS Pathog.* **20**, e1012205. doi:10.1371/journal.ppat.1012205
- Zihad, S., Sifat, N., Islam, M. A., Monjur-Al-Hossain, A. S. M., Sikdar, K., Sarker, M. M. R., Shilpi, J. A. and Uddin, S. J. (2023). Role of pattern recognition receptors in sensing *Mycobacterium tuberculosis*. *Heliyon* **9**, e20636. doi:10.1016/j.heliyon.2023.e20636
- Zulauf, K. E., Sullivan, J. T. and Braunstein, M. (2018). The SecA2 pathway of *Mycobacterium tuberculosis* exports effectors that work in concert to arrest phagosome and autophagosome maturation. *PLoS Pathog.* **14**, e1007011. doi:10.1371/journal.ppat.1007011