

## **Archive ouverte UNIGE**

https://archive-ouverte.unige.ch

Article scientifique

Commentaire 2011

**Published** version

Open Access

This is the published version of the publication, made available in accordance with the publisher's policy.

Invited commentary

Licker, Marc

## How to cite

LICKER, Marc. Invited commentary. In: The annals of thoracic surgery, 2011, vol. 92, n° 4, p. 1194. doi: 10.1016/j.athoracsur.2011.06.085

This publication URL: <a href="https://archive-ouverte.unige.ch/unige:32610">https://archive-ouverte.unige.ch/unige:32610</a>

Publication DOI: 10.1016/j.athoracsur.2011.06.085

© This document is protected by copyright. Please refer to copyright holder(s) for terms of use.

- 18. Licker M, de Perrot M, Höhn L, et al. Perioperative mortality and major cardio-pulmonary complications after lung surgery for non-small cell carcinoma. Eur J Cardiothorac Surg 1999;15:314–9.
- 19. Dancewicz M, Kowalewski J, Peplinski J. Factors associated with perioperative complications after pneumonectomy for primary carcinoma of the lung. Interact Cardiovasc Thorac Surg 2006;5:97–100.
- 20. Marret E, Miled F, Bazelly B, et al. Risk and protective factors for major complications after pneumonectomy for lung cancer. Interact Cardiovasc Thorac Surg 2010;10:936–9.
- 21. Thomas P, Michelet P, Barlesi F, et al. Impact of blood transfusions on outcome after pneumonectomy for thoracic malignancies. Eur Respir J 2007;29:565–70.
- 22. van der Werff YD, van der Houwen HK, Heijmans PJ, et al. Postpneumonectomy pulmonary edema. A retrospective analysis of incidence and possible risk factors. Chest 1997; 111:1278–84.
- 23. Linden PA, Yeap BY, Chang MY, et al. Morbidity of lung resection after prior lobectomy: results from the Veterans Affairs National Surgical Quality Improvement Program. Ann Thorac Surg 2007;83:425–32.
- 24. Licker M, Diaper J, Villiger Y, et al. Impact of intraoperative lung-protective interventions in patients undergoing lung cancer surgery. Crit Care 2009;13:R41.
- Michelet P, D'Journo X, Roch A, et al. Protective ventilation influences systemic inflammation after esophagectomy: a randomized controlled study. Anesthesiology 2006;105:911–9.

## **INVITED COMMENTARY**

Several observational studies have already stressed the importance of procedural risk factors such as intraoperative ventilation with large tidal volume, excessive fluid hydration, blood transfusion, absence of epidural thoracic analgesia, and the extent of lung resection [1].

In the current study, Blank and colleagues [2] found a significant association between the occurrence of major respiratory complications and intraoperative administration of homologous blood transfusion (packed red blood cell, fresh frozen plasma) and large amounts of crystalloids. Not surprisingly, these adverse events were associated with prolonged hospital length of stay.

Limitations to this study are not only the small number of patients with respiratory failure (13% in a cohort of 129 pneumonectomies) but also the wide time window that was examined (12 years) and the lack of standardized care bundles. The multivariate logistic model (adjusted for confounding factors) is a valuable hypothesisgenerating tool and, therefore, randomized clinical trials should be designed to question the impact of goal-directed therapy with blood components, crystalloids and colloids aimed to prevent anemia, coagulation disturbances, and hypo/hypervolemia.

The rationale for implementing a "restrictive" transfusion strategy (targeting a hematocrit of 24% instead of 30%), is supported by many studies that have shown a lack of benefit as well as increased adverse effects and costs associated with packed red blood cell transfusion [3]. Likewise, the administration of fresh frozen plasma is known to be ineffective in reducing the amount of blood loss and is even associated with an increased risk of infection [3, 4].

Currently, a variety of noninvasive hemodynamic monitor (eg, transesophageal Doppler, pulse contour analysis of the arterial pressure wave, tissue oximetry) and "point of care" devices for monitoring coagulation (eg, thromboelastometry/graphy) are available for clinical application in the operating room, the postanesthesia care unit and the intensive care unit. Importantly, the effectiveness of goal-directed approach using blood flow monitors has been demonstrated in decreasing morbidity and mortality in high-risk surgical patients [5]. Unfortunately, none of these studies has involved any patients

undergoing lung resection although the use of semiinvasive hemodynamic monitoring devices has been validated during one-lung ventilation [6]. Likewise, bedside application of thromboelastography is better than conventional coagulation assays to monitor perioperative coagulopathy and to predict transfusion requirements [7].

The authors deserve merit and recognition for drawing our attention to preventable adverse events. Future efforts are needed to test "protective" interventions and to devise an integrated bundle of care to manage the high-risk thoracic patients efficiently and safely.

Marc Licker, MD

Anesthesiology, Pharmacology, and Intensive Care University Hospital Geneva rue Micheli-du-crest Geneva, Switzerland 1211 e-mail: marc-joseph.licker@hcuge.ch

## References

- 1. Licker M, Schnyder JM, Frey JG, et al. Impact of aerobic exercise capacity and procedure-related factors in lung cancer surgery. Eur Respir J 2011;37:1189–98.
- Blank RS, Hucklenbruch C, Gurka KK, et al. Intraoperative factors and the risk of respiratory complications after pneumonectomy. Ann Thorac Surg 2011;92:1188–94.
- 3. Society of Thoracic Surgeons Blood Conservation Guideline Task Force, Ferraris VA, Brown JR, et al. 2011 update to the Society of Thoracic Surgeons and the Society of Cardiovascular Anesthesiologists blood conservation clinical practice guidelines. Ann Thorac Surg 2011;91:944–82.
- Sarani B, Dunkman WJ, Dean L, Sonnad S, Rohrbach JI, Gracias VH. Transfusion of fresh frozen plasma in critically ill surgical patients is associated with an increased risk of infection. Crit Care Med 2008;36:1114–8.
- Lees N, Hamilton M, Rhodes A. Clinical review: goal-directed therapy in high risk surgical patients. Crit Care 2009;13:231.
- 6. Diaper J, Ellenberger C, Villiger Y, et al. Comparison of cardiac output as assessed by transesophageal echo-Doppler and transpulmonary thermodilution in patients undergoing thoracic surgery. J Clin Anesth 2010;22:97–103.
- 7. Dickinson KJ, Troxler M, Homer-Vanniasinkam S. The surgical application of point-of-care haemostasis and platelet function testing. Br J Surg 2008;95:1317–30.