



Article scientifique

Lettre

2021

Published version

Open Access

This is the published version of the publication, made available in accordance with the publisher's policy.

Reply to 'Oxic methanogenesis is only a minor source of lake-wide diffusive CH₄ emissions from lakes'

Günthel, Marco; Donis, Daphné; Kirillin, Georgiy; Ionescu, Danny; Bizic, Mina; McGinnis, Daniel Frank; Grossart, Hans-Peter; Tang, Kam W.

How to cite

GÜNTHEL, Marco et al. Reply to 'Oxic methanogenesis is only a minor source of lake-wide diffusive CH₄ emissions from lakes'. In: Nature communications, 2021, vol. 12, n° 1. doi: 10.1038/s41467-021-21216-1

This publication URL: <https://archive-ouverte.unige.ch/unige:166972>

Publication DOI: [10.1038/s41467-021-21216-1](https://doi.org/10.1038/s41467-021-21216-1)

Reply to 'Oxic methanogenesis is only a minor source of lake-wide diffusive CH₄ emissions from lakes'

Marco Günthel ^{1,2,3✉}, Daphne Donis ⁴, Georgiy Kirillin⁵, Danny Ionescu ², Mina Bizic ², Daniel F. McGinnis ^{4✉}, Hans-Peter Grossart ^{2,6✉} & Kam W. Tang^{3✉}

REPLYING TO F. Peeters & H. Hofmann *Nature Communications* <https://doi.org/10.1038/s41467-021-21215-2> (2021)

The prevailing paradigm in methane research is that biological methane production is exclusive to anoxic or near-anoxic habitats such as sediments and oxygen-deficient bottom waters in lakes. Paradoxically, methane supersaturation in oxic lake waters is widely reported. To resolve this paradox while preserving the paradigm, some researchers assume this methane originates entirely from anoxic sources and is then transported to the oxic waters through physical processes^{1–3}. However, multiple recent studies have repeatedly shown, methane production can and does occur under oxic conditions on land, in the seas and in freshwaters, driven by diverse organisms within different life domains (Table 1 and references therein) and via photochemical conversion⁴. These findings raise legitimate questions about the nature of the environmental dynamics and global budget of methane. Because oxic methane production (OMP) is a recent discovery, its contribution to atmospheric emission is unknown. We conducted a whole-lake basin methane mass balance and analysed relevant literature data to estimate the contribution of OMP to surface emission versus lake morphometry.

Because the dynamics of methane concentration and isotope signal in lake waters are influenced by different and opposing processes simultaneously, one cannot meaningfully deduce the presence or absence of OMP without properly accounting for modulations by physical and biological processes. For example, underestimating surface emission or ignoring oxidation would lead to incorrect interpretation of methane concentration and isotope data and incorrect dismissal of OMP (Supplementary Note 1).

By balancing the gains and losses of epilimnetic methane in a stratified water column, we estimated the contribution of oxic versus anoxic methane to surface emission (Supplementary Fig. 1). Epilimnetic methane may originate from lateral and

vertical transport from anoxic zones, ebullition, and internal oxic production (OMP); surface emission and oxidation are the loss terms.

Surface methane emission can be measured directly using a flux chamber, or, in the absence of direct measurements, it is often modelled from surface-water methane concentrations and wind speeds. Both methods are commonly used but the results can differ considerably, and there exist many different wind-based models (for a more detailed discussion we refer readers to the literature^{5,6}). Notably in their manuscript, Peeters and Hofmann excluded our direct measurements of methane fluxes to the atmosphere and exclusively rely on modelling approaches (Supplementary Note 2). We instead combined direct measurements with models that were established for the target lake. Therefore, we consider that our direct measurement approach minimises methodological and model biases, and better represents reality.

For Lake Hallwil, we used the littoral sediment-to-water methane flux as determined by Donis et al.⁷ who implemented two littoral sediment core measurements sampled at 3 and 7 m depth and applying Fick's law. In contrast, Peeters and Hofmann implemented only the upper sediment core into their re-analysis. They justify this choice by stating the cores' methane isotope signature vary. As the depth of Lake Hallwil's surface mixed layer increased over the seasonal progression⁷, both sediment cores should be considered in the mass balance especially in the light of natural variability. For Lake Stechlin, we used data from two mesocosms and the open-water to resolve littoral methane input (Supplementary Notes 3 and 4). We estimated ebullitive methane fluxes as negligible in Lake Stechlin^{8,9}. We further applied an ebullitive flux of $1.2 \pm 0.8 \text{ mmol m}^{-2} \text{ d}^{-1}$ to Lake Hallwil¹⁰, giving a total sediment

¹Department of Environmental Microbiology and Biotechnology, University of Warsaw, Warsaw, Poland. ²Department of Experimental Limnology, Leibniz Institute of Freshwater Ecology and Inland Fisheries, Stechlin, Germany. ³Department of Biosciences, Swansea University, Swansea, UK. ⁴Aquatic Physics Group, Department F.-A. Forel for Environmental and Aquatic Sciences, Faculty of Science, University of Geneva, Geneva, Switzerland. ⁵Department of Ecohydrology, Leibniz Institute of Freshwater Ecology and Inland Fisheries, Berlin, Germany. ⁶Institute of Biochemistry and Biology, Potsdam University, Potsdam, Germany. ✉email: marcoguenthel@gmail.com; daniel.mcginnis@unige.ch; hgrossart@igb-berlin.de; k.w.tang@swansea.ac.uk

Table 1 Literature examples of oxic methane production (OMP) in different habitats and by different domains of life.

Organism	Domain	CH ₄ production rate	Evidence	Reference
TERRESTRIAL				
Plants	Eukaryote		INC, ISO	Keppler et al. (2006) ²⁰
Plants	Eukaryote		INC	Messenger et al. (2009) ²¹
Methanogens	Archaea		INC, ISO, OMIC	Angel et al. (2011) ²²
Fungi	Eukaryote		INC, ISO	Lenhart et al. (2012) ²³
Plants	Eukaryote		INC, ISO	Althoff et al. (2014) ²⁴
Methanogens	Archaea		MB, OMIC, PHYS	Angle et al. (2017) ¹⁴
Cyanobacteria	Prokaryote		INC, ISO	Bizic et al. (2020) ¹⁵
MARINE				
Mixed assemblage			INC, OMIC	Karl et al. (2008) ¹⁷
Bacteria	Prokaryote		INC, STAT	Damm et al. (2010) ²⁵
Cyanobacteria	Prokaryote		INC	White et al. (2010) ²⁶
α-Proteobacteria	Prokaryote		INC, OMIC	Carini et al. (2014) ²⁷
Haptophytes	Eukaryote		INC, ISO	Lenhart et al. (2016) ²⁸
Bacteria	Prokaryote		INC, OMIC	Repeta et al. (2016) ²⁹
Haptophytes	Eukaryote		INC, ISO	Klitzsch et al. (2019) ³⁰
Cyanobacteria	Prokaryote		INC, ISO	Bizic et al. (2020) ¹⁵
γ-Proteobacteria	Prokaryote		INC	Ye et al. (2020) ³²
Haptophytes	Eukaryote		INC	Klitzsch et al. (2020) ³¹
FRESHWATER				
Methanogens, algae	Archaea, Eukaryote	38–58 nmol l ⁻¹ day ⁻¹ (Lake Stechlin)	INC	Grossart et al. (2011) ¹¹
Methanogens, algae	Archaea, Eukaryote	210–240 nmol l ⁻¹ day ⁻¹ (Lake Cromwell)	ISO, MB	Bogard et al. (2014) ³³
α-, γ-proteobacteria	Prokaryote		INC, OMIC	Yao et al. (2016) ¹³
Mixed assemblage		110 nmol l ⁻¹ day ⁻¹ (Lake Hallwil)	MB	Donis et al. (2017) ⁷
γ-Proteobacteria	Prokaryote	0.2–0.7 nmol l ⁻¹ day ⁻¹ (Yellowstone Lake)	INC, ISO, OMIC	Wang et al. (2017) ³⁴
Mixed assemblage			ISO, MB, PHYS	DelSontro et al. (2018) ³⁵
Proteobacteria	Prokaryote	54–257 nmol l ⁻¹ day ⁻¹ (Lake Bonney)	INC, OMIC	Li et al. (2019) ³⁶
Cyanobacteria	Prokaryote		INC, OMIC	Khatun et al. (2019) ³⁷
Mixed assemblages		72–88 nmol l ⁻¹ day ⁻¹ (Lake Stechlin)	MB	Günthel et al. (2019) ³⁹
		78–138 nmol l ⁻¹ day ⁻¹ (Lake Hallwil)		
Cyanobacteria	Prokaryote		INC, ISO	Bizic et al. (2020) ¹⁵
Cyanobacteria	Prokaryote		STAT	Khatun et al. (2020) ³⁸
Green algae, diatoms, cryptophytes	Eukaryote	50–210 nmol l ⁻¹ day ⁻¹ (Lake Stechlin)	INC, ISO, MB, STAT	Hartmann et al. (2020) ¹⁸
Picoeukaryotes, diatoms	Eukaryote		STAT	Leon-Palmero et al. (2020) ⁴¹
Proteobacteria	Prokaryote	24–547 nmol l ⁻¹ day ⁻¹ (5 Lakes)	INC, ISO, OMIC	Perez-Coronel and Beman (2020) ⁴²

OMP evidence type: INC incubation experiments, ISO isotope techniques, MB mass balance approaches, OMIC molecular biological methods, PHYS physical modelling, STAT statistical analyses. OMP has been observed in different limnic systems, e.g. temperate and arctic regions (DelSontro et al. 2018³⁵, Li et al. 2019)³⁶, high-elevation (Perez-Coronel and Beman, 2020)⁴², and throughout the oligo-to-eutrophic nutrient spectrum (DelSontro et al., 2018³⁵, Khatun et al., 2020³⁸, Ye et al., 2020)³².

methane input of 3 mmol m⁻² d⁻¹ when combined with the diffusive flux, which is higher than the value assumed by Peeters and Hofmann. Vertical diffusive input was calculated from empirically measured methane concentration profiles and turbulent diffusivities. We parameterised methane oxidation as 30% of internal production for Lake Stechlin; in a sensitivity analysis, we evaluated this assumption and also considered the most conservative scenario, e.g., OMP set to minimum. For Lake Hallwil, methane oxidation rates were measured by experiments.

By balancing the different input and output fluxes, we produced the first system-wide OMP estimate for Lake Stechlin, which agrees well with direct bottle incubation measurements reported earlier¹¹. To further account for (seasonal) variabilities and measurement uncertainties, we conducted Monte Carlo simulations and sensitivity analysis applying various conservative scenarios to the mass balance. It is, however, worth noting that the mass balance is sensitive to the flux parameterisation and the accuracy of its result is hinged on how

reliably one accounts for these fluxes. To better resolve OMP and allow for more general and firm statements about OMP (including different lake systems), future studies should aim to reduce uncertainties associated with the littoral methane input (e.g. methodological uncertainty in sediment core measurements¹²) and methane oxidation—two key parameters in the epilimnetic methane budget.

OMP by diverse organisms (Table 1) and pathways^{13–15} point to its wider potential relevance on a global scale. To examine how OMP may vary according to lake characteristics, we combined our results with analysis of literature data to estimate OMP contribution in relation to basin morphometry (Supplementary Note 5). The epilimnetic methane sources considered here are littoral sediment and OMP. On a whole-system level, the relative contributions of these sources are proportional to the total littoral sediment area and the epilimnion volume, respectively. Because the ratio of littoral sediment area to epilimnion volume decreases with increasing lake size, the contribution of OMP to surface emission is

expected to increase with lake size. This trend does not change even when we assume a larger littoral sediment area by decreasing the sediment slope as suggested by Peeters and Hofmann (Fig. 1). As the current OMP dataset is limited to only a few lakes (four data points based on mass balance and seven based on transport modelling), future studies should aim to increase the number and types of lakes to verify the trend on a larger scale.

Note, as Peeters' and Hofmann's re-analysis excludes internal methane modulation, their OMP estimates reflect net rates while our study presents gross rates. Accordingly, their contribution pattern of oxic versus anoxic methane source to surface emission (NOMC) cannot be directly compared to our estimates (OMC) (further discrepancy is explained by Supplementary Note 5).

Oxic methane production defies the century-old teaching of anoxic methanogenesis and the convention of considering only anoxic sources in methane research; as such, skepticism is expected. While some may dismiss OMP as irrelevant¹⁶, others take a more practical approach and investigate the phenomenon at the ecological, organismal, and molecular levels^{13,17}. However, the novelty of OMP also means researchers are still trialling different methods, each with their limitations (Table 2).

A better understanding of production, storage, consumption, and distribution processes of methane, including methane produced in oxic environments, is needed to improve the assessment of the global methane cycle. This requires better spatio-temporal data resolution and better constraints of data uncertainties by using multiple methods. For instance, OMP rates determined by bottle incubations can complement results




Fig. 1 Oxic methane contribution (OMC) to surface emission in relation to lake morphology. Comparison of (a) the original relationship and (b) the alternative parameterisation using a smaller sediment slope angle. A_{sed} is the littoral sediment area and V is the surface mixed layer volume. Note, OMC is defined as in our original study; the x-axis is linearly scaled, and the y-axis is scaled to $\log_{2.7}$.

Table 2 Overview on approaches to investigate oxic methane production (OMP) in lake waters.			
Approach	Description	Caveats	Reference examples
Incubation of • Lake water • Enrichment cultures	Cultivating microbes in closed containers and recording CH_4 concentration over time. Additionally, the change in $^{13}C/^{12}C$ carbon isotope ratio in dissolved methane can be measured.	Bottle enclosure may alter the light and nutrient conditions versus in situ. Long-term incubations (exceeding hours) may not reflect in situ conditions due to changes to the production-consumption equilibrium (e.g., nutrient depletion, community alterations).	Grossart et al. (2011) ¹¹ , Bizic et al. (2020) ¹⁵ , Günthel et al. (2020) ⁴⁰ , Hartmann et al. (2020) ¹⁸ , Klintzsch et al. (2019, 2020) ^{30,31}
Metagenomics	Molecular analysis of relevant enzyme machinery or genes.	Qualitative evidence. Presence of relevant genes and enzymes indicates production potential, but actual production rate can be affected by inhibitors, missing precursors, unfavourable conditions, epigenetic modulation, etc.	Carini et al. (2014) ²⁷ , Yao et al. (2016) ¹³ , Perez-Coronel and Beman (2020) ⁴²
Statistical analysis	Methane concentration is measured together with other lake parameters. Statistical models are applied to test for correlative significance and predictive power.	Individual methane sources and sinks can be overlooked due to the complex lake water methane cycling. Results lack mechanistic understanding of the underlying processes.	Fernandez et al. (2016) ³ , Günthel et al. (2020) ⁴⁰ , Khatun et al. (2020) ³⁸ , Leon-Palmero et al. (2020) ⁴¹
Physical modelling	Combining physical mechanistic aspects with correlative analysis.	Underrepresentation of internal biological modulation (oxidation and OMP).	Peeters et al. (2019) ¹⁶
Mass balance of epilimnion in • Whole-lake basin or • Mesocosms/enclosures	Methane input and output fluxes for the epilimnion are experimentally determined and balanced. Discrepancy is attributed to OMP.	Accuracy of OMP production rates depends on how reliably methane fluxes have been determined. Spatio-temporal data resolution is often limited.	Bogard et al. (2014) ³³ , Donis et al. (2017) ⁷ , Günthel et al. (2019) ³⁹ , Peeters et al. (2019) ¹⁶ , Hartmann et al. (2020) ¹⁸
Methane isotope analysis • Comparing ambient signatures or • Isotope budgets	Analysing carbon (and hydrogen) stable isotope signatures of methane sources and considering isotope fractionation by biochemical and physical reactions (e.g., oxidation, OMP, phase exchange). Analogue to mass balance.	This analysis requires knowing (i) the quantity of all mass fluxes, (ii) isotope characteristics of all methane sources, (iii) isotope fractionation by biochemical and physical processes. Different precursors and biochemical production/consumption pathways can result in different isotope signatures.	Tang et al. (2014) ⁹ , DelSontro et al. (2018) ³⁵ , Günthel et al. (2020) ⁴⁰ , Hartmann et al. (2020) ¹⁸ , Tsunogai et al. (2020) ¹⁹

based on mass budgets, as we did in our study. The incorporation of methane carbon¹⁸ and hydrogen¹⁹ isotope data into mass budgets is a promising way to further tease apart the different methane sources. Omic approaches can be used to investigate the different OMP pathways and the organisms involved.

We have discussed the caveats of our mass balance analysis, such as the limited amount of OMP and littoral flux data, limited types of lakes being considered, and the influence by other compounding factors. The global significance of OMP can only be fully assessed when more relevant data become available, but this also requires researchers to look beyond the anoxic paradigm and consider OMP in future methane measurements. We hope our and others' work will continue to stimulate more research and constructive discussions on this topic.

Data availability

Data are made available in graphical or tabular form throughout the paper and Supplementary Information. Source data are provided with this paper.

Received: 30 January 2020; Accepted: 11 January 2021;

Published online: 22 February 2021

References

- Bastviken, D., Cole, J., Pace, M. & Tranvik, L. Methane emissions from lakes: dependence of lake characteristics, two regional assessments, and a global estimate. *Glob. Biogeochem. Cycles* **18**, GB4009 (2004).
- Murase, J., Sakai, Y., Kametani, A. & Sugimoto, A. Dynamics of methane in mesotrophic Lake Biwa, Japan. *Ecol. Res.* **20**, 377–385 (2005).
- Fernandez, J. E., Peeters, F. & Hofmann, H. On the methane paradox: transport from shallow water zones rather than in situ methanogenesis is the major source of CH₄ in the open surface water of lakes. *J. Geophys. Res. Biogeosci.* **121**, 2717–2726 (2016).
- Li, Y., Fichot, C. G., Geng, L., Scarratt, M. G. & Xie, H. The contribution of methane photoproduction to the oceanic methane paradox. *Geophys. Res. Lett.* **47**, e2020GL088362 (2020).
- Schilder, J., Bastviken, D., van Hardenbroek, M. & Heiri, O. Spatiotemporal patterns in methane flux and gas transfer velocity at low wind speeds: Implications for upscaling studies on small lakes. *J. Geophys. Res. Biogeosci.* **121**, 1456–1467 (2016).
- Klaus, M. & Vachon, D. Challenges of predicting gas transfer velocity from wind measurements over global lakes. *Aquat. Sci.* **82**, 53 (2020).
- Donis, D. et al. Full-scale evaluation of methane production under oxic conditions in a mesotrophic lake. *Nat. Commun.* **8**, 1661 (2017).
- Casper, P. Methane production in littoral and profundal sediments of an oligotrophic and a eutrophic lake. *Arch. Hydrobiol. Spec. Issues Adv. Limnol.* **48**, 253–259 (1996).
- Tang, K. W. et al. Paradox reconsidered: Methane oversaturation in well-oxygenated lake waters. *Limnol. Oceanogr.* **59**, 275–284 (2014).
- Flury, S., McGinnis, D. F. & Gessner, M. O. Methane emissions from a freshwater marsh in response to experimentally simulated global warming and nitrogen enrichment. *J. Geophys. Res.* **115**, G01007 (2010).
- Grossart, H.-P., Frindt, K., Dzallas, C., Eckert, W. & Tang, K. W. Microbial methane production in oxygenated water column of an oligotrophic lake. *Proc. Natl Acad. Sci. USA* **108**, 19657–19661 (2011).
- Bussmann, I. Methane release through resuspension of littoral sediment. *Biogeochemistry* **74**, 283–302 (2005).
- Yao, M. C., Henny, C. & Maresca, J. A. Freshwater bacteria release methane as a by-product of phosphorus acquisition. *J. Appl. Environ. Microbiol.* **82**, 6994–7003 (2016).
- Angle, J. C. et al. Methanogenesis in oxygenated soils is a substantial fraction of wetland methane emissions. *Nat. Commun.* **8**, 1567 (2017).
- Bizic, M. et al. Aquatic and terrestrial cyanobacteria produce methane. *Sci. Adv.* **6**, eaax5343 (2020).
- Peeters, F., Fernandez, E. J. & Hofmann, H. Sediment fluxes rather than oxic methanogenesis explain diffusive CH₄ emissions from lakes and reservoirs. *Sci. Rep.* **9**, 243 (2019).
- Karl, D. et al. Aerobic production of methane in the sea. *Nat. Geosci.* **1**, 473–478 (2008).
- Hartmann, J. F. et al. High spatiotemporal dynamics of methane production and emission in oxic surface water. *Environ. Sci. Technol.* **54**, 1451–1463 (2020).
- Tsunogai, U. et al. Dual stable isotope characterization of excess methane in oxic waters of a mesotrophic lake. *Limnol. Oceanogr.* **9999**, 1–16 (2020).
- Keppler, F., Hamilton, J., Brass, M. & Röckmann, T. Methane emissions from terrestrial plants under aerobic conditions. *Nature* **439**, 187–191 (2006).
- Messenger, D. J., McLeod, A. R. & Fry, S. C. The role of ultraviolet radiation, photosensitizers, reactive oxygen species and ester groups in mechanisms of methane formation from pectin. *Plant Cell Environ.* **32**, 1–9 (2009).
- Angel, R., Matthies, D. & Conrad, R. Activation of methanogenesis in arid biological soil crusts despite the presence of oxygen. *PLoS ONE* **6**, e20453 (2011).
- Lenhart, K. et al. Evidence for methane production by saprotrophic fungi. *Nat. Commun.* <https://doi.org/10.1038/ncomms2049> (2012).
- Althoff, F. et al. Abiotic methanogenesis from organosulphur compounds under ambient conditions. *Nat. Commun.* **5**, 4205 (2014).
- Damm, E. et al. Methane production in aerobic oligotrophic surface water in the central Arctic Ocean. *Biogeosciences* **7**, 1099–1108 (2010).
- White, A. E., Karl, D. M., Björkman, K. M., Beversdorf, L. J. & Letelier, R. M. Phosphonate metabolism by Trichodesmium IMS101 and the production of greenhouse gases. *Limnol. Oceanogr.* **55**, 1755–1767 (2010).
- Carini, P., White, A., Campbell, E. & Giovannoni, S. J. Methane production by phosphate-starved SAR11 chemoheterotrophic marine bacteria. *Nat. Commun.* **5**, 4346 (2014).
- Lenhart, K. et al. Evidence for methane production by the marine algae *Emiliania huxleyi*. *Biogeosciences* **13**, 3163–3174 (2016).
- Repeta, D. J. et al. Marine methane paradox explained by bacterial degradation of dissolved organic matter. *Nat. Geosci.* **9**, 884–887 (2016).
- Klitzsch, T. et al. Methane production by three widespread marine phytoplankton species: release rates, precursor compounds, and potential relevance for the environment. *Biogeosciences* **16**, 4129–4144 (2019).
- Klitzsch, T. et al. Effects of temperature and light on methane production of widespread marine phytoplankton. *J. Geophys. Res. Biogeosci.* **125**, e2020JG005793 (2020).
- Ye, W. W., Wang, X. L., Zhang, X. H. & Zhang, G. L. Methane production in oxic seawater of the western North Pacific and its marginal seas. *Limnol. Oceanogr.* **65**, 2352–2365 (2020).
- Bogard, M. J. et al. Oxic water column methanogenesis as a major component of aquatic CH₄ fluxes. *Nat. Commun.* **5**, 5350 (2014).
- Wang, Q., Dore, J. E. & McDermott, T. R. Methylphosphonate metabolism by *Pseudomonas* sp. populations contributes to the methane oversaturation paradox in an oxic freshwater lake. *Environ. Microbiol.* **19**, 2366–2378 (2017).
- DelSontro, T., del Giorgio, P. A. & Prairie, Y. T. No longer a paradox: The interaction between physical transport and biological processes explains the spatial distribution of surface water methane within and across lakes. *Ecosystems* **21**, 1073–1087 (2018).
- Li, W. et al. Methane production in the oxygenated water column of a perennially ice-covered Antarctic lake. *Limnol. Oceanogr.* **65**, 143–156 (2019).
- Khatun, S. et al. Aerobic methane production by planktonic microbes in lakes. *Sci Total Environ* **696**, 133916 (2019).
- Khatun, S. et al. Linking stoichiometric organic carbon–nitrogen relationships to planktonic cyanobacteria and subsurface methane maximum in deep freshwater lakes. *Water* **12**, 402 (2020).
- Günthel, M. et al. Contribution of oxic methane production to surface methane emission in lakes and its global importance. *Nat. Commun.* **10**, 5497 (2019).
- Günthel, M. et al. Photosynthesis-driven methane production in oxic lake water as an important contributor to methane emission. *Limnol. Oceanogr.* **65**, 2853–2865 (2020).
- Leon-Palmero, E., Contreras-Ruiz, A., Sierra, A., Morales-Baquero, R. & Reche, I. Dissolved CH₄ coupled to photosynthetic picoeukaryotes in oxic waters and to cumulative chlorophyll *a* in anoxic waters of reservoirs. *Biogeosciences* **17**, 3223–3245 (2020).
- Perez-Coronel, E. & Beman, J. M. Biogeochemical and omic evidence for multiple paradoxical methane production mechanisms in freshwater lakes. *bioRxiv* <https://doi.org/10.1101/2020.07.28.225276> (2020).

Acknowledgements

We thank Frank Peeters and Hilmar Hofmann for taking an interest in our study and raising several interesting discussion points about oxic methane production.

Author contributions

M.G., D.D., G.K. and D.F.M. analysed the data. M.G., D.D., G.K., D.I., M.B., D.F.M., H.-P.G. and K.W.T. discussed and wrote the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41467-021-21216-1>.

Correspondence and requests for materials should be addressed to M.G., D.F.M., H.-P.G. or K.W.T.

Peer review information *Nature Communications* thanks John Melack and other, anonymous, reviewers for their contributions to the peer review of this work.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021