



Article scientifique

Article

2019

Published version

Open Access

This is the published version of the publication, made available in accordance with the publisher's policy.

---

## Signaling cascades governing entry into and exit from host cells by *Toxoplasma gondii*

---

Bisio Sabaris, Hugo; Soldati-Favre, Dominique

### How to cite

BISIO SABARIS, Hugo, SOLDATI-FAVRE, Dominique. Signaling cascades governing entry into and exit from host cells by *Toxoplasma gondii*. In: Annual Review of Microbiology, 2019, vol. 73, p. 579–599. doi: 10.1146/annurev-micro-020518-120235

This publication URL: <https://archive-ouverte.unige.ch/unige:123112>

Publication DOI: [10.1146/annurev-micro-020518-120235](https://doi.org/10.1146/annurev-micro-020518-120235)

© The author(s). This work is licensed under a Creative Commons Attribution-ShareALike (CC BY-SA)

<https://creativecommons.org/licenses/by-sa/4.0>

*Annual Review of Microbiology***Signaling Cascades Governing  
Entry into and Exit from Host  
Cells by *Toxoplasma gondii*****Hugo Bisio and Dominique Soldati-Favre**Département de Microbiologie et Médecine Moléculaire, Centre Médical Universitaire,  
Université de Genève, 1211 Geneva 4, Switzerland; email: Hugo.Bisio@unige.ch

Annu. Rev. Microbiol. 2019. 73:579–99

The *Annual Review of Microbiology* is online at  
micro.annualreviews.org<https://doi.org/10.1146/annurev-micro-020518-120235>Copyright © 2019 by Annual Reviews.  
All rights reserved**Keywords**Apicomplexa, *Toxoplasma gondii*, motility, egress, microneme, signaling, calcium, cyclic nucleotides, phosphatidic acid, kinase, diacylglycerol kinase, guanylate cyclase**Abstract**

The Apicomplexa phylum includes a large group of obligate intracellular protozoan parasites responsible for important diseases in humans and animals. *Toxoplasma gondii* is a widespread parasite with considerable versatility, and it is capable of infecting virtually any warm-blooded animal, including humans. This outstanding success can be attributed at least in part to an efficient and continuous sensing of the environment, with a ready-to-adapt strategy. This review updates the current understanding of the signals governing the lytic cycle of *T. gondii*, with particular focus on egress from infected cells, a key step for balancing survival, multiplication, and spreading in the host. We cover the recent advances in the conceptual framework of regulation of microneme exocytosis that ensures egress, motility, and invasion. Particular emphasis is given to the trigger molecules and signaling cascades regulating exit from host cells.

**ANNUAL  
REVIEWS CONNECT**[www.annualreviews.org](http://www.annualreviews.org)

- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media

## Contents

INTRODUCTION .....	580
ROLE OF MICRONEME SECRETION IN EGRESS .....	581
SIGNALING CASCADE TRIGGERING MICRONEME SECRETION .....	583
NATURAL EGRESS RESPONDING TO INTRINSIC SIGNALS .....	583
ENVIRONMENTAL SENSING OF EXTRINSIC SIGNALS .....	585
INTEGRATION OF EXTRINSIC AND INTRINSIC SIGNALS .....	586
CROSS TALK BETWEEN CYCLIC NUCLEOTIDES .....	587
PKG AND PKAs: MASTER REGULATORS OF PARASITE DECISIONS .....	588
CALCIUM HOMEOSTASIS, SIGNALING, AND SENSING .....	589
PHOSPHOLIPID HOMEOSTASIS AND PHOSPHATIDIC ACID SIGNALING .....	592
FUTURE DIRECTIONS .....	592

## INTRODUCTION

*Toxoplasma gondii* belongs to the phylum Apicomplexa, which includes a large group of obligate intracellular protozoan parasites responsible for important diseases in humans and animals. In the spotlight, *Plasmodium* spp., the etiological agent of malaria, are responsible for half a million deaths per year. *Cryptosporidium*, another member of the phylum, is the second-most important severe-diarrhea-causing agent in children after retroviruses (74). The focus of this review, *T. gondii*, is a medically important opportunistic pathogen responsible for toxoplasmosis encephalitis in immune-compromised patients upon reactivation of chronic infection (40). Moreover, primary infection during pregnancy can lead to abortion or congenital disease including ocular toxoplasmosis, the most common cause of eye inflammation worldwide (138).

Together with the ciliates and dinoflagellates, the apicomplexans belong to the superphylum of Alveolata, which shares the peripheral alveolar membrane system below the plasma membrane (PM), termed the inner membrane complex (IMC), as well as an elaborate apical complex including specialized secretory organelles known as micronemes and rhoptries (46). As an obligate intracellular parasite, *T. gondii* actively invades host cells and resides within a vacuole surrounded by a unique parasitophorous vacuole membrane (PVM), where it replicates safely. This parasite shares a unique mode of substrate-dependent motility crucial for egress and invasion. The exocytosis of the micronemes coupled to the activation of the actomyosin system elicits effective parasite egress from infected cells, whereas reinvasion involves the coordinated discharge of both micronemes and rhoptries (47).

Recent studies on the signaling pathways and the implicated second messengers underpinning egress have facilitated a broader understanding of the interplay between various signals that govern this complex cascade of events. The cyclic GMP (cGMP) and intracellular calcium ( $\text{Ca}^{2+}$ ) acting on protein kinase G (PKG) and calcium-dependent protein kinases (CDPKs), respectively, are key players controlling microneme secretion. Additionally phosphatidic acid (PA) is a lipid mediator that contributes in a coordinating manner to microneme secretion and gliding motility (23).

The aim of this review is to integrate current knowledge about the signaling pathways leading to microneme secretion and egress from infected cells, with particular focus on *T. gondii*.

## ROLE OF MICRONEME SECRETION IN EGRESS


Microneme exocytosis crucially participates in three key steps of the parasite lytic cycle (13): egress, gliding motility, and invasion (26). Egress from infected cells depends on multiple highly synchronized processes including conoid (an apical organelle composed of tubulin fibers) protrusion, microneme secretion, actomyosin activation, and separation of parasites at their basal pole (**Figure 1**).

The micronemes are apical specialized organelles that exocytose their content in a tightly regulated fashion, releasing perforins, lipases, proteases, and adhesins that are critical for egress, gliding, and invasion (71, 113). Most notably, perforin-like protein 1 (TgPLP1) ensures the lysis of the PVM and host PM (69). Micronemes also ensure gliding motility via the delivery of adhesins at the PM of parasites that participate in a coordinated fashion with the actomyosin system to generate forward motion. These adhesins are typically termed MICs and exist as complexes that are discharged at the apical tip of the parasite. Some of these complexes bind to host cell receptors or the extracellular matrix and are translocated to the basal pole (64, 112). Most of these MICs undergo pre- and postexocytosis proteolytic cleavage, which includes post-Golgi aspartyl protease 3-mediated maturation (37), cell surface processing by subtilisin protease 1 (78), and shedding after secretion performed by rhomboid proteases (38).

The actomyosin system of *T. gondii* is composed of an essential actin (TgACT1) and 11 myosins that are implicated in various processes including parasite motility (MyoA, MyoC, and MyoH), organelle trafficking and positioning (MyoF), basal pole constriction, and intravacuolar cell-cell communication (MyoI). The substrate-dependent motility of apicomplexan parasites is powered by a phylum-specific molecular machine called the glideosome that lies underneath the PM in the pellicular space and is anchored in the IMC. According to the capping model, gliding relies on the rearward translocation of apically discharged MICs powered by the glideosome in order to propel the parasite forward (15, 118). The components and assembly of the glideosome have been recently reviewed, in Reference 47. According to the model, filamentous actin (F-actin) and the cytoplasmic tails of the adhesins are connected via a recently discovered protein called the glideosome-associated connector (GAC) (66). Concordantly, GAC is indispensable for gliding motility, invasion, and egress and binds both F-actin and the cytoplasmic tail of MIC2.

Successful egress also depends on the fission of the recently identified network that connects intravascular parasites at their basal pole. This cell-cell communication accounts for synchronous division of *T. gondii* during acute infection (48, 107). This process resembles *Plasmodium* blood stage division strategy via schizogony, suggesting its evolutionary conservation. During egress the basal F-actin disassembles and parasites very quickly disconnect and invade new cells as individual entities (48, 68, 107, 127). This coordinated timing implies a concerted signaling event that concomitantly triggers microneme exocytosis and ensures parasite separation. Treatment with cytochalasin D, an inhibitor of actin polymerization that primarily acts on parasite actin (36), or genetic deletion of *MyoI* resulted in asynchronous division and loss of connectivity between intravacuolar parasites (48, 107).

Nondividing extracellular *T. gondii* tachyzoites constitute a distinct biological state compared to nongliding, replicating intracellular parasites. Freshly egressed parasites harbor an extruded conoid, secrete micronemes, and are able to glide for several minutes and survive for a few hours (13). Once a suitable host cell is encountered, extracellular parasites attach and apically reorient. Host cell attachment results in a new burst of microneme secretion, which is immediately followed by the discharge of the rhoptries positioned at the apical tip of the parasites (26, 41). The rhoptries contain membranous material and proteins (6, 16) that are directly injected into the host cytoplasm and allow invasion, PVM formation, and subversion of host cellular functions (7, 55, 71) (**Figure 1**).



**Figure 1**

The lytic cycle of attachment, invasion, replication, and egress. Extracellular tachyzoites are capable of movement using gliding motility. (1) The exocytosis of micronemes (dark green) occurs at the apical tip of the parasite. (Top inset) As a result of this process, adhesins (MICs) are deposited at the parasite plasma membrane (pPM), where they interact with a host receptor (black) at the host plasma membrane (hPM), for example. Rearward translocation of the adhesin by actin (red) and myosin machinery (dark blue) at the interspace of the inner membrane complex (IMC) and the pPM produces a forward movement of the parasite. The glideosome-associated connector (GAC) connects the actin filaments with the cytosolic tail of the adhesins. Diverse molecules characteristic of extracellular environments (low  $K^+$  levels, pH lower than 7.2, and the presence of serum albumin) trigger gliding motility by providing signals for microneme exocytosis. (2) Once a suitable cell is detected, parasites attach, reorient by the action of rhomboid proteases, and (3) eventually discharge micronemes and rhoptries (light blue) to initiate invasion. (Bottom inset) The nature of the rhoptry discharge signal is still unknown but likely encompasses the recognition of a host cell receptor by a micronemal protein. A similar mechanism is utilized for gliding and the invasion process. One specific adhesin is used for invasion (AMA1), and the host receptor is introduced by the parasite in the hPM (RONs) during the rhoptry discharge process. (4) Once the invasion process is finished, *T. gondii* resides in the parasitophorous vacuole (PV) composed of host- and rhoptry-derived lipid and proteins, switching into the replicative mode. Parasite division is performed without completion of cytokinesis, resulting in cytosolic-connected daughter cells. This process is dependent on filamentous actin and myosin. (5) Egress is triggered by a decrease of host  $K^+$  levels or pH, an increase in host calcium levels, or accumulation of vacuolar/extracellular phosphatidic acid (ePA) and requires the coordination of microneme secretion, actomyosin motor activation, and parasite fission. During this process, microneme content destabilizes the hPM and the PV and furnishes the pPM with MICs, which will allow gliding motility. In addition, basal actin filaments collapse during the process of parasite fission and render independent parasites capable of starting a new lytic cycle. Adapted from Reference 10.

## SIGNALING CASCADE TRIGGERING MICRONEME SECRETION


The signaling cascade leading to microneme secretion initiates with the production of cGMP by a large and atypical guanylate cyclase (GC), which is fused to a P4-ATPase domain (10, 21, 140). In turn, induction of the cGMP-dependent protein kinase G (PKG) (19) induces  $\text{Ca}^{2+}$  mobilization, potentially by phosphorylating and activating key enzymes involved in the formation of phosphatidylinositol (3,4,5) triphosphate [ $\text{PI}_{(4,5)}\text{P}_2$ ].  $\text{PI}_{(4,5)}\text{P}_2$  is converted by the PM-associated phosphoinositide phospholipase C (PI-PLC) (18) into inositol triphosphate ( $\text{IP}_3$ ) and diacylglycerol (DAG) (122). In eukaryotic model organisms,  $\text{IP}_3$  is known to bind to an  $\text{IP}_3$  receptor ( $\text{IP}_3\text{R}$ ) that serves as a  $\text{Ca}^{2+}$  channel, and it releases  $\text{Ca}^{2+}$  from the endoplasmic reticulum (ER) (53). In *T. gondii*,  $\text{IP}_3$  presumably stimulates  $\text{Ca}^{2+}$  release from an unknown  $\text{Ca}^{2+}$  store (85). The rise in intracellular  $\text{Ca}^{2+}$  activates the calcium-dependent protein kinases 1 and 3 (CDPK1 and CDPK3), as well as other members of this family (86, 87).

DAG can be phosphorylated to produce PA through a reversible reaction involving diacylglycerol kinase 1 (DGK1) and phosphatidic acid phosphatases (PAPs/lipins). The accumulation of PA in the inner leaflet of the parasite PM is sensed by an acylated pleckstrin homology domain-containing protein (APH) at the surface of the micronemes (22). Ultimately, a machinery promoting membrane fusion, most likely dependent on SNARE (soluble NSF-attachment protein receptor) proteins, leads to microneme exocytosis (22, 23, 44) (**Figure 2**).

## NATURAL EGRESS RESPONDING TO INTRINSIC SIGNALS

Egress is a tightly controlled process, and several extrinsic signals (derived from the host environment) have been shown to act as triggers or mediators of microneme secretion. *T. gondii* atypically divides by endodyogeny and thus remains invasive at almost any point during the intracellular growth phase. This adaptation allows a quick response to insults of the infected host cells by the host immune system or to sudden limitation in nutrients. On the other hand, nonstimulated egress has been termed natural egress (24) and controls the maximum number of division cycles that the parasite undergoes prior to exit. Until recently it was not clear whether natural egress was dictated by an intrinsic signal (derived from the parasite), such as abscisic acid (ABA) (100), or whether the parasites would eventually exit the host cell by mechanical breakage of the PVM and host PM. ABA was proposed to regulate natural egress and conversion to latent stages in *T. gondii* via the production of cADP ribose (100). Although *T. gondii* possesses ADP ribosyl cyclase and hydrolase (31) and a putative ABA-binding G protein-coupled receptor (GPCR) (GPR89, TGGT1\_286490), no pathway for the biosynthesis of ABA has thus far been found in apicomplexans. Consequently, the absence of an assigned molecular mechanism behind natural egress has precluded addressing the relevance of this process in vivo.

Recently an alternative and plausible candidate for an intrinsic signal was unveiled. *T. gondii*, as well as the other members of the cyst-forming subgroup of coccidians, encodes a second DGK (DGK2). This protein exhibits a signal peptide, is stored in the dense granules, and is secreted into the PV space, where it accumulates. DGK2 is an active enzyme that produces PA, likely in the outer leaflet of the parasite PM but possibly also on the tubular network and PVM. Of considerable relevance, parasites with deleted *DGK2* harbor a selective defect in natural egress despite complete responsiveness to extrinsic inducers (10). Two other dense granule proteins, GRA41 and GRA22, have also been shown to contribute to natural egress, particularly by affecting  $\text{Ca}^{2+}$  homeostasis (77, 103), yet the relationship with DGK2 remains to be assessed. Intriguingly, while fast-growing tachyzoites egress after five to six rounds of division, bradyzoites remain trapped in a vacuole that evolves to form a cyst that will grow and persist for much longer periods of time. This difference suggests that stage conversion from tachyzoites to bradyzoites



(Caption appears on following page)

**Figure 2** (Figure appears on preceding page)

Signaling pathways leading to microneme exocytosis. (1) Activation of guanylate cyclase (GC) in the presence of two interacting partners, unique GC organizer (UGO) and CDC50.1, at the parasite plasma membrane produces the signaling molecule cyclic guanosine monophosphate (cGMP). The concentration of cGMP depends on production by GC and degradation to GMP by phosphodiesterases (PDEs), particularly the fitness-conferring PDE2. (2) cGMP activates protein kinase G (PKG), which in turn increases the production of phosphatidylinositol-4,5-bisphosphate [PI<sub>(4,5)</sub>P<sub>2</sub>] in *Plasmodium* spp. and potentially in other apicomplexans. PI<sub>(4,5)</sub>P<sub>2</sub> is produced by subsequent reactions of PI4K and PI4P5K. The hydrolysis of PI<sub>(4,5)</sub>P<sub>2</sub> by phosphoinositide phospholipase C (PI-PLC) produces diacylglycerol (DAG) and inositol triphosphate (IP<sub>3</sub>). (3) IP<sub>3</sub> is thought to stimulate release of calcium from the endoplasmic reticulum (ER) by an unknown calcium channel. (4) On the other hand, DAG is phosphorylated to produce phosphatidic acid (PA) by a reversible reaction catalyzed by the diacylglycerol kinase 1 (DGK1) and phosphatidic acid phosphatases (PAPs). (5) Plasma membrane PA binds to the acylated pleckstrin homology domain-containing protein (APH) and allows the double C2 domain protein (DOC2.1)-mediated fusion of the micronemes and parasite plasma membrane. This process is likely dependent on a SNARE complex. (6) Calcium is sensed by a plethora of proteins, including the calcium-dependent protein kinases (CDPKs), particularly CDPK1 and CDPK3. CDPK1 is essential for microneme exocytosis, while CDPK3 likely enhances calcium-dependent signals. (7) cGMP level is controlled by PDE2 (and possibly other PDEs), which is regulated by the activity of the protein kinase A catalytic 1 domain (PKAc1). PKAc1 is itself tightly controlled by PKA regulatory domain (PKAr), which binds adenylate cyclase (AC)-generated cyclic adenosine monophosphate (cAMP). ACβ is likely the main contributor of cAMP production during tachyzoite stages and localizes to the rhoptry by its binding to armadillo repeats-only protein (ARO) and ARO-interacting protein (AIP). [cAMP] depends also on its degradation to AMP performed by PDEs. (8) Danger signals like a drop in [K<sup>+</sup>], acidification of the parasitophorous vacuole, or a rise in host cytoplasmic [Ca<sup>2+</sup>] also lead to egress. In the absence of extrinsic signals, DGK2 produces vacuolar PA and restricts intracellular growth timing by triggering egress. Adapted from Reference 68.

also implies a reprogramming of a molecular clock governing egress during the chronic phase of infection. Other reported players associated with egress include the nucleotide-triphosphate-degrading enzymes (NTPases) secreted into the PV. These proteins can be activated by addition of dithiothreitol and are postulated to deplete host cell ATP, leading to a decrease in activity of the Na<sup>+</sup>/K<sup>+</sup>-ATPase pump, a drop in K<sup>+</sup> levels, and concomitantly, egress (123).

## ENVIRONMENTAL SENSING OF EXTRINSIC SIGNALS

*T. gondii* is ready to escape from an unhealthy host cell or dangerous environment at any point during intracellular replication, and the burning question is what is the parasite able to sense and how? The PVM is permeable to small molecules (119), allowing the diffusion of host metabolites and ions. Leakiness of the host PM leads to an outward flow of K<sup>+</sup> ions; the exposed parasite responds by promoting microneme exocytosis (49, 96). Low K<sup>+</sup> could activate protein kinase A catalytic subunit 1 (PKAc1) (96) or PI-PLC by a GPCR; these are two common targets found in model organisms (56). A GPCR activated by K<sup>+</sup> has recently been identified in *Plasmodium falciparum* (94) that corresponds to TGGT1\_262610 in *T. gondii*; however, molecular evidence of its link to cAMP levels or PI-PLC activity still remains to be established.

In extracellular parasites, serum albumin acts as a natural agonist for microneme secretion in a PKG-dependent manner (20). Importantly, sudden exposure to an acidic environment has also been shown to promote parasite gliding (55) and to stimulate microneme exocytosis (114). Intracellular parasites that have undergone approximately five to six rounds of replication experience a rapid drop in pH that appears to be concomitant to egress. Interestingly exposure to an acidic environment can overcome a high-K<sup>+</sup>-induced block in microneme secretion (114), in agreement with K<sup>+</sup> and pH being sensed by different receptors. Of relevance, perforin-dependent T cell-mediated cytotoxicity triggers *T. gondii* egress in a caspase-independent manner (108), potentially directly mediating an imbalance in host ion composition.

A rise in intracellular Ca<sup>2+</sup> originates not only from parasite internal stores but also from the extracellular environment (14). Extracellular Ca<sup>2+</sup> was shown to enhance egress, invasion, and gliding (105), and hence a still unidentified channel, blocked by nifedipine (105), is responsible

for uptake of extracellular  $\text{Ca}^{2+}$ . Interestingly, putative voltage-dependent gated  $\text{Ca}^{2+}$  channels have been identified in *T. gondii* (111) but not functionally characterized to date. Changes in the environmental level of  $\text{Ca}^{2+}$  detected by the parasite before egress could act as a signal to trigger it (12), and signals known to modulate host cytosolic  $\text{Ca}^{2+}$  have been shown to impact parasite egress, such as the cell death receptor Fas-Fas ligand that depends on caspase-mediated release of intracellular  $\text{Ca}^{2+}$  to trigger egress (108). Nitric oxide (NO) was also reported to induce egress in a BAPTA-AM-dependent manner (139). Importantly BAPTA-AM is de-esterified by the host after entry, rendering it unable to cross the PM of the parasite. NO induces heme-dependent soluble GCs in mammalian cells, leading to a rise in the host cytosol  $\text{Ca}^{2+}$  (35). A host GPCR signaling cascade leading to  $\text{Ca}^{2+}$  release and calpain-mediated cortical cytoskeleton destabilization has also been involved in efficient parasite egress (93).

Although direct correlation with changes in host  $\text{Ca}^{2+}$  is missing, attachment of *Toxoplasma*-infected monocytes to endothelial cells is sufficient to trigger egress in a coordinated manner (2), potentially providing a mechanism to minimize plasma exposure before infection of solid organs, including the brain (73). Progesterone and estradiol have also been shown to induce egress with no effect on microneme secretion of extracellular parasites (143), indicating that both hormones act on the host signaling pathways. In contrast, histamine, which is known to raise  $\text{Ca}^{2+}$  in human foreskin fibroblasts, failed to trigger parasite egress (14), thus either disproving a role of host  $\text{Ca}^{2+}$  as an egress signal or indicating that the threshold of  $\text{Ca}^{2+}$  levels needed to trigger egress is not reached by histamine stimulation.

## INTEGRATION OF EXTRINSIC AND INTRINSIC SIGNALS

*T. gondii* possesses a unique GC, while two paralogs are found in *P. falciparum* to produce cGMP. These proteins are predicted to have unusual bifunctional activity, conserved between apicomplexans and ciliates like *Paramecium* and *Tetrahymena* (80), being composed of a fusion between a predicted P-type ATPase-like domain and two GC catalytic domains. The N-terminal P-type ATPase seems indispensable for proper localization of GC to the PM (21, 50), and, though the function of this domain remains elusive, recent genetic data support an essential role of its activity in *T. gondii* (21).

In intracellular parasites, the PA produced by DGK2 might act directly on the flippase-like P4-ATPase domain or indirectly via an unknown receptor, to activate GC. Interestingly, in the gametocyte stages of the rodent malaria parasite *Plasmodium berghei*, xanthurenic acid (XA) has been identified as a potent inducer of exflagellation (8), and the production cGMP could be monitored by crude membrane extracts of gametocytes (98). Although these data suggested that XA could activate GC $\beta$ , one of the two GCs present in malaria parasites, XA failed to induce activity of the recombinantly expressed catalytic domains of GC $\beta$  (98). Consequently, either the binding for XA is outside of the catalytic domains or modulation of GC $\beta$  activity is indirect and mediated by a regulatory protein. Mammalian metabotropic glutamate receptors bind to XA and modulate cAMP levels in the cerebral cortex (45), and similar processes might regulate *Plasmodium* response to XA via GPCR to regulate GC.

Reminiscent of its flippase-like nature, the P4-ATPase domains of both *Plasmodium* and *Toxoplasma* GCs bind to cell division control 50 protein (CDC50), which acts as a chaperone controlling GC stability and localization (10, 50). Of relevance, the conditional knockdown of CDC50.1 in *T. gondii* specifically affects egress from infected cells, indicating that the trafficking of GC and the P-type ATPase activity are essential to sense PV-derived molecules to trigger egress (10). In CDC50.1-depleted parasites, BIPPO [5-benzyl-3-isopropyl-1H-pyrazolo[4,3-d]pyrimidin-7(6H)-one], a phosphodiesterase (PDE) inhibitor that raises cGMP levels by

inhibition of its consumption (22, 62), still induces microneme secretion in extracellular parasites, but it fails to trigger egress. The fact that BIPPO triggers microneme secretion from CDC50.1-depleted extracellular parasites implies they may have higher cGMP levels, relative to intracellular parasites where BIPPO fails to activate egress in the absence of CDC50.1. Since CDC50.1 directly affects the function of the unique enzyme responsible for cGMP production in *T. gondii*, these data strongly suggest that the tonic synthesis of cGMP is higher in extracellular parasites, possibly to ensure a higher rate of microneme exocytosis (10). The carboxyl terminal domain of the GC presents striking structural similarities to that of G protein-dependent adenylate cyclases (ACs) (28). However, despite the absence of biochemical evidence to support GC activity, depletion of the gene in *T. gondii* completely abolishes cGMP production (10) and can be rescued by a cell-permeable cGMP analogue (21). Moreover, heterologously expressed *P. falciparum* GC $\beta$  catalytic domains or the whole *Paramecium* enzyme possesses cGMP cycling activity (28, 80).

Remarkably the GC forms a signaling platform at the parasite PM, interacting not only with CDC50.1 but also with a large polytopic protein termed unique GC organizer (UGO) that potentially participates in sensing multiple signals (10). UGO is composed of a series of transmembrane-spanning domains at both the N-terminal and C-terminal sides that are strikingly conserved in the protein across the phylum. In contrast the large, central, and predicted ectodomain is highly divergent between and even within the apicomplexan species. Conditional depletion of UGO causes a mislocalization of the GC and a complete ablation of GC activity, hampering motility, invasion, and egress. The role of the ectodomain remains to be addressed but is likely to be relevant for sensing and integrating signals in extracellular parasites.

## CROSS TALK BETWEEN CYCLIC NUCLEOTIDES

The cyclic nucleotides cAMP and cGMP are important second messengers exhibiting pleiotropic roles in eukaryotic cells. cAMP and cGMP are produced from ATP and GTP, by ACs and GCs, respectively. Conversely, hydrolysis via cyclic nucleotide PDEs inactivates these second messengers, establishing a steady state to regulate the threshold levels for activation/repression of downstream components of the signaling cascade (**Supplemental Table 1**).

Distinctive roles have been attributed to cGMP and cAMP in *T. gondii*. The parasite possesses single genes coding for GC and PKG exclusively dedicated to motility, invasion, and egress (10, 19, 21). Remarkably, cGMP signaling via PKG is associated not only with microneme exocytosis but also with separation of daughter cells prior to egress and with activation of the actomyosin system, assessed by the measurement of the apicobasal flux of F-actin (10, 21, 127). In contrast cAMP can potentially be produced by four putative ACs to modulate the activity of three distinct PKA catalytic subunits. *T. gondii* AC $\alpha_1$ , AC $\alpha_2$ , and AC $\alpha_3$  harbor transmembrane-spanning domains reminiscent of ion channel domains (21). Individual knockouts of each of the AC $\alpha$ s are dispensable, suggesting some redundancy in their function (21, 68). Interestingly, *Plasmodium* encodes only one AC $\alpha$ , which plays a role in microneme exocytosis in *P. berghei* sporozoites, but it does not have a detectable role in the blood stages or subsequent development in the mosquito (104). *T. gondii* AC $\beta$  is an ortholog of the bicarbonate-sensitive soluble ACs and localizes to the surface of the rhoptries in complex with the acylated armadillo repeats only protein (ARO) and the interacting partner (AIP) (97). Deletion of AC $\beta$  has been shown to induce a strong phenotype (21) as well as to weakly phenocopy the restless phenotype observed upon conditional deletion of PKAc1 (68, 129) (see below). This restless phenotype refers to the parasite's inability to turn off microneme secretion and motility upon infection of a new cell (68, 129). *Plasmodium* also codes for an essential blood stage bicarbonate-sensitive AC $\beta$ , which might act as a pH sensor (3, 117). So far, it is not clear whether ACs respond directly to external or internal stimuli, but the essential

Supplemental Material >

role of the PKAc (see below) suggests that cAMP is implicated in multiple processes across the lytic cycle and life stages of *T. gondii* (68, 129) (**Figure 2**).

Cyclic nucleotides are degraded by PDEs, for which there are 18 putative genes in *T. gondii* (62). Two of these PDEs are predicted to confer fitness in *T. gondii* tachyzoites (121) and have been localized but not functionally characterized (68). Importantly, microneme exocytosis and egress (22, 87) can be induced in *T. gondii* by addition of the human PDE5 inhibitor zaprinast (22, 87) or BIPPO (22, 62). Despite their extensive use in the field, the specificity of these chemicals in apicomplexan parasites is not fully assessed. Experiments in *Plasmodium* spp. cell extracts suggest that zaprinast induces a specific rise in cGMP while BIPPO inhibits both cGMP and cAMP degradation, mostly acting on PDE $\delta$ , which possesses dual cAMP/cGMP hydrolysis activity (62, 134). *Plasmodium* possesses only 4 PDEs, PDE $\alpha$ - $\delta$ , and all of them are predicted to include transmembrane-spanning domains (3). Interestingly, these proteins present stage-specific expression patterns, with PDE $\alpha$  and PDE $\beta$  predominantly expressed in late trophozoite and schizont blood stages, PDE $\gamma$  in sporozoites, and PDE $\delta$  in gametocytes (3).

### PKG AND PKAs: MASTER REGULATORS OF PARASITE DECISIONS

In Apicomplexa, cGMP universally acts as mediator upstream of the signaling events leading to microneme secretion (22, 25, 88). In *T. gondii*, PKG has been shown to be essential for microneme secretion while dispensable for intracellular growth (10, 19, 21). The apicomplexan PKGs have four cGMP-binding domains; only three of them seem to be functional. Upon binding of cGMP, a conformational change of PKG allows exposure of the active site, enabling the phosphorylation of substrate proteins (76). Interestingly *Toxoplasma* and *Eimeria* produce two isoforms of PKG (19, 39), while only one is detected in *Plasmodium* (3). These two isoforms arise from alternative translation initiation and lead to a cytosolic shorter version of PKG and a dually acylated membrane-associated longer version of the protein (39, 130). Association of PKG with the parasite PM by acylation is both necessary and sufficient to control motility and invasion (19). In contrast, *Plasmodium* PKG seems to associate with the ER membrane but lacks acylation motifs (60).

The inhibition of PKG with either compound 1, trisubstituted pyrrole 4-[2-(4-fluorophenyl)-5-(1-methylpiperidine-4-yl)-1H-pyrrol-3-yl]pyridine, or compound 2, 4-[7-[(dimethylamino)methyl]-2-(4-fluorophenyl)imidazo[1,2-a]pyridine-3-yl]pyrimidin-2-amine, has been shown to block microneme secretion and egress of *T. gondii* tachyzoites (87, 136). Interestingly, despite the fact that compound 1 and compound 2 have similar specificity in *Plasmodium* (92), compound 2 is less specific in *T. gondii* (68), potentially targeting CDPKs in addition to PKG. New-generation compounds have also been developed to target *Plasmodium* (4) or *T. gondii* (120) PKG and have proved to be attractive potential drugs against apicomplexan parasites.

PKAc is regulated by fluctuating levels of cAMP. The inactive enzyme is composed of a catalytic subunit (PKAc) bound to a regulatory subunit (PKAr) (126) that contains two cAMP-binding domains. Upon cAMP binding to the PKAr subunit, the complex disassembles and leads to the release of an active PKAc. The lack of the N-terminal dimerization domain in apicomplexan PKAr suggests that the holoenzyme is likely composed of one subunit of PKAc and one subunit of PKAr, contrasting with the mammalian relatives that assemble in a tetramer disposition (57, 75).

*T. gondii* encodes three PKAc subunits (PKAc1-3) and only one PKAr subunit. PKAc1 has recently been characterized in *T. gondii*, and its binding to PKAr has been confirmed (68, 125, 129). PKAc1 localizes to the parasite periphery upon binding of the double dually acylated PKAr by myristoylation and palmitoylation, and it is released soluble and active upon increase in cAMP concentration (68). Previous reports using the PKA-specific inhibitor H89 (82) have suggested

a role of PKA regulating parasite cell division (75). Concordantly, conditional overexpression of PKAc1 or conditional depletion of PKAr leads to a defect in intracellular growth (68). Remarkably, PKAc1 inactivation by gene transcript depletion or overexpression of a cAMP-insensitive mutant PKAr results in acidification-dependent or DGK2-dependent premature egress followed by successive restless invasion events leading to host cell destruction (10, 68, 129). The restless phenotype correlates with the inability of a PKAc1-depleted parasite to dump calcium signaling upon invasion and switch from the motile to the replicative stage. This phenotype does not resemble transmigration since rhoptry discharge and formation of a moving junction occur normally (129). Moreover, a decrease in cAMP production levels mimics the cell destruction phenotype (68). Compound 1 blocks premature egress induced by PKAc1 downregulation, illustrating important cross talk between PKA and PKG signaling pathways (68). Concordantly, phosphoproteomic studies have revealed changes in a putative cGMP-specific PDE upon PKAc1 inactivation, suggesting that PKAc1 could regulate cGMP levels in order to control natural egress (68).

PKAc2 also localizes to the parasite periphery, but it is preferentially expressed during the sexual stages in felids (125). Given the comparable localization with PKAc1, PKAc2 might have similar functions in the definitive host stages of the parasite.

In contrast, PKAc3 localizes to the cytosol, indicating that the spatial control of cAMP levels and PKAc isoforms is crucial to differentially regulating PKAc activity (125). Depletion of PKAc3 leads to an increased conversion of tachyzoites to bradyzoites, indicating that a threshold level of cAMP is central to maintaining tachyzoite commitment (125). Concordantly, treatment of tachyzoites with 3-isobutyl-1-methylxanthine, a PDE inhibitor that causes prolonged elevation of cAMP levels (72), causes bradyzoite repression (42). Since transient elevation of cAMP produced by forskolin induces bradyzoite differentiation (72), the level and prolongation of the cAMP signal appears crucial for transition decisions.

Overall, PKA should be seen as a master regulator of parasite transitions, a potential hub to incorporate environmental signals sensed directly or indirectly by ACs to govern important parasite decisions.

## CALCIUM HOMEOSTASIS, SIGNALING, AND SENSING

Recent advances in developing genetically encoded  $\text{Ca}^{2+}$  indicators have offered powerful tools to visualize  $\text{Ca}^{2+}$  throughout the life cycle of the parasite (14). This technology has been exploited to identify novel compounds modulating  $\text{Ca}^{2+}$  signaling via a cell-based phenotypic screening (120). PKG activation is associated with  $\text{Ca}^{2+}$  mobilization and PI-PLC-dependent hydrolysis of  $\text{PI}_{(4,5)}\text{P}_2$  to generate DAG and  $\text{IP}_3$ . The current model suggests that  $\text{IP}_3$  mediates  $\text{Ca}^{2+}$  release from intracellular stores and regulates diverse functions, among them conoid protrusion (11, 34) and microneme exocytosis (1, 88, 135). PI-PLC acts as a signaling node, involving upstream signaling events related to cyclic nucleotide regulation, environmental cues, DAG/PA interconversion, and  $\text{IP}_3$ -associated  $\text{Ca}^{2+}$  release (22, 43). PI-PLC has been localized to the parasite periphery and cytosol in intracellular parasites and accumulates apically in extracellular parasites (22, 43). Recent studies have also shown that PI-PLC colocalizes with centrin 2 at the apical end (61). Intriguingly, despite its clear implication in microneme secretion, conditional depletion of PI-PLC results in fast cell death, likely due to errant lipid signaling (22, 43), implicating a broader role for PI-PLC activity in PM integrity. The same is true for DGK1, for which depletion causes an imbalance between DAG and PA with rapid loss of PM integrity and parasite lysis (22).

Downstream of PI-PLC activity is  $\text{IP}_3$  production and the ensuing release of  $\text{Ca}^{2+}$ , a process sensitive to  $\text{IP}_3\text{R}$  inhibitors (88).  $\text{IP}_3\text{Rs}$  are typically located in the ER and open in response to  $\text{IP}_3$  to mobilize  $\text{Ca}^{2+}$  (5). Alcohols such as ethanol and acetaldehyde, as well as the ryanodine receptor

Supplemental Material >

(RyR) agonists caffeine and ryanodine, efficiently stimulate microneme secretion by elevating intracellular  $\text{Ca}^{2+}$  in *T. gondii* (22, 25, 88). Despite the pharmacological evidence for their existence, canonical genes encoding either  $\text{IP}_3\text{R}$  or RyR are absent in apicomplexan parasites (101); however, given these data, it is likely that noncanonical  $\text{IP}_3\text{R}$  and RyR genes exist and are awaiting discovery. Following  $\text{IP}_3$  production,  $\text{Ca}^{2+}$  is rapidly mobilized (89, 131) and subsequently quenches generating of waves of cytosolic  $\text{Ca}^{2+}$  (14), which correlate with multiple rounds of microneme exocytosis (22). These waves of  $\text{Ca}^{2+}$  potentially ensure an efficient usage of the adhesins pool during gliding motility and provide a mechanism to rapidly extinguish microneme secretion and motility upon successful entry into the host cell (122). Putative  $\text{Ca}^{2+}$  stores in *T. gondii* include the ER, IMC, Golgi apparatus, mitochondrion, acidocalcisome, and plant-like vacuole (85). While the store utilized during egress is unknown, the ER is the likely  $\text{Ca}^{2+}$  contributor to triggering microneme secretion (27, 95, 99). Treatment of *T. gondii* parasites with the sarcoplasmic/ER  $\text{Ca}^{2+}$ -ATPase inhibitor thapsigargin or artemisinin has been shown to prevent reentry of  $\text{Ca}^{2+}$  in the ER and stimulate microneme secretion, presumably by blocking influx of leaked  $\text{Ca}^{2+}$  from the ER (27, 95, 99). It should also be noted that the acidocalcisomes may contribute in part to  $\text{Ca}^{2+}$  release during microneme secretion (81, 90); however, the precise role that this source may play remains incompletely defined. Interestingly, as described above, extracellular  $\text{Ca}^{2+}$  can act as a stimulator, and the molecular entity is unknown; a channel for  $\text{Ca}^{2+}$  must exist to uptake extracellular  $\text{Ca}^{2+}$  and stimulate microneme exocytosis (14).

Apicomplexan parasites lack typical  $\text{Ca}^{2+}$  effector kinases (PKC/CaMK) and instead utilize phylum-specific CDPKs (reviewed in 9, 85), which contain EF-hand motifs that consist of a helix-loop-helix structural domain binding  $\text{Ca}^{2+}$  (**Supplemental Table 2**). There are 14 CDPK genes in *T. gondii*, and several of these have been implicated in microneme-secretion-dependent events such as motility, invasion, and egress. Specifically, CDPK1 has been shown to be required for  $\text{Ca}^{2+}$ -regulated microneme secretion, thereby controlling motility, invasion, and egress from host cells (86). Consistent with its role, CDPK1 is important for *T. gondii* dissemination in vivo and reactivation of chronic infections (115). Dynamamin-related protein DrpB, which has been implicated in apical organelle biogenesis (17), is a direct substrate of CDPK1, but the importance of its phosphorylation is still unknown (84). Remarkably CDPK1 has recently been shown to control conoid protrusion and activation of the actomyosin system, leading to an apicobasal flux of F-actin independently of the contribution of CDPK1 in microneme secretion (127). Substrate candidates are members of the glideosome, which has been shown to be phosphorylated, but the relevance of these findings in vivo is still unclear (52, 65). Interestingly CDPK1 activity might be regulated by association with the protein deglycase DJ-1 in an oxidation- and  $\text{Ca}^{2+}$ -dependent manner. Lack of DJ-1 results in an upregulation of CDPK1 and prevents normal exocytosis of micronemes (30).

In contrast, CDPK3 has been shown to impact only  $\text{Ca}^{2+}$ -ionophore-induced egress, and deletion of the gene does not affect invasion, gliding, or microneme secretion in extracellular conditions (51, 87, 91). Interestingly, egress induced by permeabilization of the host cell using digitonin and low  $\text{K}^+$  buffer was significantly lower in parasites lacking CDPK3 but could be increased in the presence of ionophore treatment (91). This led the authors to conclude that CDPK3 might be important to amplify the response to a  $\text{Ca}^{2+}$  stimulus, rather than being a binary switch from intracellular to extracellular parasites. CDPK3 has also been shown to be implicated in ion homeostasis (128) and metabolism (132) and might be implicated in controlling broader aspects of *T. gondii* cell biology, preparing the cell for intracellular or extracellular conditions. Like PKG, CDPK3 is dually acylated, and its association with the pellicle is essential for its function (51, 87, 91). Overexpression of CDPK1 partially rescued the block in egress experienced by CDPK3 knockout

parasites, implicating potential cross talk between the kinases (128). A recent study reported that dense granule secretion is negatively regulated by CDPK1 and CDPK3 (70). Logically, this negative regulation might reduce the secretion of proteins that are important during the intracellular phase of the lytic cycle. While the precise targets of the CDPKs and their specific contribution to microneme secretion are yet to be fully described, rapid advances are being made in the field, and identification of specific substrates will likely elucidate the molecular events underpinning  $\text{Ca}^{2+}$  sensing and signaling leading to microneme secretion.

Although CDPKs have been by far the most-studied  $\text{Ca}^{2+}$ -regulated proteins in Apicomplexa, *T. gondii* possesses 63 additional EF-hand domain-containing proteins annotated in the genome, many of which have unknown function. Calmodulin (CaM)-like proteins constitute another important branch of 20 proteins harboring EF-hand domains (101, 110). CaM-like proteins undergo conformational changes upon  $\text{Ca}^{2+}$  binding that affect their interaction with binding partners and function (76). A conserved CaM localized to the apical and basal ring of *T. gondii* tachyzoites acts as a regulator of calcineurin, impacting parasite attachment without affecting microneme secretion (106). CaM and CaM-like proteins can serve as myosin light chains (MLCs) and associate with myosin heavy chains (58). Several MLCs have been characterized to date, with MLC1 and MLC2 associated with MyoA and MyoD, respectively (59, 137). Additional CaM-like proteins are associated with the conoid and potentially bind to MyoH to empower gliding motility, invasion, and egress (54, 83).

Similar to CaM, centrin 1 (CEN1) has been found to be at the centriole, and two other centrins have been localized to the spindle pole and centrosome, where they likely participate in cell division (63). Interestingly, centrin 2 (CEN2) localizes to the preconoidal ring and basal complex and critically participates in microneme secretion by an unknown mechanism (G. Lentini, D.J. Dubois, B. Maco, D. Soldati-Favre, K. Frénil, manuscript under revision). Two more EF-hand-containing proteins have been localized to the PM and rhoptries, while their apparently dispensable function remains unknown (29). Parafusins are also known to play a role in  $\text{Ca}^{2+}$ -mediated exocytosis (109). *T. gondii* encodes two parafusins that are involved specifically in calcium-ionophore-induced microneme secretion, again by an unknown mechanism (116).

Ultimately, SNAREs and C2-domain-containing (Ferlin-like) proteins mediate vesicle-membrane fusion in diverse cell types in response to  $\text{Ca}^{2+}$  and membrane curvature. Using a forward-genetics approach, researchers have shown that a conserved C2-domain-containing protein termed DOC2.1 plays a central role in microneme secretion, with a concomitant impact on egress, motility, and invasion in both *T. gondii* and *P. falciparum* (44). Furthermore DOC2.2 was shown to bind to  $\text{Ca}^{2+}$  in *Plasmodium* (67). Although DOC2.1 is yet to be localized in *Toxoplasma*, it has been proposed to mediate microneme release by controlling fusion of SNARE-like proteins/complexes in a  $\text{Ca}^{2+}$ -dependent manner (44). Three ferlins are encoded in the genome of *T. gondii* (32). Ferlin 2 participates in membrane fusion during rhoptry discharge, likely in a  $\text{Ca}^{2+}$ -dependent manner (32). Moreover, the homolog of *T. gondii* Ferlin 1 has been shown to be involved in gametocyte egress (102), potentially implicating Ferlin 1 in microneme exocytosis.

Although the signaling events culminating in microneme release are slowly being unraveled, little is known about the actual fusion event at the parasite PM. Membrane fusion involves a universal machinery that includes SNARE and SM (Sec1/Munc18-like) proteins. During fusion, vesicular and target SNARE proteins assemble into an  $\alpha$ -helical *trans*-SNARE complex that forces the two membranes tightly together, and SM proteins likely wrap around assembling *trans*-SNARE complexes to catalyze membrane fusion (124). This machinery is controlled by  $\text{Ca}^{2+}$  via C2-containing proteins, like ferlins and DOC2 (133, 141).

## PHOSPHOLIPID HOMEOSTASIS AND PHOSPHATIDIC ACID SIGNALING

The production of IP<sub>3</sub> by PI-PLC and concurrent Ca<sup>2+</sup> mobilization preceding microneme secretion are linked to the generation of DAG (22). Importantly DAG can be interconverted to PA through a reversible reaction that requires ATP and the action of DGKs and PAPs. As in eukaryotic model organisms, PA is linked to exocytosis (142), and the conversion of DAG to PA at the PM appears essential for microneme secretion in Apicomplexa. Concordantly, blocking PA production either with specific DGK inhibitors or through conditional depletion of the PM-associated DGK1 reduces PA levels and hampers microneme secretion in *T. gondii* (22). Conversely microneme secretion and egress can be induced by treatment of tachyzoites with the PAP inhibitor propranolol, implying that the ensuing buildup of PA facilitates exocytosis (22).

The study that delineated the importance of PA signaling at the parasite PM also revealed the presence of APH, an acylated protein at the surface of the micronemes (22). Of relevance, recombinant APH from both *Plasmodium* and *T. gondii* was shown to bind specifically to PA through both PIP-strip and liposome-binding assays, supporting the view that the protein indeed acts as a PA sensor. Furthermore, structural studies have demonstrated that APH inserts into and clusters multiple phosphate head groups at the bilayer-binding surface (33). A conditional depletion of APH resulted in a selective block in microneme secretion, leading to impairments in parasite motility, invasion, and egress (22). Based on these observations, a plausible model emerged whereby microneme-associated APH binds to PA produced by DGK1 at the inner leaflet of the parasite PM and critically participates in the signaling cascade leading to microneme exocytosis. In this context, DOC2.1 also binds to the membrane, but its lipid specificity is unknown (67) (**Supplemental Table 3**).

PA produced by DGK1 during egress is also postulated to coordinate the concerted action between adhesins and the glideosome. Indeed, GAC possesses a PH domain at the carboxyl terminus of the protein, which has been found to bind to PA and likely contributes to the positioning of GAC to the cytosolic face of the apical PM. The dual binding of GAC to MIC-tails and PA that concentrates at the parasite PM (22) presumably increases selectivity and affinity for secreted adhesins by combining two weak-binding interactions to engender strong binding (79).

## FUTURE DIRECTIONS

Recent studies have shed new light on the signaling pathways governing parasite egress from infected cells and provided critical insights into the molecular mechanisms involved. Yet many outstanding questions remain to be addressed. Is the signaling platform composed of the atypical CDC50.1, GC, and UGO, integrating both intrinsic and extrinsic signals, and if yes how? What are the molecular entities responsible for Ca<sup>2+</sup> homeostasis and acting as Ca<sup>2+</sup> channels at the parasite PM and the intracellular storage compartments? What are the substrates of the CDPK1 accounting for the essential role of this kinase in microneme secretion, conoid protrusion, and activation of R-actin flux? Is *T. gondii* relying on natural and/or induced egress to spread during chronic infection? What are the signaling events governing rhopty discharge in coordination with microneme secretion to produce the moving junction and initiate invasion?

## DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

## ACKNOWLEDGMENTS

We thank Mathieu Brochet for critical reading of the manuscript. Funding received from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program under grant agreement number 695596 and the Swiss National Foundation to D.S.-F. (FN3100A0-116722).

## LITERATURE CITED

1. Alves E, Bartlett PJ, Garcia CR, Thomas AP. 2011. Melatonin and IP3-induced  $\text{Ca}^{2+}$  release from intracellular stores in the malaria parasite *Plasmodium falciparum* within infected red blood cells. *J. Biol. Chem.* 286:5905–12
2. Baba M, Batanova T, Kitoh K, Takashima Y. 2017. Adhesion of *Toxoplasma gondii* tachyzoite-infected vehicle leukocytes to capillary endothelial cells triggers timely parasite egression. *Sci. Rep.* 7:5675
3. Baker DA, Drought LG, Flueck C, Nofal SD, Patel A, et al. 2017. Cyclic nucleotide signalling in malaria parasites. *Open Biol.* 7:170213
4. Baker DA, Stewart LB, Large JM, Bowyer PW, Ansell KH, et al. 2017. A potent series targeting the malarial cGMP-dependent protein kinase clears infection and blocks transmission. *Nat. Commun.* 8:430
5. Berridge MJ. 2009. Inositol trisphosphate and calcium signalling mechanisms. *Biochim. Biophys. Acta* 1793:933–40
6. Besteiro S, Bertrand-Michel J, Lebrun M, Vial H, Dubremetz JF. 2008. Lipidomic analysis of *Toxoplasma gondii* tachyzoites rhoptries: further insights into the role of cholesterol. *Biochem. J.* 415:87–96
7. Besteiro S, Dubremetz JF, Lebrun M. 2011. The moving junction of apicomplexan parasites: a key structure for invasion. *Cell Microbiol.* 13:797–805
8. Billker O, Lindo V, Panico M, Etienne AE, Paxton T, et al. 1998. Identification of xanthurenic acid as the putative inducer of malaria development in the mosquito. *Nature* 392:289–92
9. Billker O, Lourido S, Sibley LD. 2009. Calcium-dependent signaling and kinases in apicomplexan parasites. *Cell Host Microbe* 5:612–22
10. Bisio H, Lunghi M, Brochet M, Soldati-Favre D. 2019. Phosphatidic acid governs natural egress in *Toxoplasma gondii* via a guanylate cyclase receptor platform. *Nat. Microbiol.* 4:420–28
11. Black MW, Arrizabalaga G, Boothroyd JC. 2000. Ionophore-resistant mutants of *Toxoplasma gondii* reveal host cell permeabilization as an early event in egress. *Mol. Cell Biol.* 20:9399–408
12. Black MW, Boothroyd JC. 2000. Lytic cycle of *Toxoplasma gondii*. *Microbiol. Mol. Biol. Rev.* 64:607–23
13. Blader IJ, Coleman BI, Chen CT, Gubbels MJ. 2015. Lytic cycle of *Toxoplasma gondii*: 15 years later. *Annu. Rev. Microbiol.* 69:463–85
14. Borges-Pereira L, Budu A, McKnight CA, Moore CA, Vella SA, et al. 2015. Calcium signaling throughout the *Toxoplasma gondii* lytic cycle: a study using genetically encoded calcium indicators. *J. Biol. Chem.* 290:26914–26
15. Boucher LE, Bosch J. 2015. The apicomplexan glideosome and adhesins—structures and function. *J. Struct. Biol.* 190:93–114
16. Bradley PJ, Ward C, Cheng SJ, Alexander DL, Coller S, et al. 2005. Proteomic analysis of rhoptry organelles reveals many novel constituents for host-parasite interactions in *Toxoplasma gondii*. *J. Biol. Chem.* 280:34245–58
17. Breinich MS, Ferguson DJ, Foth BJ, van Dooren GG, Lebrun M, et al. 2009. A dynamin is required for the biogenesis of secretory organelles in *Toxoplasma gondii*. *Curr. Biol.* 19:277–86
18. Brochet M, Collins MO, Smith TK, Thompson E, Sebastian S, et al. 2014. Phosphoinositide metabolism links cGMP-dependent protein kinase G to essential  $\text{Ca}^{2+}$  signals at key decision points in the life cycle of malaria parasites. *PLoS Biol.* 12:e1001806
19. Brown KM, Long S, Sibley LD. 2017. Plasma membrane association by N-acylation governs PKG function in *Toxoplasma gondii*. *mBio* 8:e00375-17
20. Brown KM, Lourido S, Sibley LD. 2016. Serum albumin stimulates protein kinase G-dependent microneme secretion in *Toxoplasma gondii*. *J. Biol. Chem.* 291:9554–65

21. Brown KM, Sibley LD. 2018. Essential cGMP signaling in *Toxoplasma* is initiated by a hybrid P-type ATPase-guanylate cyclase. *Cell Host Microbe* 24:804–16.e6
22. Bullen HE, Jia Y, Yamaryo-Botte Y, Bisio H, Zhang O, et al. 2016. Phosphatidic acid-mediated signaling regulates microneme secretion in *Toxoplasma*. *Cell Host Microbe* 19:349–60
23. Bullen HE, Soldati-Favre D. 2016. A central role for phosphatidic acid as a lipid mediator of regulated exocytosis in apicomplexa. *FEBS Lett.* 590:2469–81
24. Caldas LA, Attias M, de Souza W. 2018. A structural analysis of the natural egress of *Toxoplasma gondii*. *Microbes Infect.* 20:57–62
25. Carruthers VB, Moreno SN, Sibley LD. 1999. Ethanol and acetaldehyde elevate intracellular  $[Ca^{2+}]$  and stimulate microneme discharge in *Toxoplasma gondii*. *Biochem. J.* 342(Part 2):379–86
26. Carruthers VB, Sibley LD. 1997. Sequential protein secretion from three distinct organelles of *Toxoplasma gondii* accompanies invasion of human fibroblasts. *Eur. J. Cell Biol.* 73:114–23
27. Carruthers VB, Sibley LD. 1999. Mobilization of intracellular calcium stimulates microneme discharge in *Toxoplasma gondii*. *Mol. Microbiol.* 31:421–28
28. Carucci DJ, Witney AA, Muhia DK, Warhurst DC, Schaap P, et al. 2000. Guanylyl cyclase activity associated with putative bifunctional integral membrane proteins in *Plasmodium falciparum*. *J. Biol. Chem.* 275:22147–56
29. Chang L, Dykes EJ, Li J, Moreno SNJ, Hortua Triana MA. 2018. Characterization of two EF-hand domain-containing proteins from *Toxoplasma gondii*. *J. Eukaryot. Microbiol.* 66:343–53
30. Child MA, Garland M, Foe I, Madzlan P, Trecek M, et al. 2017. *Toxoplasma* DJ-1 regulates organelle secretion by a direct interaction with calcium-dependent protein kinase 1. *mBio* 8:e02189-16
31. Chini EN, Nagamune K, Wetzel DM, Sibley LD. 2005. Evidence that the cADPR signalling pathway controls calcium-mediated microneme secretion in *Toxoplasma gondii*. *Biochem. J.* 389:269–77
32. Coleman BI, Saha S, Sato S, Engelberg K, Ferguson DJP, et al. 2018. A member of the ferlin calcium sensor family is essential for *Toxoplasma gondii* rhoptry secretion. *mBio* 9:e01510-18
33. Darvill N, Dubois DJ, Rouse SL, Hammoudi PM, Blake T, et al. 2018. Structural basis of phosphatidic acid sensing by APH in apicomplexan parasites. *Structure* 26:1059–71.e6
34. Del Carmen MG, Mondragón M, González S, Mondragón R. 2009. Induction and regulation of conoid extrusion in *Toxoplasma gondii*. *Cell Microbiol.* 11(6):967–82
35. Denninger JW, Marletta MA. 1999. Guanylate cyclase and the  $\cdot NO/cGMP$  signaling pathway. *Biochim. Biophys. Acta Bioenerg.* 1411:334–50
36. Dobrowolski JM, Sibley LD. 1996. *Toxoplasma* invasion of mammalian cells is powered by the actin cytoskeleton of the parasite. *Cell* 84:933–39
37. Dogga SK, Mukherjee B, Jacot D, Kockmann T, Molino L, et al. 2017. A druggable secretory protein maturase of *Toxoplasma* essential for invasion and egress. *eLife* 6:e27480
38. Dogga SK, Soldati-Favre D. 2016. Biology of rhomboid proteases in infectious diseases. *Semin. Cell Dev. Biol.* 60:38–45
39. Donald RG, Liberator PA. 2002. Molecular characterization of a coccidian parasite cGMP dependent protein kinase. *Mol. Biochem. Parasitol.* 120:165–75
40. Dubey JP. 2004. Toxoplasmosis—a waterborne zoonosis. *Vet. Parasitol.* 126:57–72
41. Dubremetz JF. 2007. Rhoptries are major players in *Toxoplasma gondii* invasion and host cell interaction. *Cell Microbiol.* 9:841–48
42. Eaton MS, Weiss LM, Kim K. 2006. Cyclic nucleotide kinases and tachyzoite-bradyzoite transition in *Toxoplasma gondii*. *Int. J. Parasitol.* 36:107–14
43. Fang J, Marchesini N, Moreno SN. 2006. A *Toxoplasma gondii* phosphoinositide phospholipase C (TgPI-PLC) with high affinity for phosphatidylinositol. *Biochem. J.* 394:417–25
44. Farrell A, Thirugnanam S, Lorestani A, Dvorin JD, Eidell KP, et al. 2012. A DOC2 protein identified by mutational profiling is essential for apicomplexan parasite exocytosis. *Science* 335:218–21
45. Fazio F, Lionetto L, Curto M, Iacovelli L, Cavallari M, et al. 2015. Xanthurenic acid activates mGlu2/3 metabotropic glutamate receptors and is a potential trait marker for schizophrenia. *Sci. Rep.* 5:17799
46. Francia ME, Dubremetz JF, Morrissette NS. 2015. Basal body structure and composition in the apicomplexans *Toxoplasma* and *Plasmodium*. *Cilia* 5:3

47. Frenal K, Dubremetz JF, Lebrun M, Soldati-Favre D. 2017. Gliding motility powers invasion and egress in Apicomplexa. *Nat. Rev. Microbiol.* 15:645–60
48. Frenal K, Jacot D, Hammoudi PM, Graindorge A, Maco B, Soldati-Favre D. 2017. Myosin-dependent cell-cell communication controls synchronicity of division in acute and chronic stages of *Toxoplasma gondii*. *Nat. Commun.* 8:15710
49. Fruth IA, Arrizabalaga G. 2007. *Toxoplasma gondii*: induction of egress by the potassium ionophore nigericin. *Int. J. Parasitol.* 37:1559–67
50. Gao H, Yang Z, Wang X, Qian P, Hong R, et al. 2018. ISP1-anchored polarization of GC $\beta$ /CDC50A complex initiates malaria ookinete gliding motility. *Curr. Biol.* 28:2763–76.e6
51. Garrison E, Treeck M, Ehret E, Butz H, Garbuz T, et al. 2012. A forward genetic screen reveals that calcium-dependent protein kinase 3 regulates egress in *Toxoplasma*. *PLoS Pathog.* 8:e1003049
52. Gilk SD, Gaskins E, Ward GE, Beckers CJ. 2009. GAP45 phosphorylation controls assembly of the *Toxoplasma* myosin XIV complex. *Eukaryot. Cell* 8:190–96
53. Gill DL, Ghosh TK, Mullaney JM. 1989. Calcium signalling mechanisms in endoplasmic reticulum activated by inositol 1,4,5-trisphosphate and GTP. *Cell Calcium* 10:363–74
54. Graindorge A, Frenal K, Jacot D, Salamun J, Marq JB, Soldati-Favre D. 2016. The conoid associated motor MyoH is indispensable for *Toxoplasma gondii* entry and exit from host cells. *PLoS Pathog.* 12:e1005388
55. Hakansson S, Charron AJ, Sibley LD. 2001. *Toxoplasma* evacuoles: a two-step process of secretion and fusion forms the parasitophorous vacuole. *EMBO J.* 20:3132–44
56. Hanlon CD, Andrew DJ. 2015. Outside-in signaling—a brief review of GPCR signaling with a focus on the *Drosophila* GPCR family. *J. Cell Sci.* 128:3533–42
57. Haste NM, Talabani H, Doo A, Merckx A, Langsley G, Taylor SS. 2012. Exploring the *Plasmodium falciparum* cyclic-adenosine monophosphate (cAMP)-dependent protein kinase (PfPKA) as a therapeutic target. *Microbes Infect.* 14:838–50
58. Heissler SM, Sellers JR. 2014. Myosin light chains: Teaching old dogs new tricks. *Bioarchitecture* 4:169–88
59. Herm-Gotz A, Weiss S, Stratmann R, Fujita-Becker S, Ruff C, et al. 2002. *Toxoplasma gondii* myosin A and its light chain: a fast, single-headed, plus-end-directed motor. *EMBO J.* 21:2149–58
60. Hopp CS, Flueck C, Solyakov L, Tobin A, Baker DA. 2012. Spatiotemporal and functional characterization of the *Plasmodium falciparum* cGMP-dependent protein kinase. *PLoS ONE* 7:e48206
61. Hortua Triana MA, Márquez-Nogueras KM, Chang L, Stasic AJ, Li C, et al. 2018. Tagging of weakly expressed *Toxoplasma gondii* calcium-related genes with high-affinity tags. *J Eukaryot. Microbiol.* 65(5): 709–21
62. Howard BL, Harvey KL, Stewart RJ, Azevedo MF, Crabb BS, et al. 2015. Identification of potent phosphodiesterase inhibitors that demonstrate cyclic nucleotide-dependent functions in apicomplexan parasites. *ACS Chem. Biol.* 10:1145–54
63. Hu K. 2008. Organizational changes of the daughter basal complex during the parasite replication of *Toxoplasma gondii*. *PLoS Pathog.* 4:e10
64. Huynh MH, Carruthers VB. 2006. *Toxoplasma* MIC2 is a major determinant of invasion and virulence. *PLoS Pathog.* 2:e84
65. Jacot D, Frenal K, Marq JB, Sharma P, Soldati-Favre D. 2014. Assessment of phosphorylation in *Toxoplasma* glideosome assembly and function. *Cell Microbiol.* 16:1518–32
66. Jacot D, Tosetti N, Pires I, Stock J, Graindorge A, et al. 2016. An apicomplexan actin-binding protein serves as a connector and lipid sensor to coordinate motility and invasion. *Cell Host Microbe* 20:731–43
67. Jean S, Zapata-Jenks MA, Farley JM, Tracy E, Mayer DC. 2014. *Plasmodium falciparum* double C2 domain protein, PDOC2, binds to calcium when associated with membranes. *Exp. Parasitol.* 144:91–95
68. Jia Y, Marq JB, Bisio H, Jacot D, Mueller C, et al. 2017. Crosstalk between PKA and PKG controls pH-dependent host cell egress of *Toxoplasma gondii*. *EMBO J.* 36:3250–67
69. Kafsack BF, Pena JD, Coppens I, Ravindran S, Boothroyd JC, Carruthers VB. 2009. Rapid membrane disruption by a perforin-like protein facilitates parasite exit from host cells. *Science* 323:530–33

70. Katris NJ, Ke H, McFadden GI, van Dooren GG, Waller RF. 2019. Calcium negatively regulates secretion from dense granules in *Toxoplasma gondii*. *Cell Microbiol.* 21:e13011
71. Kemp LE, Yamamoto M, Soldati-Favre D. 2013. Subversion of host cellular functions by the apicomplexan parasites. *FEMS Microbiol. Rev.* 37:607–31
72. Kirkman LA, Weiss LM, Kim K. 2001. Cyclic nucleotide signaling in *Toxoplasma gondii* bradyzoite differentiation. *Infect. Immun.* 69:148–53
73. Konradt C, Ueno N, Christian DA, Delong JH, Pritchard GH, et al. 2016. Endothelial cells are a replicative niche for entry of *Toxoplasma gondii* to the central nervous system. *Nat. Microbiol.* 1:16001
74. Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, et al. 2013. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet* 382:209–22
75. Kurokawa H, Kato K, Iwanaga T, Sugi T, Sudo A, et al. 2011. Identification of *Toxoplasma gondii* cAMP dependent protein kinase and its role in the tachyzoite growth. *PLOS ONE* 6:e22492
76. Kursula P. 2014. The many structural faces of calmodulin: a multitasking molecular jackknife. *Amino Acids* 46:2295–304
77. LaFavers KA, Marquez-Nogueras KM, Coppens I, Moreno SNJ, Arrizabalaga G. 2017. A novel dense granule protein, GRA41, regulates timing of egress and calcium sensitivity in *Toxoplasma gondii*. *Cell Microbiol.* 19:e12749
78. Lagal V, Binder EM, Huynh MH, Kafsack BF, Harris PK, et al. 2010. *Toxoplasma gondii* protease TgSUB1 is required for cell surface processing of micronemal adhesive complexes and efficient adhesion of tachyzoites. *Cell Microbiol.* 12(12):1792–808
79. Lemmon MA. 2008. Membrane recognition by phospholipid-binding domains. *Nat. Rev. Mol. Cell Biol.* 9:99–111
80. Linder JU, Engel P, Reimer A, Kruger T, Plattner H, et al. 1999. Guanylyl cyclases with the topology of mammalian adenylyl cyclases and an N-terminal P-type ATPase-like domain in *Paramecium*, *Tetrahymena* and *Plasmodium*. *EMBO J.* 18:4222–32
81. Liu J, Pace D, Dou Z, King TP, Guidot D, et al. 2014. A vacuolar-H<sup>+</sup>-pyrophosphatase (TgVP1) is required for microneme secretion, host cell invasion, and extracellular survival of *Toxoplasma gondii*. *Mol. Microbiol.* 93:698–712
82. Lochner A, Moolman JA. 2006. The many faces of H89: a review. *Cardiovasc. Drug Rev.* 24:261–74
83. Long S, Brown KM, Drewry LL, Anthony B, Phan IQH, Sibley LD. 2017. Calmodulin-like proteins localized to the conoid regulate motility and cell invasion by *Toxoplasma gondii*. *PLOS Pathog.* 13:e1006379
84. Lourido S, Jeschke GR, Turk BE, Sibley LD. 2013. Exploiting the unique ATP-binding pocket of *Toxoplasma* calcium-dependent protein kinase 1 to identify its substrates. *ACS Chem. Biol.* 8:1155–62
85. Lourido S, Moreno SN. 2015. The calcium signaling toolkit of the Apicomplexan parasites *Toxoplasma gondii* and *Plasmodium* spp. *Cell Calcium* 57:186–93
86. Lourido S, Shuman J, Zhang C, Shokat KM, Hui R, Sibley LD. 2010. Calcium-dependent protein kinase 1 is an essential regulator of exocytosis in *Toxoplasma*. *Nature* 465:359–62
87. Lourido S, Tang K, Sibley LD. 2012. Distinct signalling pathways control *Toxoplasma* egress and host-cell invasion. *EMBO J.* 31:4524–34
88. Lovett JL, Marchesini N, Moreno SN, Sibley LD. 2002. *Toxoplasma gondii* microneme secretion involves intracellular Ca<sup>2+</sup> release from inositol 1,4,5-triphosphate (IP<sub>3</sub>)/ryanodine-sensitive stores. *J. Biol. Chem.* 277:25870–76
89. Lovett JL, Sibley LD. 2003. Intracellular calcium stores in *Toxoplasma gondii* govern invasion of host cells. *J. Cell Sci.* 116:3009–16
90. Luo S, Ruiz FA, Moreno SN. 2005. The acidocalcisome Ca<sup>2+</sup>-ATPase (TgA1) of *Toxoplasma gondii* is required for polyphosphate storage, intracellular calcium homeostasis and virulence. *Mol. Microbiol.* 55:1034–45
91. McCoy JM, Whitehead L, van Dooren GG, Tonkin CJ. 2012. TgCDPK3 regulates calcium-dependent egress of *Toxoplasma gondii* from host cells. *PLOS Pathog.* 8:e1003066
92. McRobert L, Taylor CJ, Deng W, Fivelman QL, Cummings RM, et al. 2008. Gametogenesis in malaria parasites is mediated by the cGMP-dependent protein kinase. *PLOS Biol.* 6:e139

93. Millholland MG, Mishra S, Dupont CD, Love MS, Patel B, et al. 2013. A host GPCR signaling network required for the cytolysis of infected cells facilitates release of apicomplexan parasites. *Cell Host Microbe* 13:15–28
94. Moraes MS, Budu A, Singh MK, Borges-Pereira L, Levano-García J, et al. 2017. *Plasmodium falciparum* GPCR-like receptor SR25 mediates extracellular K<sup>+</sup> sensing coupled to Ca<sup>2+</sup> signaling and stress survival. *Sci. Rep.* 7:9545
95. Moreno SN, Zhong L. 1996. Acidocalcisomes in *Toxoplasma gondii* tachyzoites. *Biochem. J.* 313(Part 2):655–59
96. Moudy R, Manning TJ, Beckers CJ. 2001. The loss of cytoplasmic potassium upon host cell breakdown triggers egress of *Toxoplasma gondii*. *J. Biol. Chem.* 276:41492–501
97. Mueller C, Samoo A, Hammoudi PM, Klages N, Kallio JP, et al. 2016. Structural and functional dissection of *Toxoplasma gondii* armadillo repeats only protein. *J. Cell Sci.* 129:1031–45
98. Muhia DK, Swales CA, Deng W, Kelly JM, Baker DA. 2001. The gametocyte-activating factor xanthurenic acid stimulates an increase in membrane-associated guanylyl cyclase activity in the human malaria parasite *Plasmodium falciparum*. *Mol. Microbiol.* 42:553–60
99. Nagamune K, Beatty WL, Sibley LD. 2007. Artemisinin induces calcium-dependent protein secretion in the protozoan parasite *Toxoplasma gondii*. *Eukaryot. Cell* 6:2147–56
100. Nagamune K, Hicks LM, Fux B, Brossier F, Chini EN, Sibley LD. 2008. Abscisic acid controls calcium-dependent egress and development in *Toxoplasma gondii*. *Nature* 451:207–10
101. Nagamune K, Sibley LD. 2006. Comparative genomic and phylogenetic analyses of calcium ATPases and calcium-regulated proteins in the Apicomplexa. *Mol. Biol. Evol.* 23:1613–27
102. Obrova K, Cyrklaff M, Frank R, Mair GR, Mueller AK. 2018. Transmission of the malaria parasite requires ferlin for gamete egress from the red blood cell. *Cell Microbiol.* 21(5):e12999
103. Okada T, Marmansari D, Li ZM, Adilbish A, Canko S, et al. 2013. A novel dense granule protein, GRA22, is involved in regulating parasite egress in *Toxoplasma gondii*. *Mol. Biochem. Parasitol.* 189:5–13
104. Ono T, Cabrita-Santos L, Leitao R, Bettiol E, Purcell LA, et al. 2008. Adenylyl cyclase alpha and cAMP signaling mediate *Plasmodium* sporozoite apical regulated exocytosis and hepatocyte infection. *PLoS Pathog.* 4:e1000008
105. Pace DA, McKnight CA, Liu J, Jimenez V, Moreno SN. 2014. Calcium entry in *Toxoplasma gondii* and its enhancing effect of invasion-linked traits. *J. Biol. Chem.* 289:19637–47
106. Paul AS, Saha S, Engelberg K, Jiang RH, Coleman BI, et al. 2015. Parasite calcineurin regulates host cell recognition and attachment by apicomplexans. *Cell Host Microbe* 18:49–60
107. Periz J, Whitelaw J, Harding C, Gras S, Del Rosario Minina MI, et al. 2017. *Toxoplasma gondii* F-actin forms an extensive filamentous network required for material exchange and parasite maturation. *eLife* 6:e24119
108. Persson EK, Agnarson AM, Lambert H, Hitziger N, Yagita H, et al. 2007. Death receptor ligation or exposure to perforin trigger rapid egress of the intracellular parasite *Toxoplasma gondii*. *J. Immunol.* 179:8357–65
109. Plattner H, Kissmehl R. 2005. Molecular aspects of rapid, reversible, Ca<sup>2+</sup>-dependent dephosphorylation of pp63/parafusin during stimulated exo-endocytosis in *Paramecium* cells. *Cell Calcium* 38:319–27
110. Polonais V, Javier Foth B, Chinthalapudi K, Marq JB, Manstein DJ, et al. 2011. Unusual anchor of a motor complex (MyoD-MLC2) to the plasma membrane of *Toxoplasma gondii*. *Traffic* 12:287–300
111. Prole DL, Taylor CW. 2011. Identification of intracellular and plasma membrane calcium channel homologues in pathogenic parasites. *PLoS ONE* 6:e26218
112. Reiss M, Viebig N, Brecht S, Fourmaux MN, Soete M, et al. 2001. Identification and characterization of an escorter for two secretory adhesins in *Toxoplasma gondii*. *J. Cell Biol.* 152:563–78
113. Roiko MS, Carruthers VB. 2009. New roles for perforins and proteases in apicomplexan egress. *Cell Microbiol.* 11:1444–52
114. Roiko MS, Svezhova N, Carruthers VB. 2014. Acidification activates *Toxoplasma gondii* motility and egress by enhancing protein secretion and cytolytic activity. *PLoS Pathog.* 10:e1004488

115. Rutaganira FU, Barks J, Dhason MS, Wang Q, Lopez MS, et al. 2017. Inhibition of calcium dependent protein kinase 1 (CDPK1) by pyrazolopyrimidine analogs decreases establishment and reoccurrence of central nervous system disease by *Toxoplasma gondii*. *J. Med. Chem.* 60:9976–89
116. Saha S, Coleman BI, Dubey R, Blader IJ, Gubbels MJ. 2017. Two phosphoglucomutase paralogs facilitate ionophore-triggered secretion of the *Toxoplasma* micronemes. *mSphere* 2:e00521-17
117. Salazar E, Bank EM, Ramsey N, Hess KC, Deitsch KW, et al. 2012. Characterization of *Plasmodium falciparum* adenyllyl cyclase-beta and its role in erythrocytic stage parasites. *PLOS ONE* 7:e39769
118. Santos JM, Lebrun M, Daher W, Soldati D, Dubremetz JF. 2009. Apicomplexan cytoskeleton and motors: key regulators in morphogenesis, cell division, transport and motility. *Int. J. Parasitol.* 39:153–62
119. Schwab JC, Beckers CJ, Joiner KA. 1994. The parasitophorous vacuole membrane surrounding intracellular *Toxoplasma gondii* functions as a molecular sieve. *PNAS* 91:509–13
120. Sidik SM, Hortua Triana MA, Paul AS, El Bakkouri M, Hackett CG, et al. 2016. Using a genetically encoded sensor to identify inhibitors of *Toxoplasma gondii* Ca<sup>2+</sup> signaling. *J. Biol. Chem.* 291:9566–80
121. Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, et al. 2016. A genome-wide CRISPR screen in *Toxoplasma* identifies essential apicomplexan genes. *Cell* 166:1423–35.e12
122. Singh S, Alam MM, Pal-Bhowmick I, Brzostowski JA, Chitnis CE. 2010. Distinct external signals trigger sequential release of apical organelles during erythrocyte invasion by malaria parasites. *PLOS Pathog.* 6:e1000746
123. Stommel EW, Ely KH, Schwartzman JD, Kasper LH. 1997. *Toxoplasma gondii*: dithiol-induced Ca<sup>2+</sup> flux causes egress of parasites from the parasitophorous vacuole. *Exp. Parasitol.* 87:88–97
124. Sudhof TC, Rizo J. 2011. Synaptic vesicle exocytosis. *Cold Spring Harb. Perspect. Biol.* 3:a005637
125. Sugi T, Ma YF, Tomita T, Murakoshi F, Eaton MS, et al. 2016. *Toxoplasma gondii* cyclic AMP-dependent protein kinase subunit 3 is involved in the switch from tachyzoite to bradyzoite development. *mBio* 7:e00755-16
126. Taylor SS, Ilouz R, Zhang P, Kornev AP. 2012. Assembly of allosteric macromolecular switches: lessons from PKA. *Nat. Rev. Mol. Cell Biol.* 13:646–58
127. Tosetti N, Dos Santos Pacheco N, Soldati-Favre D, Jacot D. 2019. Three F-actin assembly centers regulate organelle inheritance, cell-cell communication and motility in *Toxoplasma gondii*. *eLife* 8:e42669
128. Treeck M, Sanders JL, Gaji RY, LaFavers KA, Child MA, et al. 2014. The calcium-dependent protein kinase 3 of *Toxoplasma* influences basal calcium levels and functions beyond egress as revealed by quantitative phosphoproteome analysis. *PLOS Pathog.* 10:e1004197
129. Uboldi AD, Wilde ML, McRae EA, Stewart RJ, Dagley LF, et al. 2018. Protein kinase A negatively regulates Ca<sup>2+</sup> signalling in *Toxoplasma gondii*. *PLOS Biol.* 16:e2005642
130. Vaandrager AB, Ehlert EM, Jarchau T, Lohmann SM, de Jonge HR. 1996. N-terminal myristoylation is required for membrane localization of cGMP-dependent protein kinase type II. *J. Biol. Chem.* 271:7025–29
131. Vieira MC, Moreno SN. 2000. Mobilization of intracellular calcium upon attachment of *Toxoplasma gondii* tachyzoites to human fibroblasts is required for invasion. *Mol. Biochem. Parasitol.* 106:157–62
132. Wallbank BA, Dominicus CS, Broncel M, Legrave N, Kelly G, et al. 2019. Characterisation of the *Toxoplasma gondii* tyrosine transporter and its phosphorylation by the calcium-dependent protein kinase 3. *Mol. Microbiol.* 111:1167–81
133. Wang S, Li Y, Ma C. 2016. Synaptotagmin-1 C2B domain interacts simultaneously with SNAREs and membranes to promote membrane fusion. *eLife* 5:e14211
134. Wentzinger L, Bopp S, Tenor H, Klar J, Brun R, et al. 2008. Cyclic nucleotide-specific phosphodiesterases of *Plasmodium falciparum*: PfPDE $\alpha$ , a non-essential cGMP-specific PDE that is an integral membrane protein. *Int. J. Parasitol.* 38:1625–37
135. Wetzel DM, Chen LA, Ruiz FA, Moreno SN, Sibley LD. 2004. Calcium-mediated protein secretion potentiates motility in *Toxoplasma gondii*. *Cell Sci.* 117(Part 24):5739–48
136. Wiersma HI, Galuska SE, Tomley FM, Sibley LD, Liberator PA, Donald RG. 2004. A role for coccidian cGMP-dependent protein kinase in motility and invasion. *Int. J. Parasitol.* 34:369–80
137. Williams MJ, Alonso H, Enciso M, Egarter S, Sheiner L, et al. 2015. Two essential light chains regulate the MyoA lever arm to promote *Toxoplasma* gliding motility. *mBio* 6:e00845-15

138. Wong SY, Remington JS. 1994. Toxoplasmosis in pregnancy. *Clin. Infect. Dis.* 18:853–61
139. Yan X, Ji Y, Liu X, Suo X. 2015. Nitric oxide stimulates early egress of *Toxoplasma gondii* tachyzoites from human foreskin fibroblast cells. *Parasites Vectors* 8:420
140. Yang L, Uboldi AD, Seizova S, Wilde ML, Coffey MJ, et al. 2019. An apically located hybrid guanylate cyclase-ATPase is critical for the initiation of Ca<sup>2+</sup> signalling and motility in *Toxoplasma gondii*. *J. Biol. Chem.* 294:8959–72
141. Yao J, Gaffaney JD, Kwon SE, Chapman ER. 2011. Doc2 is a Ca<sup>2+</sup> sensor required for asynchronous neurotransmitter release. *Cell* 147:666–77
142. Zeniou-Meyer M, Zabari N, Ashery U, Chasserot-Golaz S, Haerberle AM, et al. 2007. Phospholipase D1 production of phosphatidic acid at the plasma membrane promotes exocytosis of large dense-core granules at a late stage. *J. Biol. Chem.* 282:21746–57
143. Zhang X, Zhang H, Fu Y, Liu J, Liu Q. 2018. Effects of estradiol and progesterone-induced intracellular calcium fluxes on *Toxoplasma gondii* gliding, microneme secretion, and egress. *Front. Microbiol.* 9:1266

# Contents

The Way It Was <i>M. J. Osborn</i> .....	1
Life Cycle of <i>Cryptococcus neoformans</i> <i>Youbao Zhao, Jianfeng Lin, Yumeng Fan, and Xiaorong Lin</i> .....	17
Posttranscription Initiation Control of Gene Expression Mediated by Bacterial RNA-Binding Proteins <i>Paul Babitzke, Ying-Jung Lai, Andrew J. Renda, and Tony Romeo</i> .....	43
Ecology and Evolution of Plant Microbiomes <i>Viviane Cordovez, Francisco Dini-Andreote, Víctor J. Carrión, and Jos M. Raaijmakers</i> .....	69
Cellular Functions and Mechanisms of Action of Small Heat Shock Proteins <i>Axel Mogk, Carmen Ruger-Herreros, and Bernd Bukau</i> .....	89
Protein Acetylation in Bacteria <i>Chelsey M. VanDrisse and Jorge C. Escalante-Semerena</i> .....	111
Coordination of the Cell Cycle in Trypanosomes <i>Richard J. Wheeler, Keith Gull, and Jack D. Sunter</i> .....	133
Phage Therapy in the Twenty-First Century: Facing the Decline of the Antibiotic Era; Is It Finally Time for the Age of the Phage? <i>Shayla Hesse and Sankar Adhya</i> .....	155
Structural Basis of Response Regulator Function <i>Rong Gao, Sophie Bouillet, and Ann M. Stock</i> .....	175
Multikinase Networks: Two-Component Signaling Networks Integrating Multiple Stimuli <i>Vanessa I. Francis and Steven L. Porter</i> .....	199
Functional Regulators of Bacterial Flagella <i>Sundharraman Subramanian and Daniel B. Kearns</i> .....	225
Tc Toxin Complexes: Assembly, Membrane Permeation, and Protein Translocation <i>Daniel Roderer and Stefan Raunser</i> .....	247

Algal Sex Determination and the Evolution of Anisogamy <i>James Umen and Susana Coelho</i> .....	267
The Ultimate Guide to Bacterial Swarming: An Experimental Model to Study the Evolution of Cooperative Behavior <i>Jinyuan Yan, Hilary Monaco, and Joao B. Xavier</i> .....	293
Responses of Microorganisms to Osmotic Stress <i>Erhard Bremer and Reinhard Krämer</i> .....	313
Biogeography of the Oral Microbiome: The Site-Specialist Hypothesis <i>Jessica L. Mark Welch, Floyd E. Dewhirst, and Gary G. Borisy</i> .....	335
Bacterial Persisters and Infection: Past, Present, and Progressing <i>Bridget Gollan, Grzegorz Grabe, Charlotte Michaux, and Sophie Helaine</i> .....	359
Multiple Roles of c-di-GMP Signaling in Bacterial Pathogenesis <i>Martina Valentini and Alain Filloux</i> .....	387
Chromatic Acclimation in Cyanobacteria: A Diverse and Widespread Process for Optimizing Photosynthesis <i>Joseph E. Sanfilippo, Laurence Garczarek, Frédéric Partensky, and David M. Keboe</i> .....	407
Diversity, Genomics, and Distribution of Phytoplankton-Cyanobacterium Single-Cell Symbiotic Associations <i>Rachel A. Foster and Jonathan P. Zebr</i> .....	435
Bent Bacteria: A Comparison of Cell Shape Mechanisms in <i>Proteobacteria</i> <i>Jennifer A. Taylor, Sophie R. Sichel, and Nina R. Salama</i> .....	457
<i>Acinetobacter baumannii</i> : Envelope Determinants That Control Drug Resistance, Virulence, and Surface Variability <i>Edward Geisinger, Wenwen Huo, Juan Hernandez-Bird, and Ralph R. Isberg</i> .....	481
Two-Component Sensing and Regulation: How Do Histidine Kinases Talk with Response Regulators at the Molecular Level? <i>Alejandro Buschiazso and Felipe Trajtenberg</i> .....	507
Human Coronavirus: Host-Pathogen Interaction <i>To Sing Fung and Ding Xiang Liu</i> .....	529
Small Is Mighty—Chemical Communication Systems in <i>Pseudomonas aeruginosa</i> <i>Stephen Dela Abator and LianHui Zhang</i> .....	559
Signaling Cascades Governing Entry into and Exit from Host Cells by <i>Toxoplasma gondii</i> <i>Hugo Bisio and Dominique Soldati-Favre</i> .....	579

Metabolic Basis of Pathogenesis and Host Adaptation in Rice Blast <i>Yi Zhen Deng and Naweed I. Naqvi</i> .....	601
Assembly and Subcellular Localization of Bacterial Type VI Secretion Systems <i>Jing Wang, Maj Brodmann, and Marek Basler</i> .....	621
Paleomicrobiology: Diagnosis and Evolution of Ancient Pathogens <i>Kirsten I. Bos, Denise Kühnert, Alexander Herbig, Luis Roger Esquivel-Gomez, Aida Andrades Valtueña, Rodrigo Barquera, Karen Giffin, Aditya Kumar Lankapalli, Elizabeth A. Nelson, Susanna Sabin, Maria A. Spyrou, and Johannes Krause</i> .....	639
Defense and Counterdefense During Plant-Pathogenic Oomycete Infection <i>Yan Wang, Brett M. Tyler, and Yuanchao Wang</i> .....	667

## Indexes

Cumulative Index of Contributing Authors, Volumes 69–73 .....	697
---	-----

## Errata

An online log of corrections to *Annual Review of Microbiology* articles may be found at <http://www.annualreviews.org/errata/micro>